

DEPARTMENT OF WATER AND SANITATION


NO. 2428

26 August 2022

**NATIONAL WATER ACT, 1998
(ACT NO. 36 OF 1998)****RESERVE DETERMINATION FOR WATER RESOURCES OF THE BREEDE-GOURITZ WATER
MANAGEMENT AREA**

I, Senzo Mchunu, in my capacity as Minister of Water and Sanitation, and duly authorised in terms of sections 16(1) of the National Water Act, 1998 (Act No. 36 of 1998), hereby publish the Reserve determination for water resources of the Breede-Gouritz Water Management Area, as set out in the Schedule to this Notice.

Director: Reserve Determination
Attention: Mr Yakeen Atwaru
Department of Water and Sanitation
Ndinaye Building 185 Francis Baard Street
Private Bag X313
Pretoria
0001
Email: atwaruy@dws.gov.za



SENZO MCHUNU (MP)
MINISTER OF WATER AND SANITATION
DATE: 13/07/22

**RESERVE DETERMINATION FOR WATER RESOURCES OF THE BREEDE-GOURITZ WATER
MANAGEMENT AREA IN TERMS OF SECTION 16(1) AND (2) OF THE NATIONAL WATER ACT,
1998 (ACT NO. 36 OF 1998)**

SCHEDULE

1. DESCRIPTION OF WATER RESOURCE

- 1.1. The Reserve is determined for all or part of every significant water resource within the Breede-Gouritz Water Management Area as set out below:

Water Management Area:	Breede-Gouritz
Drainage Regions:	G40-G50, H10- H90, J11-J40, K10-K70 Tertiary Drainage Region
Rivers:	Breede Overberg Area: Breede River, Rivieronderend River, Overberg River, as well as other smaller coastal rivers. Gouritz Coastal Area: Gouritz River, Buffels River, Touws River, Groot River, Gamka River, Olifants River, Kammanassie River, and smaller coastal rivers.

- 1.2. The Minister has in terms of section 12 of the National Water Act, 1998 (Act No.36 of 1998) ("the Act"), prescribed a system for classifying water resources by issuing Government Notice No. R. 810, published in *Government Gazette* No. 33541 dated 17 September 2010. In terms of section 16(1) of the Act, the Minister must, as soon as reasonably practicable after the class of all or part of a water resource has been determined, by Notice in the *Gazette*, determine the Reserve for all or part of that water resource.

- 1.3. The Minister, in terms of section 16(1) of the Act, determines the Reserve for the Breede-Gouritz Water Management Area.

2. PROPOSED RESERVE DETERMINATION AS REQUIRED IN TERMS OF SECTION 16(1) AND (2) OF THE NATIONAL WATER ACT, 1998

- 2.1. The proposed Reserve which includes the Ecological Water Requirements (EWRs) and the Basic Human Needs Reserve (BHN) for the Rivers at EWR sites and selected biophysical nodes in the Breede-Gouritz Water Management Area is set out in **Section 4**. The Gouritz and Breede-Overberg Catchments and EWR sites are indicated in **Figure 1 and 2 respectively**.
- 2.2. The Water Quality component of the proposed Reserve for the Rivers at the EWR sites in Breede-Gouritz Water Management Area in terms of section 16(1) of the Act is set out in **Section 5**.
- 2.3. The proposed Groundwater Reserve for Water Quantity and Quality in terms of section 16(1) of the Act for the Breede-Gouritz Water Management Area is set out in **Section 6**.
- 2.4. The proposed Estuarine Reserve in terms of section 16(1) of the Act for the Breede-Gouritz Water Management Area is set out in **Section 7**.
- 2.5. The proposed Wetland Reserve in terms of section 16(1) of the Act for the Breede-Gouritz Water Management Area catchment is set out in **Section 8**.
- 2.6. The Reserve will apply from the date signed off as determined in terms of section 16(1) of the Act, unless otherwise specified by the Minister.

3. ACRONYMS AND DEFINITIONS

3.1 Acronyms

BHN	Basic Human Needs
EC	Ecological Category
EcoSpecs	Ecological Specifications
EIS	Ecological Importance and Sensitivity
ER	Ecological Reserve
EWR	Ecological Water Requirement
MAR	Mean Annual Runoff
MCM	Million Cubic Metres
PES	Present Ecological Status
RC	Reference conditions
REC	Recommended Ecological Category
TPCs	Thresholds of Potential Concern
WQSU	Water quality sub-unit

3.2 Definitions

Baseflow is a sustained low flow in rivers during dry or fair-weather conditions, but not necessarily all contributed by groundwater; includes contribution from delayed interflow and groundwater discharge.

Ecological Importance and Sensitivity (EIS): Key indicators in the ecological classification of water resources. Ecological importance relates to the presence, representativeness and diversity of species of biota and habitat. Ecological sensitivity relates to the vulnerability of the habitat and biota to modifications that may occur in flows, water levels and physico-chemical conditions.

Ecological Water Requirements (EWR): The flow patterns (magnitude, timing and duration) and water quality needed to maintain a riverine ecosystem in a particular condition. This term is used to refer to both the quantity and quality components.

Ecological Water Requirement (EWR) Sites: Specific points on the river as determined through the site selection process. An EWR site consists of a length of river which may consist of various cross-sections for both hydraulic and ecological purposes. These sites provide sufficient indicators to assess environmental flows and assess the condition of biophysical components (drivers such as hydrology, geomorphology and physico-chemical) and biological responses (viz. fish, invertebrates and riparian vegetation).

Present Ecological State (PES): A category indicating the current health or integrity of various biological attributes of the water resource, compared to the natural or close to natural reference conditions. The results of the process are provided as Ecological Categories (ECs) ranging from A (near natural) to F (completely modified) for the PES.

Recharge is the addition of water to the zone of saturation, either by downward percolation of precipitation or surface water and/ or the lateral migration of groundwater from adjacent aquifers.

Recommended Ecological Category (REC): An ecological category indicating the ecological management target for a water resource based on its ecological classification that should be attained. Categories range from Category A (unmodified, natural) to Category D (largely modified).

Reserve: is the quantity and quality of the water required to satisfy the basic human needs by securing a basic water supply and to protect the aquatic ecosystem in order to secure ecologically sustainable development and use of the relevant water resource.

River Node (biophysical node): These are modelling point's representative of an upstream reach or area of an aquatic eco-system (rivers, wetlands, estuaries and groundwater) for which a suite of relationships apply.

Sub-quaternary catchments: A finer subdivision of the quaternary catchments (the catchment areas of tributaries of main stem rivers in quaternary catchments).

4. SURFACE WATER QUANTITY COMPONENT FOR RIVERS AT SELECTED EWR SITES AND NODES

The Reserve consists of two parts – the Basic Human Need (BHN) Reserve and the Ecological Reserve (ER). The BHN Reserve provides for the essential needs of individuals served by the water resource in question and includes water for drinking, food preparation and for personal hygiene. The ER relates to the water required to protect the aquatic ecosystems of the water resource. The Reserve refers to both the **quantity** and **quality** of the water in the resource and will vary depending on the class of the resource (Class I, II and III).

BREEDE-OVERBERG AREA (ECOLOGICAL RESERVE)

Table 4.1: Summary of the data for Nodes and EWR sites. EWR Sites are indicated in bold.

Quaternary Catchment	Node/ EWR site	Water Resource	PES	EIS	REC	nMAR (MCM)	EWR nMAR (%)
G40C	Piii1	Palmiet	B	High	B	39.9	19.1
G40C	Piv10	Witklippiesskl oof	D	High	D	15.1	21.5
G40C	Piv9	Palmiet	D	High	B	78.8	21.5
G40C	Piv8	Klipdrift	D	High	D	13.6	21.5
G40D	Piv4	Klein-Palmiet	D	High	D	13.7	21.5
G40D	Piv7	Krom/Ribbok	D	Very High	A	27.5	21.5
G40D	Piii2	Palmiet	C	Very High	B/C	206.6	31.2
G40D	Piv12	Dwars/Louws	C	Very High	C	25.2	100.0
G40D	Piii3	Palmiet	C	Very High	B	250.4	34.5
G40G	Niii5	Bot	C	Very High	A	31.9	21.3
G40H	Nx6	Onrus	E	High	B	5.1	13.4
G40F	Niv43	Swart	E	High	B	42.1	13.3
G40K	Niv45	Steenbok	E	Very High	A	10.8	12.2
G40J	Nii4	Hartebees	D	Very High	B	18.4	12.5
G40K	Nv23	Klein	D	Moderate	C	43.0	19.3
G40M	Nx8	Uilkraal	C	Very High	A	2.4	19.2
G50B	Ni4	Nuwejaar	D	Moderate	D	12.5	13.0
G50C	Nvii15	Heuninges	D	-	D	17.8	13.1
G50C	Niv44	Heuninges	D	Very High	B	18.8	13.1
G50C	Nii5	Kars	E	Very High	B	21.6	20.4
G50E	Nv24	Kars	C	Moderate	B	15.4	30.3
G50H	Nii7	DeHoop Vlei	B	High	B	27.1	30.0
G50H	Nii6	Sout	D	-	B	4.2	12.6
H10B	Nvii3	Rooikloof	B	High	B	6.807	37.95
H10B	Niv3	Titus	C	Moderate	C	26.2	22.0
H10C	Niv1	Koekedou	D	Very High	A	18.8	14.2
H10C	Niv2	Dwars	C	High	B	74.9	22.0
H10D	Nvi4	Breede	D	Moderate	D	175.509	17.51
H10D	Niv4	Witels	A	Very High	A	84.3	43.3
H10D	Nvi3	Breede	C	High	B	252.8	31.7
H10E	Nvi2	Wit	A	Very High	A	42.6	46.6
H10F	Niv6	Wabooms	D	High	B	7.4	14.4
H10F	Nviii1/ EWR1	Breede	D/E	High	D	434.90	31.7
H10G	Niv7	Slanghoek	D	High	B	32.6	14.5
H10G	Niii1	Breede	D	High	B	497.6	25.4
H10J	Niv40	Elands	B	Very High	A	58.1	50.8
H10J	Niv41	Krom	B	Very High	A	9.0	50.8
H10J	Nvii2/ EWR2	Molenaars	C	Very High	B	105.6	35.0
H10J	Niv42	Molenaars (Smalblaar)	E	High	B	191.2	17.4

Quaternary Catchment	Node/ EWR site	Water Resource	PES	EIS	REC	nMAR (MCM)	EWR nMAR (%)
H10K	Niv12	Holsloot	C	High	B	119.5	35.0
H10H	Nvii6	Hartbees	D	Very High	A	4.0	14.4
H10H	Niv9	Hartbees/ Wetskloof	D	Very High	A	10.2	14.4
H10L	Nv3	Breede	C	High	B	850.9	31.7
H20G	Nvii7	Hex	C	Moderate	C	102.8	22.3
H20H	Niv10	Hex	D	High	B	107.1	22.3
H40B	Nvii5	Koo	D	High	B	0.9	13.1
H40C	Niv11	Nuy	E	High	B	29.4	13.2
H40D	Niv13	Doring	E	High	B	47.4	12.9
H40F	Nvii8/ EWR3	Breede	C/D	Moderate	C/D	1042.8	45.5
H40G	Nvii11	Poesnels	D	High	B	16.1	12.8
H40K	Niv14	Keisers	D	Very High	A	12.6	12.5
H40H	Niv15	Vink	D	Very High	A	15.6	12.4
H30C	Niv20	Pietersfontein	D	Moderate	C	17.3	12.0
H30B	Niv18	Kingna	D	High	B	27.1	12.3
H30D	Nvii9	Keisie	D	High	B	21.5	11.9
H30E	Nii2	Kogmanskloof	D	Very High	B	52.0	18.9
H50B	Ni2	Breede	D	High	B	1170.1	17.3
H60B	Nvii10	Du Toits	B	Very High	A	43.9	50.8
H60D	Nv7	Riviersonderend	C	Very High	A	370.2	30.1
H60E	Niv28/ EWR6	Baviaans	B	High	B	7.9	70.90
H60E	Niv29	Sersants	D	High	B	4.6	29.9
H60F	Niv30	Gobos	C	Very High	A	12.4	48.1
H60F	Nv9/EWR5	Riviersonderend	D	High	D	413.7	24.5
H60G	Niv31	Kwartel	D	High	B	10.7	13.4
H60H	Niv33	Soetmelksvlei	D	Very High	A	4.0	29.9
H60H	Niv34	Slang	D	Very High	A	2.1	29.9
H60H	Nv10	Riviersonderend	D	Very High	A	442.9	24.5
H60K	Niv35	Kwassadie	E	Very High	A	5.9	17.3
H60L	Ni3	Riviersonderend	D	High	B	483.8	24.5
H70A	Niv24	Leeu	E	Very High	A	5.8	12.6
H70B	Nv2	Breede	C	High	B	1701.4	26.4
H70D	Nii3	Tradouw	B	Very High	A	19.4	29.9
H70F	Niv25	Buffeljags	E	High	B	119.4	14.1
H70G	Niii4/ EWR4	Breede	C	Very High	B/C	1832.7	40.1
H70J	Niv26	Slang	E	High	B	10.0	14.2

GOURITZ COASTAL AREA (ECOLOGICAL RESERVE)

Table 4.2: Summary of the data for Nodes and EWR sites. EWR sites are indicated in bold.

Quaternary Catchment	Node/ site	EWR	Water Resource	PES	EIS	REC	nMAR (MCM)	EWR nMAR (%)
J11C	giv34		Buffels	B	High	B	13.1	26.5
J11F	gv25		Buffels	B	High	B	24.2	17.8
J11H	J1BUFF- EWR5		Buffels	C	Moderate	C	27.4	17.9
J11K	giv32		Groot	D	High	B	30.5	17.9
J12D	giv28		Touws	D	High	B	16.4	11.3
J12H	giv27		Touws	B	Moderate	C	26.4	26.8
J12K	giv26		Brak	C	High	B	2.9	17.7
J12L	J1DORI- EWR7		Doring	C/D	Low	C/D	2.9	12.0
J12L	J12L Modelled		Huis	D		D	1.56	40.3
J12M	J1TOUW- EWR3		Touws	B/C	High	B/C	33.5	17.8
J13B	gv7		Groot	C	High	B	72.7	18.0
J13C	gii3		Groot	B	High	B	78.1	27.0
J21D	giv3		Gamka	B	High	A	31.9	27.1
J22F	giv1		Koekemoers	C	Very High	A	7.4	17.9
J22K	giv2		Leeu	C	Very High	A	17.1	17.9
J23F	gv17		Gamka	B	High	B	58.1	27.0
J23J	gv27		Gamka	C	High	B	69.6	18.3
J24E	gv14		Dwyka	A	High	B	4.0	39.1
J25A	J2GAMK- EWR4		Gamka	C/D	High	C	79.8	14.9
J25E	gii2		Gamka	C	High	B	111.8	15.2
J31D	J3OLIF- EWR9		Olifants	C	Moderate	C	11.8	17.8
J32E	giv15		Traka	C	High	C	2.7	17.9
J33B	gv33		Olifants	D	High	B	25.0	11.9
J33E	gv21		Meirings	C	Very High	A	21.4	19.1
J33F	giv11		Olifants	E	High	B	80.0	12.4
J34C	J3KAMM- EWR10		Kammanassie	C/D	Low	C/D	41.2	15.3
J34F	giv10		Leeu	E	Very High	A	59.2	12.1
J35E	gv19		Olifants	E	High	B	224.5	12.9
J35F	giv17		Olifants	D	High	B	253.4	12.9
J40B	J4GOUR- EWR6		Gouritz	C	Moderate	C	489.1	14.8
J40E	gv9		Gouritz	C	High	B	571.8	14.8
H80B	giii5		Duiwenhoks	E	Very High	A	62.5	20.1
H80E	H8DUIW- EWR1		Duiwenhoks	D	Low	D	83.2	20.9
H90C	giv27		Korinte	D	High	B	34.1	14.5
H90C	H9GOUK- EWR2		Goukou	C/D	Moderate	C/D	50.9	24.2
H90E	gv41		Goukou	C	High	B	105.0	28.2
K10D	giv25		Brandwag	D	High	B	17.9	9.9
K20A	gvii7		Groot-Brak	B/C	Very High	A	27.0	26.5
K20A	gviii2- GB1-BC	EWR	Groot-Brak	B/C	Very High	B/C	15.3	26.5
K20A	gvii3-EWR Var 3		Varing	D	High	C/D	8.4	20.9
K20A	gviii12- EWR Var2		Varing	D	High	C/D	6.0	20.9
K30A	gviii4-EWR		Maalgate	B	Very High	A	15.3	46.0

Quaternary Catchment	Node/ EWR site	Water Resource	PES	EIS	REC	nMAR (MCM)	EWR nMAR (%)
K30A	gvii8	Maalgate	B	High	D	30.1	16.4
K30B	gvii9	Malgas	C	Very High	C	17.3	31.6
K30B	gviii6 EWR Gwa1 -D	Gwaing	E	High	D	34.1	16.4
K30C	gviii7 EWR Sw1 - D	Swart	D	High	D	16.1	14.5
K30C	gvii11 EWR Ka1 - D	Kaaimans	B	High	B	18.6	50.2
K30C	gviii8 EWR Si1 -B	Silver	B	Very High	B	14.9	50.2
K30D	gvii12	Touws	B	Very High	A	16.7	30.3
K30D	gx8	Klein Keurbooms	D	Very High	B	2.5	14.1
K40A	giii10 EWR 2 Diep -B	Diep	B	Very High	B	12.4	30.3
K40B	giii13	Hoekraal	B	Very High	A	27.9	30.3
K40C	gvii13 EWR 4 Karatara-AB	Karatara	B	Very High	A/B	11.2	40.2
K40C	giii11	Karatara	A/B	Very High	A	33.8	40,2
K40E	Gou 1	Goukamma	B/C	Very High	B/C	30.4	38.5
K50A	EWR 1	Knysna	B	High	B	26.5	32.1
K50A	Kny 2	Knysna	B	-	B	46.5	32.1
K50B	EWR 2	Gouna	A/B	Very High	A/B	27.6	53.4
K60C	K6KEUR-EWR8	Keurbooms	C	Very High	B/C	46.1	34.9
K60D	giv5	Palmiet	A	Very High	A	42.1	48.3
K60F	giv4	Bitou	C	Very High	A	23.6	22.8
K60G	Noe 1	Noetsie	B	Very High	A/B	4.8	63.4
K60G	gx3	Piesang	D	Very High	A	7.3	28.5
K60E	gx9	Keurbooms	C	Very High	A	91.3	34.9
K70A	gx4	Buffels	B	Very High	B	1.8	34.3
K70A	gx5	Sout	B	Very High	B	3.8	34.3
K70B	gvii15	Bloukrans	B	Very High	B	31.2	33.9

Table 4.3: Basic Human Needs for the Breede-Gouritz WMA

Quaternary Catchment	Water Resource	BHN (%NMAR)	Quaternary Catchment	Water Resource	BHN (%NMAR)
G40C	Palmiet	0.008	J11C	Buffels	0.02
G40D	Palmiet	0	J11F	Buffels	0.03
G40G	Bot	0.50	J11H	Buffels	0.03
G40H	Onrus	5.88	J11K	Groot	0.36
G40F	Swart	0.17	J12D	Touws	0.03
G40K	Steenbok	0	J12H	Touws	0
G40J	Hartebees	0.08	J12K	Brak	0
G40K	Klein	0	J12L	Doring	0.21
G40M	Uilkraal	0.125	J12M	Touws	0
G50B	Nuwejaar	1.12	J13B	Groot	0.01
G50C	Heuninges	0	J13C	Groot	0
G50E	Kars	0.84	J21D	Gamka	0
G50G	DeHoop Vlei	0.03	J22F	Koekemoers	0.14
G50H	Sout	0.04	J22K	Leeu	0
H10B	Rooikloof	0	J23F	Gamka	0.10
H10C	Dwars	0.70	J23J	Gamka	0
H10D	Breede	0	J24E	Dwyka	0
H10E	Wit	0	J25A	Gamka	0.003
H10F	Breede	0.04	J25E	Gamka	0.03
H10G	Slanghoek	0	J31D	Olifants	0.02
H10J	Elands	0.02	J32E	Traka	0.03
H10K	Holsloot	0	J33B	Olifants	0.02
H10H	Breede	0	J33E	Meirings	0.56
H10L	Breede	0.005	J34C	Olifants	0.01
H20G	Hex	0.01	J34F	Kammanassie	0.05
H20H	Hex	1.20	J35E	Leeu	0.004
H40B	Koo	2.22	J35F	Olifants	0.01
H40C	Nuy	0.07	J40B	Olifants	0
H40D	Doring	0	J40E	Gouritz	0.003
H40F	Breede	0.002	H80B	Gouritz	0
H40G	Poesnels	0	H80E	Duiwenhoks	0.005
H40K	Keisers	0.32	H90C	Goukou	0.33
H40H	Vink	0.06	H90E	Goukou	0.06
H30C	Pietersfontein	0.07	K10D	Brandwag	0.06
H30B	Kingna	1.33	K20A	Varing	1.50
H30D	Keisie	0.04	K30A	Maalgate	0.39
H30E	Kogmanskloof	0.17	K30B	Malgas	0.18
H50B	Breede	0.004	K30C	Swart	8.99
H60B	Du Toits	0	K30D	Touws	0.54
H60D	Riviersonderend	0.001	K40A	Diep	0.04
H60E	Baviaans	1.14	K40B	Hoekraal	0
H60F	Gobos	0.007	K40C	Karatara	0.54
H60G	Kwartel	0	K40E	Goukamma	0.39
H60H	Soetmelksvlei	0	K50A	Knysna	0.002
H60K	Kwassadie	0	K50B	Gouna	0.47
H60L	Riviersonderend	0	K60C	Keurbooms	0.03
H70A	Leeu	0	K60D	Palmiet	25.26
H70B	Breede	0.01	K60F	Bitou	0.38
H70D	Tradouw	0	K60G	Piesang	4.34
H70F	Buffeljags	0.02	K60E	Keurbooms	0.02
H70G	Breede	0	K70A	Buffels	1.32
H70J	Slang	0.40	K70B	Bloukrans	0.03

5. SURFACE WATER QUALITY COMPONENT FOR RIVERS AT EWR SITES

BREDE-OVERBERG

Table 5.1: PES categories and overall site assessment for Breede River at Node Nviii1 (represented by EWR Site 1)

RIVER	Breede River		WATER QUALITY MONITORING POINTS	
WQRU	1 (Upper Breede River to Wit River confluence)	RC	DWA monitoring station @ Koekedou River in Ceres (H1H013Q01) (1998 -2002, n=38)	
EWR SITE	EWR Site 1	PES	DWA monitoring station @ Witbrug (H1H006Q01) (1998 -2002, n=143)	
Confidence assessment	Medium. EWR site is further downstream of DWA monitoring station.			
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L) 1:95 th Percentile values	MgSO ₄	7	21	Category B
	Na ₂ SO ₄	8	4	Category A
	MgCl ₂	5	6	Category A
	CaCl ₂	12	24	Category B
	NaCl	6	27	Category A
	CaSO ₄	0	0	Category A
Nutrients (mg/L)	SRP	-	-	No data
	TIN	2.8011	0.318	Category B
	TDS	21.8-50.6	< 45	<i>B Category.</i> Increased TDS during Winter months
	TSS	-	< 5	Acceptable
	P0 ₄	0.024	0.042	<i>C Category.</i>
Physical Variables	pH (5 th – 95 th %)	6.2-7.5	6.9 – 7.8	<i>A Category.</i>
	Temperature	-	-	No data but no concerns noted about DO concentrations
	Dissolved oxygen	-	-	
	Turbidity (NTU)	-	2	
	Electrical conductivity (mS/m)	-	-	No data
Response variables	Chl a: periphyton	-	-	No data
	Chl a: phytoplankton	-	-	No data
	Macroinvertebrates	SASS score = >110 and an ASPT score >7	SASS score = 69 and ASPT score = 5.3	<i>D/E Category.</i> Largely modified. Loss of habitat area through infilling.
	Fish community score			<i>D/E Category.</i> Introductions of alien fish species, i.e., bass, trout and blue gills.
Toxics		-	No data but pesticide residues are the concern due to intensive agriculture in Ceres	
OVERALL SITE CLASSIFICATION		<i>A/B Category</i>		

Table 5.2: PES categories and overall site assessment for Molenaars River at Node Nvii2 (represented by EWR Site 2)

RIVER	Molenaars River	WATER QUALITY MONITORING POINTS		
WQRU	2 (Complete Molenaars River)	RC	The Molenaars River @ Hawequas Forest Reserve (H1H018Q01) (1998 -1992, n=93)	
EWR SITE	EWR Site 2	PES	The Molenaars River @ Hawequas Forest Reserve (H1H018Q01) (1998 -2002, n=141)	
Confidence assessment	High. Monitoring point is very close to the EWR site. Historic data shows no trends and hence PES data was used for Reference conditions.			
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L) 1:95 th % values	MgSO ₄	6	6	Category A
	Na ₂ SO ₄	9	9	Category A
	MgCl ₂	4	4	Category A
	CaCl ₂	6	6	Category A
	NaCl	9	9	Category A
	CaSO ₄	0	0	Category A
Nutrients (mg/L)	SRP		-	
	TIN (1:50 th %)	0.151	0.151	Category A
	TDS	22	22	Category A
	TSS	< 5	< 5	Category A
	P ₀₄ (1:50 th %)	0.025	0.025	Category C
	NH ₃ -N	<0.015.	<0.015.	Within range
Physical Variables	pH (5 th – 95 th %)	5.5 – 7.3	5.5 – 7.3	A <i>Category</i> . Typical acidic Western Cape river
	Temperature			Within range, Category A
	Dissolved oxygen	80 - 120% saturation.	80 - 120% saturation.	
	Turbidity (NTU)	-	0	
	Electrical conductivity (mS/m)	-	-	No data
Response variables	Chl a: periphyton	-	-	No data
	Chl a: phytoplankton	-	-	No data
	Macroinvertebrates	SASS score ≥140 and an ASPT score >8	SASS score = 175 and ASPT score = 7.9	A/B <i>Category</i> . Catchment is relatively pristine
	Fish community score			E <i>Category</i> . No indigenous fish were recorded during the study
Toxics		-	-	No toxic substance concerns
OVERALL SITE CLASSIFICATION		A <i>Category</i>		

Table 5.3: PES categories and overall site assessment for Breede River at Node Nviii8 (represented by EWR Site 3)

RIVER	Breede River		WATER QUALITY MONITORING POINTS	
WQRU	3 (Middle Breede from Molenaars confluence to Kogmanskloof confluence)		RC	None. No WQ monitoring station could provide reference data for this Resource Unit.
EWR SITE	EWR Site 3		PES	Breede River at Le Chasseur (H4H017Q01) (1995 -1999, n=214)
Confidence assessment	Moderate			
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L) (1:95 th %)	MgSO ₄	-	39	Category D
	Na ₂ SO ₄	-	5	Category A
	MgCl ₂	-	12	Category A
	CaCl ₂	-	32	Category B
	NaCl	-	95	Category B
	CaSO ₄	-	0	Category A
Nutrients (mg/L)	SRP	-		
	TIN (1:50 th %)	-	0.242	Category A
	TDS	-	< 45	<i>B Category.</i> High TDS loads. Irrigation return flows in tributaries and main stem between Brandvlei Dam and EWR Site 3.
	TSS	-	< 5	High TSS loads. Releases from Brandvlei to alleviate high TDS.
	PO ₄ (1:50 th %)	-	0.032	C Category
Physical Variables	pH (5 th – 95 th %)	-	6.9 – 7.8	A Category
	Temperature (°C)	-	22.4	Summer temperature
	Dissolved oxygen	-	-	No observed data
	Turbidity (NTU)	-	-	No data. Low sediment production area
	Electrical conductivity (mS/m)	-	-	No data
Response variables	Chl a: periphyton	-	-	No data
	Chl a: phytoplankton	-	-	No data
	Macroinvertebrates	SASS score ≥110 and an ASPT score >7	SASS score = 91 and ASPT score = 7	<i>A Category.</i> Suggest there is no significant impact to community structure.
	Fish community score	-	-	<i>D Category.</i> Only 2 of the 7 indigenous freshwater species expected to occur there were sampled.
Toxics		-	-	No data but Pesticides are a concern as a result of intensive agriculture
OVERALL SITE CLASSIFICATION		<i>D Category.</i> Mainly due to increased summer low flows and increased sediment load.		

Table 5.4: PES categories and overall site assessment for Breede River at Node Niii4 (represented by EWR Site 4)

RIVER	Breede River		WATER QUALITY MONITORING POINTS	
WQRU	6 (Breede River from Buffelsjags River to Estuary)		RC	None. No WQ monitoring station could provide reference data for this Resource Unit.
EWR SITE	EWR Site 4		PES	Lower Breede River @ Swellendam (H7H006Q01) (1995 -1999, n=214)
Confidence assessment	Moderate (There were no observed hydrological data for this reach of the river and flow in the river during field visits was often too high to allow for discharge readings to be taken; Monitoring point quite far upstream)			
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L) (1:95 th %)	MgSO ₄	-	83	Category E/F
	Na ₂ SO ₄	-	3	Category A
	MgCl ₂	-	49	Category D
	CaCl ₂	-	62	Category C
	NaCl	-	318	Category D
	CaSO ₄	-	0	Category A
Nutrients (mg/L)	SRP	-	-	No data
	TIN (1:50 th %)	-	0.23	Category A
	TDS	-	-	C Category. High TDS loads due to Irrigation return flows in tributaries and main
	TSS	-	-	No data
	PO ₄ (1:50 th %)	-	0.024	C Category
	NH ₃ -N	-	0.3 (Median)	No historic data
Physical Variables	NO ₂ NO ₃ -N	-	<1	Recommended winter concentrations but summer can be as low as < 0.3 mg/l
	pH (5 th – 95 th %)	-	6.9 – 8.1	Category A/B. No significant change.
	Temperature (°C)	-	-	No observed data. Some concerns about short-term fluctuation in DO levels.
	Dissolved oxygen	-	-	
	Turbidity (NTU)	-	-	
Electrical conductivity (mS/m)	-	-	No data	
Response variables	Chl a: periphyton	-	-	No data
	Chl a: phytoplankton	-	-	No data
	Macroinvertebrates	SASS score ≥110 and an ASPT score >7	SASS score = 87 and ASPT score = 6.9	B Category. Moderately impacted.
	Fish community score	10 indigenous fish species		C Category. Presence of carp and bass.
Toxics		-		No data but pesticide residues are the concern due to intensive agriculture.
OVERALL SITE CLASSIFICATION		D Category. Dependent on high quality inflow from Buffelsjags River to ensure acceptable WQ in the lower Breede Resource Unit & Estuary		

Table 5.5: PES categories and overall site assessment for Riviersonderend at Node Nv7 (represented by EWR site 5)

RIVER	Riviersonderend River		WATER QUALITY MONITORING POINTS	
WQRU	7 (Middle Riviersonderend River from Theewaterskloof Dam to Bok River)		RC	Riviersonderend at Swart River/Nuweberg Forest Reserve (H6H008Q01) (1990 -1992, n=34)
EWR SITE	EWR Site 5		PES	Riviersonderend at Theewaterskloof Dam (H6H012Q01) (1998 -2002, n=39) Riviersonderend at Reenen (H6H009Q01) (1995 -1999, n=56)
Confidence assessment		High		
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L) 1:95 th value	MgSO ₄	7	12	Category A
	Na ₂ SO ₄	7	13	Category A
	MgCl ₂	4	14	Category A
	CaCl ₂	5	14	Category A
	NaCl	11	99	Category B
	CaSO ₄	0	0	Category A
Nutrients (mg/L)	SRP	-	-	No data
	TIN (1:50 th %)	0.068	0.154	Category A.
	TDS	13.6-32	Summer: <100 and winter: < 150	<i>B Category.</i> High TDS loads due to agricultural return flows in winter
	TSS	< 5	-	No data
	PO ₄ (1:50 th %)	<0.0165	0.013	Category B. Slightly higher.
Physical Variables	NO ₂ NO ₃ -N	0.02	<0.25	Recommended winter concentrations but summer can be as low as < 0.2 mg/l
	pH (5 th – 95 th %)	6.5 – 7.5	6.4 - 7.4	<i>A/B Category.</i>
	Temperature (°C)	+4 °C	-	No data
	Dissolved oxygen	80 - 120% saturation	-	
	Turbidity (NTU)	-	4	No data
Electrical conductivity (mS/m)	-	-		
Response variables	Chl a: periphyton	-	-	No data
	Chl a: phytoplankton	-	-	No data
	Macroinvertebrates	SASS score ≥110 and an ASPT score >7	SASS score = 95 and ASPT score = 6.8	<i>C/D Category.</i> Most of the reach has good water quality but poor habitat quality.
	Fish community score	Six indigenous fish species		<i>E Category.</i> Dominated by large and small mouth bass. Reduction of indigenous fish species.
Toxics		-	-	No data but pesticide residues are the concern due to intensive agriculture upstream of EWR site
OVERALL SITE CLASSIFICATION		<i>A/B Category</i> (The presence of Theewaterskloof Dam has slightly reduced the water quality in the downstream river)		

Table 5.6: PES categories and overall site assessment for Baviaans River at Node Niv28 (represented by EWR Site 6)

RIVER	Baviaans River	WATER QUALITY MONITORING POINTS		
WQRU	9 (Complete Baviaans River)	RC	Baviaans River at Genadendal Mission Station (H6H005Q01) (1972 -1994, n=346)	
EWR SITE	EWR Site 6	PES	Baviaans River at Genadendal Mission Station (H6H005Q01) (1998 -2002, n=42)	
Confidence assessment	High. Good data record to assess both reference and PES. Historic data shows no trends and hence PES data was used for Reference conditions			
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L) 1:95 th % value	MgSO ₄	9	9	Category A
	Na ₂ SO ₄	9	9	Category A
	MgCl ₂	4	4	Category A
	CaCl ₂	10	10	Category A
	NaCl	25	25	Category A
	CaSO ₄	0	0	Category A
Nutrients (mg/L)	SRP	-	-	No data
	TIN (1:50 th % value)	0.04	0.040	Category A
	TDS	41	41	Category A
	TSS	-	-	No data
	P0 ₄ (1:50 th % value)	0.022	0.022	Category A
Physical Variables	pH (5 th – 95 th %)	4.7 – 7.1	4.7 – 7.0	A Category. Regarded as natural for low pH Western Cape Rivers
	Temperature (°C)	-	-	No data but no DO concerns
	Dissolved oxygen	-	-	
	Turbidity (NTU)	-	-	
	Electrical conductivity (mS/m)	-	-	No data
Response variables	Chl a: periphyton	-	-	No data
	Chl a: phytoplankton	-	-	No data
	Macroinvertebrates	SASS score ≥140 and an ASPT score >8	SASS score = 109 and ASPT score = 8.38	A Category. Reduced habitat.
	Fish community score	Three indigenous fish species		A/B Category. All species expected to occur here historically, were recorded
Toxics		-	-	No data but no toxic substance concerns noted
OVERALL SITE CLASSIFICATION		A Category		

GOURITZ

Table 5.7: PES categories and overall site assessment for Duiwenhoks River at H8DUIW-EWR1

RIVER	Duiwenhoks River	WATER QUALITY MONITORING POINTS	
WQRU		RC	DWS gauging weir H8H001Q01 (1967 – 1979; number of samples (n) = 66 - 71, Electrical Conductivity: n = 110).
EWR SITE	H8DUIW-EWR1	PES	DWS gauging weir H8H001Q01 (2007 – 2013; n = 69, Fluorine (F) = 48).
Confidence assessment		Confidence: 3.5	
Water Quality Constituents		PES Value	Category/Comment
Inorganic salts (mg/L)	SO ₄	N/A	-
	Na	382.2	Exceeds the ≤ 70 mg/L (TWQR) for Agricultural Use: Irrigation.
	Mg	67.4	No guideline.
	Ca	55.0	No guideline.
	Cl	805.4	Exceeds the ≤ 100 mg/L (TWQR) for Agricultural Use: Irrigation.
	K	9.25	No guideline.
Nutrients (mg/L)	SRP	0.014	A
	TIN	0.118	A
Physical Variables	pH (5 th – 95 th %)	6.6 and 8.1	B
	Temperature (°C)	N/A	A/B. Impacts expected at low flows.
	Dissolved oxygen	N/A	B. Impacts expected at low flows.
	Turbidity (NTU)	N/A	B. Changes in turbidity appear to be largely related to natural with minor man-made modifications, e.g. gravel mining upstream
	Electrical conductivity (mS/m)	272	80 mS/m
Response variables	Chl a: phytoplankton	N/A	N/A
	Macroinvertebrates	50.7% SASS score = 78 ASPT score = 56	D
	Diatoms	11.1	C/D (n = 1, Jan 2014)
	Fish community score	51.6%	D (all estuarine spp. that moved into the freshwater zone and aliens).
Toxics	Ammonia (as N)	0.003	A
	Fluoride (as F)	0.33	A
OVERALL SITE CLASSIFICATION		C Category	

Table 5.8: Water quality EcoSpecs and TPCs (C category) for Duiwenhoks River at H8DUIW-EWR1

Metrics	EcoSpecs	TPCs
Inorganic salt ions		
Sulphate as SO ₄	N/A	N/A
Sodium as Na	The 95 th percentile of the data must be ≤ 380 mg/L.	The 95 th percentile of the data must be 300 - 380 mg/L.
Magnesium as Mg	The 95 th percentile of the data must be ≤ 67 mg/L.	The 95 th percentile of the data must be 53.5 - 67 mg/L.
Calcium as Ca	The 95 th percentile of the data must be ≤ 55 mg/L.	The 95 th percentile of the data must be 44 - 55 mg/L.
Chloride as Cl	The 95 th percentile of the data must be ≤ 800 mg/L.	The 95 th percentile of the data must be 640 - 800 mg/L.
Potassium as K	The 95 th percentile of the data must be ≤ 9 mg/L.	The 95 th percentile of the data must be 7 - 9 mg/L.
Physical Variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data must be ≤ 270 mS/m.	The 95 th percentile of the data must be 210 - 270 mS/m.
pH	The 5 th percentile of the data must be 6.5 - 8.0, and the 95 th percentile 8.0 - 8.8.	The 5 th percentile of the data is ≤ 6.3 and the 95 th percentile is ≥ 8.6.
Temperature ^(a)	Natural temperature range.	Initiate baseline monitoring for this variable.
Dissolved oxygen ^(a) (DO)	The 5 th percentile of the data must be ≥ 7.0 mg/L.	The 5 th percentile of the data must be 7.2 - 7.0 mg/L. Initiate baseline monitoring for this variable.
Turbidity ^(a)	Changes in turbidity are related to minor man-made modifications (e.g. gravel mining upstream). Some silting of habitats is expected.	Initiate baseline monitoring for this variable.
Nutrients		
TIN	The 50 th percentile of the data must be ≤ 0.25 mg/L.	The 50 th percentile of the data must be 0.2 - 0.25 mg/L.
PO ₄ -P	The 50 th percentile of the data must be ≤ 0.015 mg/L.	The 50 th percentile of the data must be 0.012 - 0.015 mg/L.
Response variables		
Chl-a phytoplankton	The 50 th percentile of the data must be < 15 µg/L.	The 50 th percentile of the data must be 12 - 15 µg/L.
Chl-a periphyton	The 50 th percentile of the data must be ≤ 12 mg/m ² .	The 50 th percentile of the data must be 10 - 12 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data must be ≤ 1.5 mg/L.	The 50 th percentile of the data must be 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data must be ≤ 0.015 mg/L.	The 50 th percentile of the data must be 0.012 - 0.015 mg/L.
Other toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

(a) N/A- No data were available for this assessment.

Table 5.9: PES categories and overall site assessment for Goukou River at H9GOUK-EWR2

RIVER	Goukou River		WATER QUALITY MONITORING POINTS
WQRU		RC	Reference Condition (RC) was represented by the A Category benchmark tables in DWAF (2008), as no other data were available to describe natural state.
EWR SITE	H9GOUK-EWR2	PES	DWS gauging weir H9H005Q01 (2007 – 2014; n = 63 - 71, F = 52).
Confidence assessment	Confidence: 3		
Water Quality Constituents		PES Value	Category/Comment
Inorganic salts (mg/L)	SO ₄	N/A	N/A
	Na	650.4	Exceeds the ≤ 70 mg/L (TWQR) for Agricultural Use: Irrigation.
	Mg	79.0	No guideline.
	Ca	57.1	No guideline.
	Cl	1081.3	Exceeds the ≤ 100 mg/L (TWQR) for Agricultural Use: Irrigation.
	K	20.4	No guideline.
Nutrients (mg/L)	SRP	0.085	D
	TIN	0.055	A
Physical Variables	pH (5 th – 95 th %)	6.6 and 8.35	B
	Temperature (°C)	N/A	A/B. Impacts expected at low flows.
	Dissolved oxygen	N/A	B. Impacts expected at low flows.
	Turbidity (NTU)	N/A	A/B. Changes in turbidity appear to be largely related to natural.
	Electrical conductivity (mS/m)	408.4	E/F
Response variables	Chl a: phytoplankton	N/A	N/A
	Macroinvertebrates	51.2% SASS score = 113 ASPT score = 6.6	D
	Diatoms	14.4 and 11.0	C/D (n = 2; Jan and July 2014)
	Fish community score	47.4%	D
Toxics	Ammonia (as N)	0.01	A
	Fluoride (as F)	0.59	A
OVERALL SITE CLASSIFICATION		C/D Category	

Table 5.10: Water quality EcoSpecs and TPCs (Category: C/D) for Goukou River at GOUK-EWR2

Metrics	EcoSpecs	TPCs
Inorganic salt ions		
Sulphate as SO ₄	N/A	N/A
Sodium as Na	The 95 th percentile of the data is between ≤ 650 mg/L.	The 95 th percentile of the data is between 520 - 650 mg/L.
Magnesium as Mg	The 95 th percentile of the data is between ≤ 80 mg/L.	The 95 th percentile of the data is between 64 - 80 mg/L.
Calcium as Ca	The 95 th percentile of the data is between ≤ 55 mg/L.	The 95 th percentile of the data is between 44 - 55 mg/L.
Chloride as Cl	The 95 th percentile of the data is between ≤ 1 000 mg/L.	The 95 th percentile of the data is between 800 - 1 000 mg/L.
Potassium as K	The 95 th percentile of the data is between ≤ 20 mg/L.	The 95 th percentile of the data is between 16 - 20 mg/L.
Physical variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data is between ≤ 400 mS/m.	The 95 th percentile of the data is between 320 - 400 mS/m.
pH	The 5 th percentile of the data must be 6.5 - 8.0, and the 95 th percentile 8.0 - 8.8.	The 5 th percentile of the data is ≤ 6.3 and the 95 th percentile is ≥ 8.6.
Temperature ^(a)	Natural temperature range.	Initiate baseline monitoring for this variable.
Dissolved oxygen ^(a)	The 5 th percentile of the data is between ≥ 7.0 mg/L.	The 5 th percentile of the data is between 7.2 - 7.0 mg/L. Initiate baseline monitoring for this variable.
Turbidity ^(a)	Changes in turbidity are related to minor man-made modifications. Some silting of habitats is expected.	Initiate baseline monitoring for this variable.
Nutrients		
TIN-N	The 50 th percentile of the data is between ≤ 0.25 mg/L.	The 50 th percentile of the data is between 0.2 - 0.25 mg/L.
PO ₄ -P	The 50 th percentile of the data is between ≤ 0.125 mg/L.	The 50 th percentile of the data is between 0.1 - 0.125 mg/L.
Response variables^(a)		
Chl-a phytoplankton	The 50 th percentile of the data is between < 15 µg/L.	The 50 th percentile of the data is between 12 - 15 µg/L.
Chl-a periphyton	The 50 th percentile of the data is between ≤ 21 mg/m ² .	The 50 th percentile of the data is between 17 - 21 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data is between ≤ 1.5 mg/L.	The 50 th percentile of the data is between 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.015 mg/L.
Other toxics	The 95 th percentile of the data is between within the TWQR as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

(a) N/A: No data were available for this assessment.

Table 5.11: PES categories and overall site assessment for Touws River at J1TOUW-EWR3

RIVER	Touws River		WATER QUALITY MONITORING POINTS	
WQRU			RC	N/A
EWR SITE	J1TOUW-EWR3		PES	J1H018Q01 (Water Management System (WMS) code 102147), located upstream of the EWR site. (Data record: 2000 – 2014; number of samples (n) = ± 128).
Confidence assessment		2.5		
Water Quality Constituents		PES Value	Category/Comment	
Inorganic salts (mg/L)	SO ₄	N/A	All guidelines exceeded due to high saline geology of the area. It is assumed that some increase in salinity may be expected due to irrigation return flows. No large urban centers are situated in this area.	
	Na	2 016.9		
	Mg	370.1		
	Ca	258.2		
	Cl	3 494.6		
	K	37.06		
Nutrients (mg/L)	SRP	0.033	D	
	TIN	0.079	A	
Physical Variables	pH (5 th – 95 th %)	7.6 and 8.6	B	
	Temperature (°C)	N/A	B. Impacts expected at low flows. B. Impacts expected at low flows, although on-site data still shows high levels. B. Changes in turbidity appear to be largely related to natural with minor man-made modifications.	
	Dissolved oxygen	N/A		
	Turbidity (NTU)	N/A		
	Electrical conductivity (mS/m)	1181.8		
Response variables	Chl a: phytoplankton	N/A	N/A	
	Macroinvertebrates	74.0%	C	
	Diatoms	8.6 (average)	D	
	Fish community score	56.8%	D	
Toxics	Ammonia (as N)	0.034	A	
	Fluoride (as F)	0.43	A	
OVERALL SITE CLASSIFICATION		<i>B/C Category</i>		

Note:

RC: Information available to the water quality specialist on water quality conditions and land-use were used as no RC data were available and the A Category benchmarks tables in DWAF (2008) were considered unsuitable due to the high geology-based salinities in the area.

Table 5.12: Water quality EcoSpecs and TPCs (Category B/C) for Touws River at J1TOUW-EWR3

Metrics	EcoSpecs	TPCs
Inorganic salt ions		
Sulphate as SO ₄	N/A	
Sodium as Na	The 95 th percentile of the data is between ≤ 2000 mg/L.	The 95 th percentile of the data is between 1600 - 2000 mg/L.
Magnesium as Mg	The 95 th percentile of the data is between ≤ 370 mg/L.	The 95 th percentile of the data is between 300 - 370 mg/L.
Calcium as Ca	The 95 th percentile of the data is between ≤ 260 mg/L.	The 95 th percentile of the data is between 200 - 260 mg/L.
Chloride as Cl	The 95 th percentile of the data is between ≤ 3500 mg/L.	The 95 th percentile of the data is between 2800 - 3500 mg/L.
Potassium as K	The 95 th percentile of the data is between ≤ 37 mg/L.	The 95 th percentile of the data is between 30 - 37 mg/L.
Physical variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data is between ≤ 1100 mS/m.	The 95 th percentile of the data is between 880 - 1100 mS/m.
pH	The 5 th percentile of the data is between 6.5 – 8.0, and the 95 th percentile 8.0 - 8.8.	The 5 th percentile of the data is ≤ 6.3 and the 95 th percentile is ≥ 8.6.
Temperature	Natural temperature range.	Initiate baseline monitoring for this variable.
Dissolved oxygen	The 5 th percentile of the data is between ≥ 7.0 mg/L.	The 5 th percentile of the data is between 7.2 - 7.0 mg/L. Initiate baseline monitoring for this variable.
Turbidity	Changes in turbidity are related to minor man-made modifications. Some silting of habitats is expected.	Initiate baseline monitoring for this variable.
Nutrients		
TIN-N	The 50 th percentile of the data is between ≤ 0.25 mg/L.	The 50 th percentile of the data is between 0.2 - 0.25 mg/L.
PO ₄ -P	The 50 th percentile of the data is between ≤ 0.075 mg/L.	The 50 th percentile of the data is between 0.06 - 0.075 mg/L.
Response variables		
Chl-a phytoplankton	The 50 th percentile of the data is between < 15 µg/L.	The 50 th percentile of the data is between 12 - 15 µg/L.
Chl-a periphyton	The 50 th percentile of the data is between ≤ 21 mg/m ² .	The 50 th percentile of the data is between 17 - 21 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data is between ≤ 1.5 mg/L.	The 50 th percentile of the data is between 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.015 mg/L.
Other toxics	The 95 th percentile of the data is between within the TWQR as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

(a) N/A: No data were available for this assessment.

Table 5.13: PES categories and overall site assessment for Gamka River at J2GAMK-EWR4

RIVER	Gamka River		WATER QUALITY MONITORING POINTS	
WQRU			RC	A category benchmark tables from DWAF (2008) were used.
EWR SITE	J2GAMK-EWR4		PES	Data were sourced from DWS gauging weir J2H016Q01 (WMS code 102173), located downstream Gamkapoort Dam and upstream of the EWR site. (Data record: 2007 – 2014; n = 127).
Confidence assessment		3.0		
Water Quality Constituents			PES Value	Category/Comment
Inorganic salts (mg/L)	SO ₄		N/A	
	Na		114.0	Exceeds the 70mg/l (TWQR) for Agricultural Use: Irrigation
	Mg		20.5	No guideline
	Ca		57.6	No guideline
	Cl		155.5	Exceeds the 100mg/l (TWQR) for Agricultural Use: Irrigation
	K		7.9	No guideline
Nutrients (mg/L)	SRP		0.07	D
	TIN		0.523	B
Physical Variables	pH (5 th – 95 th %)		7.4 and 8.6	
	Temperature (°C)		N/A	Impact expected as the site is downstream of the Gamkapoort Dam (constructed in 1970)
	Dissolved oxygen		N/A	
	Turbidity (NTU)		N/A	
	Electrical conductivity (mS/m)		97.5	C. Natural salinity expected to be higher than the 30Ms/m A category benchmark value in DWAF (2008)
Response variables	Chl a: phytoplankton		N/A	N/A
	Macroinvertebrates		61.4%	C/D
	Diatoms		9.9	D
	Fish community score		60.4%	C/D
Toxics	Ammonia (as N)		0.015	A
	Fluoride (as F)		0.53	A
OVERALL SITE CLASSIFICATION			B/C Category	

Table 5.14: Water quality EcoSpecs and TPCs (PES: B/C) for Gamka River at J2GAMK-EWR4

Metrics	EcoSpecs: PES	TPCs: PES
Inorganic salt ions		
Sulphate as SO ₄	N/A	N/A
Sodium as Na	The 95 th percentile of the data is between ≤ 114 mg/L.	The 95 th percentile of the data is between 90 - 114 mg/L.
Magnesium as Mg	The 95 th percentile of the data is between ≤ 20 mg/L.	The 95 th percentile of the data is between 16 - 20 mg/L.
Calcium as Ca	The 95 th percentile of the data is between ≤ 58 mg/L.	The 95 th percentile of the data is between 47 - 58 mg/L.
Chloride as Cl	The 95 th percentile of the data is between ≤ 155 mg/L.	The 95 th percentile of the data is between 124 - 155 mg/L.
Potassium as K	The 95 th percentile of the data is between ≤ 8 mg/L.	The 95 th percentile of the data is between 6.5 - 8.0 mg/L.
Physical variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data is between ≤ 100 mS/m.	The 95 th percentile of the data is between 80 - 100 mS/m.
pH	The 5 th percentile of the data is between 5.9 - 6.5, and the 95 th percentile 8.0 - 8.8.	The 5 th percentile of the data is ≤ 5.7 and the 95 th percentile is ≥ 8.6.
Temperature ^(a)	Moderate change to temperature due to upstream Gamkapoort Dam.	Initiate baseline monitoring for this variable.
Dissolved oxygen ^(a)	The 5 th percentile of the data is between ≥ 7.0 mg/L. Although some impacts are expected due to the upstream Gamkapoort Dam, the size of the river will mitigate the effects.	The 5 th percentile of the data is between 7.2 - 7.0 mg/L. Initiate baseline monitoring for this variable.
Turbidity ^(a)	Changes in turbidity are related to minor man-made modifications. Some silting of habitats is expected.	Initiate baseline monitoring for this variable.
Nutrients		
TIN-N	The 50 th percentile of the data is between ≤ 0.7 mg/L.	The 50 th percentile of the data is between 0.56 - 0.7 mg/L.
PO ₄ -P	The 50 th percentile of the data is between ≤ 0.125 mg/L.	The 50 th percentile of the data is between 0.1 - 0.125 mg/L.
Response variables		
Chl-a phytoplankton	The 50 th percentile of the data is between < 15 µg/L.	The 50 th percentile of the data is between 12 - 15 µg/L.
Chl-a periphyton	The 50 th percentile of the data is between ≤ 21 mg/m ² .	The 50 th percentile of the data is between 17 - 21 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data is between ≤ 1.5 mg/L.	The 50 th percentile of the data is between 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.015 mg/L.
Other toxics	The 95 th percentile of the data is between within the TWQR as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

(a) N/A: No data were available for this assessment.

Table 5.15: PES categories and overall site assessment for Buffels River at J1BUFF-EWR5

RIVER	Buffels	WATER QUALITY MONITORING POINTS	
WQRU		RC	Data were sourced from DWS gauging weir J1H028Q01 (WMS code 102152), located downstream Floriskraal Dam and upstream of the EWR site. Note that the monitoring point is not in the same Level II EcoRegion as the EWR site; however, this was the only data point between the dam and the site. (Data record: 1972 – 1977; n = 54, Conductivity: n = 33).
EWR SITE	J1BUFF-EWR5	PES	Data were sourced from DWS gauging weir J1H028Q01 (WMS code 102152) (Data record: 2010 – 2014; n = 44).
Confidence assessment		Confidence: 2.5	
Water Quality Constituents		PES Value	Category/Comment
Inorganic salts (mg/L)	SO ₄	61.42	No guideline
	Na	81.44	Exceeds the ≤ 70 mg/L (TWQR) for Agricultural Use: Irrigation.
	Mg	25.2	No guideline.
	Ca	48.68	No guideline.
	Cl	124.0	Exceeds the ≤ 100 mg/L (TWQR) for Agricultural Use: Irrigation.
	K	6.11	No guideline.
Nutrients (mg/L)	SRP	0.015	B. Levels have decreased since the 1970s.
	TIN	0.26	A/B. No change from the 1970s.
Physical Variables	pH (5 th – 95 th %)	7.5 and 8.5	B. No change from the 1970s.
	Temperature (°C)	N/A	Impact expected as the site is downstream the large Floriskraal Dam (constructed in 1965).
	Dissolved oxygen	N/A	
	Turbidity (NTU)	N/A	
	Electrical conductivity (mS/m)	78.1	No change from the 1970s.
Response variables	Chl a: phytoplankton	N/A	N/A
	Macroinvertebrates	72.0%	C
	Diatoms	11.2 (average)	C/D
	Fish community score	83.7%	B
Toxics	Ammonia (as N)	0.017	A
	Fluoride (as F)	0.66	A. No change from the 1970s.
OVERALL SITE CLASSIFICATION		B/C Category	

Table 5.16: Water quality EcoSpecs and TPCs (Category B/C) for Buffels River at J1BUFF-EWR5

Metrics	EcoSpecs: PES	TPCs: PES
Inorganic salt ions		
Sulphate as SO ₄	The 95 th percentile of the data is between ≤ 60 mg/L.	The 95 th percentile of the data is between 48 - 60 mg/L.
Sodium as Na	The 95 th percentile of the data is between ≤ 80 mg/L.	The 95 th percentile of the data is between 64 - 80 mg/L.
Magnesium as Mg	The 95 th percentile of the data is between ≤ 25 mg/L.	The 95 th percentile of the data is between 20 - 25 mg/L.
Calcium as Ca	The 95 th percentile of the data is between ≤ 50 mg/L.	The 95 th percentile of the data is between 40 - 50 mg/L.
Chloride as Cl	The 95 th percentile of the data is between ≤ 125 mg/L.	The 95 th percentile of the data is between 100 - 125 mg/L.
Potassium as K	The 95 th percentile of the data is between ≤ 6.0 mg/L.	The 95 th percentile of the data is between 4.8 - 6.0 mg/L.
Physical variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data is between ≤ 85 mS/m.	The 95 th percentile of the data is between 68 - 85 mS/m.
pH	The 5 th percentile of the data is between 5.9 - 6.5, and the 95 th percentile 8.0 - 8.8.	The 5 th percentile of the data is ≤ 5.7 and the 95 th percentile is ≥ 8.6.
Temperature	Moderate change to temperature expected due to upstream Floriskraal Dam.	Initiate baseline monitoring for this variable.
Dissolved oxygen	The 5 th percentile of the data is between ≥ 7.0 mg/L. Although some impacts are expected due to the upstream Floriskraal Dam, the size of the river should mitigate the effects.	The 5 th percentile of the data is between 7.2 - 7.0 mg/L. Initiate baseline monitoring for this variable.
Turbidity	Changes in turbidity are related to minor man-made modifications. Some silting of habitats is expected.	Initiate baseline monitoring for this variable.
Nutrients		
TIN-N	The 50 th percentile of the data is between ≤ 0.48 mg/L.	The 50 th percentile of the data is between 0.38 - 0.48 mg/L.
PO ₄ -P	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.015 mg/L.
Response variables		
Chl-a phytoplankton	The 50 th percentile of the data is between < 10 µg/L.	The 50 th percentile of the data is between 8 - 10 µg/L.
Chl-a periphyton	The 50 th percentile of the data is between ≤ 12 mg/m ² .	The 50 th percentile of the data is between 10 - 12 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data is between ≤ 1.5 mg/L.	The 50 th percentile of the data is between 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.015 mg/L.
Other toxics	The 95 th percentile of the data is between within the TWQR as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

(a) N/A: No data were available for this assessment.

Table 5.17: PES categories and overall site assessment for Gouritz River at J4GOUR-EWR6

RIVER	Gouritz River	WATER QUALITY MONITORING POINTS	
WQRU		RC	Data were sourced from DWS gauging weir J4H002Q01 (WMS code 102201), located upstream of the EWR site. (Data record: 1965 – 1967; n = 29)
EWR SITE	J4GOUR-EWR6	PES	Data were sourced from DWS gauging weir J4H002Q01 (Data record: 2010 – 2014; n = 85).
Confidence assessment		Confidence 3	
Water Quality Constituents		PES Value	Category/Comment
Inorganic salts (mg/L)	SO ₄	693.0	No guideline, but a reduction over time.
	Na	964.0	Exceeds the ≤ 70 mg/L (TWQR) for Agricultural Use: Irrigation. <i>Significant increase over time.</i>
	Mg	127.0	No guideline.
	Ca	123.3	No guideline.
	Cl	1 289.3	Exceeds the ≤ 100 mg/L (TWQR) for Agricultural Use: Irrigation.
	K	9.81	No guideline.
Nutrients (mg/L)	SRP	0.015	B/C
	TIN	0.05	A
Physical Variables	pH (5 th – 95 th %)	7.8 and 8.65	B
	Temperature (°C)	N/A	Impact expected at low flows.
	Dissolved oxygen	N/A	
	Turbidity (NTU)	N/A	
	Electrical conductivity (mS/m)	542.5	No change from the 1970s.
Response variables	Chl a: phytoplankton	N/A	N/A
	Macroinvertebrates	75.0%	C
	Diatoms	10.2 (average)	C/D
	Fish community score	50.1%	D
Toxics	Ammonia (as N)	0.015	A
	Fluoride (as F)	1.082	A. Substantial increase from the 1960s.
OVERALL SITE CLASSIFICATION		B/C Category	

Table 5.18: Water quality) EcoSpecs and TPCs (Category B/C) for Gouritz River at J4GOUR-EWR6

Metrics	EcoSpecs	TPCs
Inorganic salt ions		
Sulphate as SO ₄	The 95 th percentile of the data is between ≤ 690 mg/L.	The 95 th percentile of the data is between 550 - 690 mg/L.
Sodium as Na	The 95 th percentile of the data is between ≤ 960 mg/L.	The 95 th percentile of the data is between 770 - 960 mg/L.
Magnesium as Mg	The 95 th percentile of the data is between ≤ 130 mg/L.	The 95 th percentile of the data is between 105 - 130 mg/L.
Calcium as Ca	The 95 th percentile of the data is between ≤ 120 mg/L.	The 95 th percentile of the data is between 95 - 120 mg/L.
Chloride as Cl	The 95 th percentile of the data is between ≤ 1300 mg/L.	The 95 th percentile of the data is between 1050 - 1300 mg/L.
Potassium as K	The 95 th percentile of the data is between ≤ 10 mg/L.	The 95 th percentile of the data is between 8 - 10 mg/L.
Physical variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data is between ≤ 550 mS/m.	The 95 th percentile of the data is between 450 - 550 mS/m.
pH	The 5 th percentile of the data is between 5.9 - 6.5, and the 95 th percentile 8.0 - 8.8.	The 5 th percentile of the data is ≤ 5.7 and the 95 th percentile is ≥ 8.6.
Temperature	Natural temperature range.	Initiate baseline monitoring for this variable.
Dissolved oxygen	The 5 th percentile of the data is between ≥ 7.0 mg/L.	The 5 th percentile of the data is between 7.2 - 7.0 mg/L. Initiate baseline monitoring for this variable.
Turbidity	Changes in turbidity are related to minor man-made modifications. Some silting of habitats is expected.	Initiate baseline monitoring for this variable.
Nutrients		
TIN-N	The 50 th percentile of the data is between ≤ 0.25 mg/L.	The 50 th percentile of the data is between 0.2 - 0.25 mg/L.
PO ₄ -P	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.015 mg/L.
Response variables		
Chl-a phytoplankton	The 50 th percentile of the data is between < 10 µg/L.	The 50 th percentile of the data is between 8 - 10 µg/L.
Chl-a periphyton	The 50 th percentile of the data is between ≤ 12 mg/m ² .	The 50 th percentile of the data is between 10 - 12 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data is between ≤ 1.5 mg/L.	The 50 th percentile of the data is between 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.015 mg/L.
Other toxics	The 95 th percentile of the data is between within the TWQR as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

(a) N/A: No data were available for this assessment.

Table 5.19: PES categories and overall site assessment for Keurbooms River at K6KEUR-EWR8

RIVER	Keurbooms River		WATER QUALITY MONITORING POINTS
WQRU		RC	A Category benchmark tables from DWAF (2008) were used.
EWR SITE	K6KEUR-EWR8	PES	Data were sourced from DWS gauging weir K6H001Q01 (WMS code 102295), located far (about 20 km) upstream of the EWR site. (Data record: 2007 – 2014; n = 121; Fluorine (F) = 107)
Confidence assessment	Confidence 3		
Water Quality Constituents		PES Value	Category/Comment
Inorganic salts (mg/L)	SO ₄	27.90	No guideline.
	Na	70.24	Just outside the ≤ 70 mg/L (TWQR) for Agricultural Use: Irrigation.
	Mg	11.25	No guideline.
	Ca	12.08	No guideline.
	Cl	129.02	Just exceeds the ≤ 100 mg/L (TWQR) for Agricultural Use: Irrigation.
	K	2.76	No guideline.
Nutrients (mg/L)	SRP	0.012	B
	TIN	0.06	A
Physical Variables	pH (5 th – 95 th %)	6.6 and 7.8	B
	Temperature (°C)	N/A	B. Some impacts expected at low flows, although on-site data still shows high levels. B. Changes in turbidity appear to be largely related to natural with minor man-made modifications.
	Dissolved oxygen	N/A	
	Turbidity (NTU)	N/A	
	Electrical conductivity (mS/m)	54.6	B
Response variables	Chl a: phytoplankton	N/A	
	Macroinvertebrates	64.0%	C
	Diatoms	9.9 (average)	C/D
	Fish community score	76.4%	C
Toxics	Ammonia (as N)	0.001	A
	Fluoride (as F)	0.26	A
OVERALL SITE CLASSIFICATION		B Category	

Table 5.20: Water quality EcoSpecs and TPCs (Category B) for Keurbooms River at K6KEUR-EWR8

Metrics	Eco Specs	TPCs
Inorganic salt ions		
Sulphate as SO ₄	The 95 th percentile of the data is between ≤ 28 mg/L.	The 95 th percentile of the data is between 22 - 28 mg/L.
Sodium as Na	The 95 th percentile of the data is between ≤ 70 mg/L.	The 95 th percentile of the data is between 56 - 70 mg/L.
Magnesium as Mg	The 95 th percentile of the data is between ≤ 12 mg/L.	The 95 th percentile of the data is between 10 - 12 mg/L.
Calcium as Ca	The 95 th percentile of the data is between ≤ 12 mg/L.	The 95 th percentile of the data is between 10 - 12 mg/L.
Chloride as Cl	The 95 th percentile of the data is between ≤ 130 mg/L.	The 95 th percentile of the data is between 104 - 130 mg/L.
Potassium as K	The 95 th percentile of the data is between ≤ 3 mg/L.	The 95 th percentile of the data is between 2.4 - 3.0 mg/L.
Physical variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data is between ≤ 55 mS/m.	The 95 th percentile of the data is between 45 - 55 mS/m.
pH	The 5 th and 95 th percentile of the data is between 6.5 - 8.0.	The 5 th and 95 th percentile of the data is ≥ 6.3 and ≤ 8.2.
Temperature	Natural temperature range.	Initiate baseline monitoring for this variable.
Dissolved oxygen	The 5 th percentile of the data is between ≥ 7.0 mg/L.	The 5 th percentile of the data is between 7.2 - 7.0 mg/L. Initiate baseline monitoring for this variable.
Turbidity	Changes in turbidity are related to minor man-made modifications. Some silting of habitats is expected.	Initiate baseline monitoring for this variable.
Nutrients		
TIN-N	The 50 th percentile of the data is between ≤ 0.25 mg/L.	The 50 th percentile of the data is between 0.2 - 0.25 mg/L.
PO ₄ -P	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.0715 mg/L.
Response variables		
Chl-a phytoplankton	The 50 th percentile of the data is between < 10 µg/L.	The 50 th percentile of the data is between 8 - 10 µg/L.
Chl-a periphyton	The 50 th percentile of the data is between ≤ 12 mg/m ² .	The 50 th percentile of the data is between 9.6 - 12.0 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data is between ≤ 1.5 mg/L.	The 50 th percentile of the data is between 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data is between ≤ 0.015 mg/L.	The 50 th percentile of the data is between 0.012 - 0.015 mg/L.
Other toxics	The 95 th percentile of the data is between within the TWQR as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

(a) N/A: No data were available for this assessment.

Table 5.21: PES categories and overall site assessment for Olifants River at J3OLIF-EWR9

RIVER	Olifants River	WATER QUALITY MONITORING POINTS	
WQRU		RC	Information available to the water quality specialist on water quality conditions and land-use were available and the A Category benchmark tables in DWAF (2008) were considered unsuitable.
EWR SITE	J3OLIF-EWR9	PES	Data were sourced from DWS gauging weir J3H021Q01 (WMS code 102192) was used for the present state assessment located downstream of the EWR site and upstream of Stompdrift Dam. (Data record: 1982 – 1993; n = 127).
Confidence assessment		Confidence: 2.5	
Water Quality Constituents		PES Value	Category/Comment
Inorganic salts (mg/L)	SO ₄	1 353.4	No guideline but concentrations are high.
	Na	1 774.5	Exceeds the ≤ 70 mg/L (TWQR) for Agricultural Use: Irrigation.
	Mg	336.0	No guideline
	Ca	284.4	No guideline
	Cl	3 113	Exceeds the ≤ 100 mg/L (TWQR) for Agricultural Use: Irrigation.
	K	30.16	No guideline.
Nutrients (mg/L)	SRP	0.019	B/C
	TIN	0.11	A
Physical Variables	pH (5 th – 95 th %)	7.3 and 9.0	B/C but assumed to be linked to the groundwater signature.
	Temperature (°C)	N/A	C. Impact expected when little surface flow.
	Dissolved oxygen	N/A	
	Turbidity (NTU)	N/A	B/C. Impact expected due to extensive livestock farming and erosion in the area.
	Electrical conductivity (mS/m)	1 078.7	Natural salinity expected to be high due to the geology of the area
Response variables	Chl a: phytoplankton	N/A	N/A
	Macroinvertebrates	69.0%	C
	Diatoms	6.0 (average)	D/E
	Fish community score	N/A	
Toxics	Ammonia (as N)	0.038	B
	Fluoride (as F)	0.678	A
OVERALL SITE CLASSIFICATION		C Category	

(a) N/A: No data were available for this assessment.

Table 5.22: Water quality EcoSpecs and TPCs (Category C) for Olifants River at J3OLIF-EWR9

Metrics	Eco Specs	TPCs
Inorganic salt ions		
Sulphate as SO ₄	The 95 th percentile of the data is between ≤ 1350 mg/L.	The 95 th percentile of the data is between 1080 - 1350 mg/L.
Sodium as Na	The 95 th percentile of the data is between ≤ 1775 mg/L.	The 95 th percentile of the data is between 1420 - 1775 mg/L.
Magnesium as Mg	The 95 th percentile of the data is between ≤ 335 mg/L.	The 95 th percentile of the data is between 270 - 335 mg/L.
Calcium as Ca	The 95 th percentile of the data is between ≤ 285 mg/L.	The 95 th percentile of the data is between 230 - 285 mg/L.
Chloride as Cl	The 95 th percentile of the data is between ≤ 3000 mg/L.	The 95 th percentile of the data is between 2400 - 3000 mg/L.
Potassium as K	The 95 th percentile of the data is between ≤ 30 mg/L.	The 95 th percentile of the data is between 24 - 30 mg/L.
Physical variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data is between ≤ 1100 mS/m.	The 95 th percentile of the data is between 880 - 1100 mS/m.
pH	The 5 th percentile of the data is between 5.9 - 6.5, and the 95 th percentile 8.8 - 9.2.	The 5 th percentile of the data is ≤ 5.7 and the 95 th percentile is ≥ 9.0.
Temperature	The upper Olifants mostly runs underground, with water appearing in places. This is not groundwater from a deep aquifer, but water from the vadose zone. Elevated temperatures and low DO levels would be expected under these conditions. EcoSpecs and TPCs are therefore difficult to set for these variables, and should rather be linked to meeting biotic requirements and monitoring biotic responses.	Initiate baseline monitoring for this variable.
Dissolved oxygen		Initiate baseline monitoring for this variable where and if possible.
Turbidity	Changes in turbidity are related to minor man-made modifications. Some silting of habitats and temporary high turbidity levels are expected.	Initiate baseline monitoring for this variable.
Nutrients		
TIN-N	The 50 th percentile of the data is between ≤ 0.25 mg/L.	The 50 th percentile of the data is between 0.2 - 0.25 mg/L.
PO ₄ -P	The 50 th percentile of the data is between ≤ 0.025 mg/L.	The 50 th percentile of the data is between 0.02 - 0.025 mg/L.
Response variables^(a)		
Chl-a phytoplankton	The 50 th percentile of the data is between < 15 µg/L.	The 50 th percentile of the data is between 12 - 15 µg/L.
Chl-a periphyton	The 50 th percentile of the data is between ≤ 21 mg/m ² .	The 50 th percentile of the data is between 17 - 21 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data is between ≤ 1.5 mg/L.	The 50 th percentile of the data is between 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data is between ≤ 0.044 mg/L.	The 50 th percentile of the data is between 0.035 - 0.044 mg/L.
Other toxics	The 95 th percentile of the data is between within the TWQR as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

Table 5.23: PES categories and overall site assessment for Kammanassie River at J3KAMM-EWR10

RIVER	Kammanassie River		WATER QUALITY MONITORING POINTS
WQRU		RC	No data were available for the water quality assessment. Land use and available information, diatom data, <i>in situ</i> water quality data and survey notes were used to provide an expert opinion and generate a PAI model and integrated water quality category for the site.
EWR SITE	J3KAMM-EWR10	PES	N/A
Confidence assessment	Confidence: 2		
Water Quality Constituents	PES Value	Category/Comment	
Inorganic salts (mg/L)	SO ₄	N/A	N/A
	Na	N/A	N/A
	Mg	N/A	N/A
	Ca	N/A	N/A
	Cl	N/A	N/A
	K	N/A	N/A
Nutrients (mg/L)	SRP	N/A	N/A
	TIN	N/A	N/A
Physical Variables	pH (5 th – 95 th %)	N/A	N/A
	Temperature (°C)	N/A	N/A
	Dissolved oxygen	N/A	N/A
	Turbidity (NTU)	N/A	N/A
	Electrical conductivity (mS/m)	N/A	N/A
Response variables	Chl a: phytoplankton	N/A	N/A
	Macroinvertebrates	C/D	
	Diatoms	C/D	The biological water quality at this site was Moderate. Nutrient levels, organic pollution and salinity were elevated with salinity and organic pollution levels becoming problematic. The improvement in diatom-based water quality could mainly be ascribed to higher flows during July 2014 which allowed for the flushing of pollutants as diatom species associated with elevated flows were abundant.
	Fish community score	D	
Toxics	Ammonia (as N)		
	Fluoride (as F)		
OVERALL SITE CLASSIFICATION		C Category	

a) N/A: No data were available for this assessment.

Note that limited water quality data exists for the Kammanassie River system. The water quality assessment is therefore based on available information and expert judgement.

Table 5.24: PES categories and overall site assessment for Groot Brak WQSU 1 & 2. (expert judgment)

RIVER	Groot Brak River		WATER QUALITY MONITORING POINTS	
WQSU	WQSU 1 + 2		RC	Default boundary tables for "A" category river
EWR SITE	None		PES	Groot Brak R. at Ernest Robertson dam - K2H005-Q01 (1983 – 1996; n = 29)
Confidence assessment		Very low because extrapolated from outflow from a dam, and not current data.		
Water Quality Constituents			Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄			No data
	Na ₂ SO ₄			
	MgCl ₂			
	CaCl ₂			
	NaCl			
	CaSO ₄			
Nutrients (mg/L)	SRP (mg/l)		0.042	Unreliable data used
	TIN (mg/l)		<0.25	A category
Physical Variables	pH (5 th – 95 th %)		4.62	Naturally acidic
	Temperature			No data
	Dissolved oxygen			Expected to be high
	Turbidity (NTU)			Expected to be low
	Electrical conductivity (mS/m)		<30	A category
Response variables	Chl a: periphyton			No data
	Chl a: phytoplankton			No data
	Macroinvertebrates (ASPT)			No data
	Fish community score			No data
Toxics				No data – expected to be minimal
OVERALL SITE CLASSIFICATION			A/B (from expert judgment)	

Table 5.25: Water quality Ecospecs and TPC's (Category A/B) for Groot Brak River at WQSU 1&2

RIVER	Groot Brak River	WATER QUALITY MONITORING POINTS			
WQSU	WQSU 1&2	DWAF WQ WMS	Currently, no monitoring station		
EWR SITE	GB 1	RHP	Currently, no monitoring site		
Confidence in PES assessment					
Very Low because extrapolated from WQSU 3					
Water Quality Constituents	PES Category	WQ Ecospecs	Improvement required?		
TPC	Monitoring frequency				
Inorganic salts (mg/L)	MgSO ₄	≤23 mg/L		95 th percentile to be < 23 mg/L	Monthly
	Na ₂ SO ₄	≤33 mg/L		95 th percentile to be < 33 mg/L	
	MgCl ₂	≤30 mg/L	N/A	95 th percentile to be < 30 mg/L	
	CaCl ₂	≤57 mg/L		95 th percentile to be < 57 mg/L	
	NaCl	≤191 mg/L		95 th percentile to be < 191 mg/L	
Nutrients (mg/L)	SRP	≤0.012 mg/L	Yes to A	50 th percentile to be < 0.012 mg/L	Monthly
	TIN	≤0.25 mg/L	No	50 th percentile to be < 0.25 mg/L	Monthly
Physical Variables	pH	< 7.9	No	95 th percentile to be < 7.9	Monthly
	Temperature	Maintain range	N/A	Maintain natural range	Monthly
	Dissolved oxygen	7 – 8 mg/L	N/A	5 th percentile to be > 7 mg/L	Monthly
	Turbidity (NTU)	Moderate change	N/A	Moderate change allowed	Monthly
	Electrical conductivity (mS/m)	Category = A	≤30 mS/m	No	95 th percentile to be < 30 mS/m
Response variables	Chl a: periphyton	≤12 mg/m ² (B category)	N/A	50 th percentile to be < 12 mg/m ²	Quarterly
	Chl a: phytoplankton	≤ 15 µg/L (B category)		50 th percentile to be < 15 µg/L	
Toxics	Macroinvertebrates (ASPT)	See Ecospecs for fish and invertebrates respectively			
	Fish community score				
	Instream toxicity	No data	Assess only if the biomonitoring results indicate that there is a serious problem and the cause is unknown.		

Table 5.26: PES categories and overall site assessment for Groot Brak Water at WQSU 3

RIVER	1.1.1 WATER QUALITY MONITORING POINTS	
	Groot Brak River	RC
WQSU	WQSU 3	Moordkull R. at Banff - K1H005 (1979 – 1982 n = 91)
EWR SITE	GB 1	K1H005 (2002 – 2006 n = 51)
Confidence assessment	Medium. NB: Because extrapolated from another catchment. Biological data support inferred water quality	
Water Quality Constituents		
Inorganic salts (mg/L)	MgSO ₄	
	Na ₂ SO ₄	
	MgCl ₂	
	CaCl ₂	
	NaCl	
	CaSO ₄	
Nutrients (mg/L)	SRP	0.006*
	TIN	0.04
	pH (5 th – 95 th %)	Not calculated
Physical Variables	Temperature	
	Dissolved oxygen	
	Turbidity (NTU)	
	Electrical conductivity (mS/m)	40
	Fish community score	-
Response variables	Chl a: periphyton	
	Chl a: phytoplankton	
Toxics	Macroinvertebrates (ASPT)	A
	Fish community score	C
OVERALL SITE CLASSIFICATION		
		B (from PAI model)

Table 5.27: Water quality Ecospecs and TPC's (Category B) for Groot Brak River at WQSU 3

RIVER	Groot Brak River	WATER QUALITY MONITORING POINTS				
WQSU	3	DWAF WQ WMS	Currently, no monitoring station			
EWR SITE	GB 1	RHP	Currently, no monitoring site			
Confidence in PES assessment						
Low – medium, because extrapolated from Moordkuil R. Biological data supports inferred PES for water quality.						
Water Quality Constituents	PES Category	WQ Ecospecs	Improvement required?	TPC	Monitoring frequency	
Inorganic salts (mg/L)	MgSO ₄	≤23 mg/L		95 th percentile to be < 23 mg/L		
	Na ₂ SO ₄	≤33 mg/L		95 th percentile to be < 33 mg/L		
	MgCl ₂	≤30 mg/L	N/A	95 th percentile to be < 30 mg/L	Monthly	
	CaCl ₂	≤57 mg/L		95 th percentile to be < 57 mg/L		
Nutrients (mg/L)	NaCl	≤191 mg/L		95 th percentile to be < 191 mg/L		
	SRP	≤0.02 mg/L	Yes to B	50 th percentile to be < 0.02 mg/L	Monthly	
Physical Variables	TIN	≤0.25 mg/L	No	50 th percentile to be < 0.25 mg/L	Monthly	
	pH	< 7.9	No	95 th percentile to be < 7.9	Monthly	
	Temperature	Maintain range	N/A	Maintain natural range	Monthly	
	Dissolved oxygen	No data, but not considered to be problem in this river.	N/A	5 th percentile to be > 7 mg/L	Monthly	
	Turbidity (NTU)	Moderate change	N/A	Moderate change allowed	Monthly	
	Electrical conductivity (mS/m)	Category = A	≤30 mS/m	No	95 th percentile to be < 30 mS/m	Monthly
	Chl a: periphyton	No data. Visual inspection March & June. No obvious sign of algae.	≤12 mg/m ² (B category)	N/A	50 th percentile to be < 12 mg/m ²	Quarterly
	Chl a: phytoplankton		≤ 15 µg/L (B category)		50 th percentile to be < 15 µg/L	
Response variables	Macroinvertebrates (ASPT)	See Ecospecs for fish and invertebrates respectively				
	Fish community score					
	Instream toxicity	Assess only if the biomonitoring results indicate that there is a serious problem and the cause is unknown.				
Toxics	No data. Possibly some pesticides					

Table 5.28: PES categories and overall site assessment for Groot Brak WQSU 4 (*boundary value recalibrated) (expert judgment)

RIVER		Groot Brak		WATER QUALITY MONITORING POINTS	
WQSU		4		RC	Groot Brak R. at Wolwedans - K2H002-Q01 (1976 – 1978; n = 68)
EWR SITE		None		PES	K2H002 (2002 – 2006; n = 57)
Confidence assessment		Very good. Monitoring station located in WQSU, above point source of village. Data collected from > 15 years before dam construction			
Water Quality Constituents		RC Value	PES Value	Category/Comment	
Inorganic salts (mg/L)	MgSO ₄			No data	
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
Nutrients (mg/L)	SRP	0.016*	0.037	Category = C. Trend increasing	
	TIN	0.04	0.075	Category = A. Trend increasing	
Physical Variables	pH (5 th – 95 th %)		6.8 – 8.1	Based on Monitoring station located in WQSU, above point source of village	
	Temperature		No data	May be a problem considering downstream of dam. Requires monitoring	
	Dissolved oxygen		No data		
	Turbidity (NTU)		No data		
	Electrical conductivity (mS/m)	30	52	Category = B. Trend increasing	
Response variables	Chl a: periphyton			No data	
	Chl a: phytoplankton			No data	
	Macroinvertebrates (ASPT)			No data	
	Fish community score			No data	
Toxics				No data	
OVERALL SITE CLASSIFICATION		B/C (expert judgment)			

Table 5.29: Water quality Ecospecs and TPC's (Category B/C) for Groot Brak River at WQSU 4

RIVER		WATER QUALITY MONITORING POINTS					
WQSU	Groot Brak River	DWAF WQ WMS	Currently, no monitoring station				
EWR SITE	GB 1	RHP	Currently, no monitoring site				
Confidence in PES assessment							
		Very Low, because extrapolated from WQSU 3.					
Water Quality Constituents		PES Category	WQ Ecospecs	Improvement required?	TPC	Monitoring frequency	
Inorganic salts (mg/L)	MgSO ₄	Not available	≤23 mg/L		95 th percentile to be < 23 mg/L	Monthly	
	Na ₂ SO ₄		≤33 mg/L		95 th percentile to be < 33 mg/L		
	MgCl ₂		≤30 mg/L	N/A	95 th percentile to be < 30 mg/L		
	CaCl ₂		≤57 mg/L		95 th percentile to be < 57 mg/L		
	NaCl		≤191 mg/L		95 th percentile to be < 191 mg/L		
Nutrients (mg/L)	SRP	Category = C.	≤0.025 mg/L	Yes to B/C	50 th percentile to be < 0.025 mg/L	Monthly	
	TIN	Category = A	≤0.25 mg/L	No	50 th percentile to be < 0.25 mg/L	Monthly	
Physical Variables	pH	Naturally acidic river	< 7.9	No	95 th percentile to be < 7.9	Monthly	
	Temperature	No data, but not considered to be problem in this river.	Maintain range	N/A	Maintain natural range	Monthly	
	Dissolved oxygen		7 – 8 mg/L	N/A	5 th percentile to be > 7 mg/L	Monthly	
	Turbidity (NTU)		Moderate change	N/A	Moderate change allowed	Monthly	
	Electrical conductivity (mS/m)		Category = A	≤30 mS/m	No	95 th percentile to be < 30 mS/m	Monthly
	Chl a: periphyton		No data. Visual inspection March & June. No obvious sign of algae.	≤12 mg/m ² (B category)	N/A	50 th percentile to be < 12 mg/ m ²	Quarterly
Chl a: phytoplankton			≤ 15 µg/L (B category)		50 th percentile to be < 15 µg/L		
Response variables		Macroinvertebrates (ASPT)	See Ecospecs for fish and invertebrates respectively				
Fish community score		C (this study)					
Instream toxicity		No data					
Toxics		No data. Possibly some pesticides		Assess only if the biomonitoring results indicate that there is a serious problem and the cause is unknown.			

Table 5.30: Water quality Ecospecs and TPC's for Malgas River at WQSU2

RIVER	Malgas River	WATER QUALITY MONITORING POINTS			
WQSU	WQSU 2	RC	Default values		
EWR SITE	Mal 1	PES	Malgas R. at Blanco - K3H004-Q01 ('01 – '06 n = 53)		
Confidence assessment	Good for the overall WQSU. Low for the EWR Site.				
Water Quality Constituents		RC Value	PES Value	Category/Comment	G-power (Confidence)
Inorganic salts (mg/L)	MgSO ₄				No data
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
Nutrients (mg/L)	SRP	≤0.005	0.038	Category = C. Trend increasing.	0.116 (Low)
	TIN	≤0.25	0.413	Category = A/B. Trend increasing	
Physical Variables	pH (5 th + 95 th %ile)		4.3 + 7.2	Naturally acidic.	0.84 (High)
	Temperature	16.3 (50%ile)	-	No PES data. Not expected to be a problem since no dam upstream	
	Dissolved oxygen			No data. Could be a problem due to quarry	
	Turbidity (NTU)			Category = A. Trend = slight increase	0.64 (Medium)
Response variables	Chl a: periphyton			No data	No data (but visual inspection at EWR site indicated localised problem).
	Chl a: phytoplankton			No data	
	Macroinvertebrates			A category; ASPT = 8.2, SASS = 164 (this study)	
	Fish community score			C/D (this study)	
Toxics				No data. Possible toxicity from quarry/cement/asphalt works, outflows from village and agricultural return flows	
OVERALL SITE CLASSIFICATION		B Category (from PAI model)			

Table 5.31: PES categories and overall site assessment for Maalgate River

RIVER		Moeras/Maalgate River	WATER QUALITY MONITORING POINTS	
WQSU		N/A	RC	Default boundary tables for A category river
EWR SITE		Moe 1 & Maa 2	PES	Maalgate R. @ Noetze Kamma K3H003 ('02 – '06; n = 52)
Confidence assessment		Good.		
Water Quality Constituents		Value	Category/Comment	
Nutrients (mg/L)	SRP	0.019	Category = B (TP = 0.015 mg/L)	
	TIN	0.1	Category = A	
Physical Variables	pH (5 th – 95 th %)	5.7 – 7.7	Naturally acidic	
	Temperature		No data. May be impacted (by abstraction)	
	Dissolved oxygen		No data. May be impacted (due to excessive abstraction)	
	Turbidity (NTU)		No data. Site visit indicated not elevated.	
	Electrical conductivity (mS/m)	63	Category = C	
Response variables	Chl a: periphyton		No data. Site visit indicated extensive periphyton	
	Chl a: phytoplankton		No data	
	Macroinvertebrates (ASPT)		D (present study) Fair (RHP)	
	Fish community score		D (present study)	
Toxics			No data – expected to be pesticides from intensive agricultural activity	
OVERALL SITE CLASSIFICATION		C Category (from expert judgement)		
Boundary values				
Determinant		Lower boundary value	Upper boundary value	
TIN (mg/L) – 50 th %ile			2.0	
PHYTOPLANKTON Chl a (µg/L) – 50 th %ile		15	20	
SRP or PO ₄ (mg/L) – 50 th %ile			0.058	
PERIPHYTON Chl a (mg/m ²) – 50 th %ile		12	21	
pH – 5 th %ile and 95 th %ile		5 th percentile: 5.00 – 5.23		
TEMPERATURE (°C) – 10 th %ile and 90 th %ile		Vary by no more than 2°C from natural range		
Dissolved oxygen (mg/L) - 5 th %ile		6		
BIOLOGICAL INDICATOR (ASPT)		5.67		

Table 5.32: PES categories and overall site assessment for Gwaing (Malgas/Keur River) at WQSU1

RIVER	Malgas River (Keur River)		WATER QUALITY MONITORING POINTS	
WQSU	WQSU 1	RC	Rooi R. @ George K3H002-Q01 ('77 – '79 n = 84)	
EWR SITE	None	PES	K3H002-Q01 ('01 – '06 n = 65)	
Confidence assessment		Low. NB: extrapolated from an adjacent catchment		
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄			No data
	Na ₂ SO ₄			
	MgCl ₂			
	CaCl ₂			
	NaCl			
	CaSO ₄			
Nutrients (mg/L)	SRP	0.014	≤0.014*	Category = A
	TIN	0.04	≤0.25*	Category = A
Physical Variables	pH (5 th – 95 th %)		3.7 – 6.7	Naturally acidic
	Temperature			No data but expected to be natural as little development in upstream catchment.
	Dissolved oxygen			
	Turbidity (NTU)			
	Electrical conductivity (mS/m)	16	17	Category = A
Response variables	Chl a: periphyton			No data
	Chl a: phytoplankton			
	Macroinvertebrates			Good (RHP)
	Fish community score			Natural (RHP)
Toxics				No data
OVERALL SITE CLASSIFICATION		A Category (from expert judgement)		
Boundary values				
Determinant		Lower boundary value	Upper boundary value	
TIN (mg/L) – 50 th %ile			≤ 0.25	
PHYTOPLANKTON Chl a (µg/L) – 50 th %ile			<10	
SRP or PO4 (mg/L) – 50 th %ile			≤ 0.005	
PERIPHYTON Chl a (mg/m ²) – 50 th %ile			< 1.7	
pH – 5 th %ile and 95 th %ile		6.5 – 8.00		
TEMPERATURE (°C) – 10 th %ile and 90 th %ile		Nat temp range		
Dissolved oxygen (mg/L) - 5 th %ile		>8		
BIOLOGICAL INDICATOR (ASPT)		7		

Table 5.33: PES categories and overall site assessment for Gwaing (Malgas River) at WQSU2

RIVER	Malgas River		WATER QUALITY MONITORING POINTS		
WQSU	WQSU 2	RC	Default values		
EWR SITE	Mal 1	PES	Malgas R. @ Blanco - K3H004-Q01 ('01 – '06 n = 53)		
Confidence assessment	Good for the overall WQSU. Low for the EWR Site.				
Water Quality Constituents		RC Value	PES Value	Category/Comment	G-power (Confidence)
Inorganic salts (mg/L)	MgSO ₄				No data
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
Nutrients (mg/L)	SRP	≤0.005	0.038	Category = C. Trend increasing.	0.116 (Low)
	TIN	≤0.25	0.413	Category = A/B. Trend increasing	
Physical Variables	pH (5 th – 95 th %)		4.3 – 7.2	Naturally acidic.	0.84 (High)
	Temperature	16.3 (50 th ile)	-	No PES data. Not expected to be a problem since no dam upstream	
	Dissolved oxygen			No data. Could be a problem due to quarry	
	Turbidity (NTU)			No data. Could be a problem due to quarry	
	Electrical conductivity (mS/m)	<30	15	Category = A. Trend = slight increase	0.64 (Medium)
Response variables	Chl a: periphyton			No data	No data (but visual inspection indicated localised problem).
	Chl a: phytoplankton			No data	
	Macroinvertebrates			A category; ASPT = 8.2, SASS = 164 (this study)	
	Fish community score			C/D (this study)	
Toxics				No data. Possible toxicity from quarry/cement/asphalt works, outflows from village and return flows	
OVERALL SITE CLASSIFICATION		B Category (from PAI model). Likely to be better at EWR site itself.			
Boundary values					
Determinant		Lower boundary value		Upper boundary value	
TIN (mg/L) – 50 th %ile				0.75	
PHYTOPLANKTON Chl a (µg/L) – 50 th %ile		5		10	
SRP or PO ₄ (mg/L) – 50 th %ile				0.02	
PERIPHYTON Chl a (mg/m ²) – 50 th %ile		1.7		12	
pH – 5 th %ile and 95 th %ile				5 th percentile: 6.00 – 6.24 95 th percentile: 8.37 – 8.69	
TEMPERATURE (°C) – 10 th %ile and 90 th %ile				Nat temp range	
Dissolved oxygen (mg/L) - 5 th %ile		7			
BIOLOGICAL INDICATOR (ASPT)		6.34			

Table 5.34: PES categories and overall site assessment for Gwaing (Gwaing River) at WQSU3

RIVER	Gwaing River		WATER QUALITY MONITORING POINTS	
WQSU	WQSU 3		RC	Default values
EWR SITE	None		PES	No DWAF monitoring station
Confidence assessment		Low for RC as default values used. Medium for the PES as no monitoring station, but supported by biomonitoring data and (limited) supplementary data.		
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄			No data
	Na ₂ SO ₄			
	MgCl ₂			
	CaCl ₂			
	NaCl			
	CaSO ₄			
Nutrients (mg/L)	SRP	≤0.005	1.4 mg/L	Category = F
	TIN	≤0.25		Category = D
Physical Variables	pH (5 th – 95 th %)			No data. Likely to be a problem with low DO and elevated turbidity
	Temperature			
	Dissolved oxygen			
	Turbidity (NTU)			
	Electrical conductivity (mS/m)	<30	49	
Response variables	Chl a: periphyton			No data
	Chl a: phytoplankton			
	Macroinvertebrates (ASPT)			RHP = "Poor water quality"
	Fish community score			
Toxics				No data, but likely to be a problem
OVERALL SITE CLASSIFICATION		D/E Category (expert judgement)		
Boundary values				
Determinant		Lower boundary value	Upper boundary value	
TIN (mg/L) – 50 th %ile			4.0	
PHYTOPLANKTON Chl a (µg/L) – 50 th %ile		20	30	
SRP or PO ₄ (mg/L) – 50 th %ile			0.125	
PERIPHYTON Chl a (mg/m ²) – 50 th %ile		21	84	
pH – 5 th %ile and 95 th %ile		5 th percentile: 5.46 – 5.7 95 th percentile: 8.56 – 10.00		
TEMPERATURE (°C) – 10 th %ile and 90 th %ile		Vary by no more than 4°C from natural range		
Dissolved oxygen (mg/L) - 5 th %ile		4		
BIOLOGICAL INDICATOR (ASPT)		5		

Where a difference in the water quality values for the Ecological Reserve and Basic Human Needs Reserve was found, the stricter or more protective value was selected for the water quality component of the Reserve.

Table 5.35: PES categories and overall site assessment for the Kaaimans River at WQSU 2

RIVER	Kaaimans River		WATER QUALITY MONITORING POINTS		
WQSU	WQSU 2	RC	Kaaimans R. @ Barbierskraal - K3H001-Q01 ('77 – '81; n = 175)		
EWR SITE	Ka1	PES	K3H001-Q01 ('01 – '06; n = 56)		
Confidence assessment		High for both the RC and PES, because the monitoring station is in the WQSU, has a good length of data, and because there is little land transformation in the catchment.			
Water Quality Constituents		RC Value	PES Value	Category/Comment	G-power (Confidence)
Inorganic salts (mg/L)	MgSO ₄				No data
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
Nutrients (mg/L)	SRP	0.011	0.028*	Category = B. Trend = slight increase	0.122 (Low)
	TIN	0.04	0.061	Category = A. Very slight increase in trend	
Physical Variables	pH (5 th – 95 th %)		4.4 – 7.4	Naturally acidic waters	0.87 (High)
	Temperature			No data. Unlikely to be a problem	
	Dissolved oxygen				
	Turbidity (NTU)			No data. Visual inspection = low turbidity. Unlikely to be a problem	
	Electrical conductivity (mS/m)	16	17	A Category. Trend = stable	0.87 (High)
Response variables	Chl a: periphyton			No data. Visual inspection = no excessive algal growth	
	Chl a: phytoplankton				
	Macroinvertebrates			A category; ASPT = 8.0, SASS = 175 (this study). Natural (RHP)	
	Fish community score			B	
Toxics				No data, but unlikely to be a problem	
OVERALL SITE CLASSIFICATION		A Category (calculated from PAI model)			

Table 5.36: Water Ecospecs for the Kaaimans River in K30C

Quality Constituent	Parameter	Ecological Reserve Requirements	Basic Human Needs Requirement ⁵	Reserve Requirement: water quality
General chemistry – major inorganic salts	MgSO ₄ (mg/l) ¹	< 16	N/A	< 16
	Na ₂ SO ₄ (mg/l) ¹	< 20	N/A	< 20
	MgCl ₂ (mg/l) ¹	< 15	N/A	< 15
	CaCl ₂ (mg/l) ¹	< 21	N/A	< 21
	NaCl (mg/l) ¹	< 45	N/A	< 45
General chemistry – Major Ions	Sodium (mg/l)	N/A	<200	<200
	Magnesium (mg/l)	N/A	<100	<100
	Chloride (mg/l)	N/A	<200	<200
	Calcium (mg/l)	N/A	<80	<80
	Sulphate (mg/l)	N/A	<400	<400
Nutrients	Phosphate (PO ₄) (mg/l) ²	<0.02mg/L	N/A	<0.02mg/L
	Total Inorganic Nitrogen (mgN/l) ²	<0.25mg/L	N/A	<0.25mg/L
Physical water quality	pH (range) 5 th percentile 95 th percentile	6.7 7.4	5 9.5	5 7.4
	Dissolved Oxygen (mg/l) ¹	>1.7 mg/L	N/A	>1.7 mg/L
	Temperature ¹	Small change from natural		Small change from natural
	Electrical conductivity (mS/m) – USE ONLY IF AGGREGATED SALTS CANNOT BE PRODUCED	≤30mS/m	0-70	≤30mS/m
Biological water quality	Chl-a: periphyton ³	<1.7 mg/m ²	N/A	
	Chl-a: phytoplankton ³	< 10 µg/L (A category)	N/A	
	Biotic community composition - macroinvertebrates	ASPT: 8 (A category)		
	In-stream toxicity	In-stream toxicity should not occur		
Toxics and complex mixtures	Toxics (as listed in DWAF, 1996 ⁶)	≤ TWQR	≤ TWQR	≤ TWQR

NOTES:

¹ : 95th percentile compliance. ²: 50th percentile compliance. ³ Chl-a is not applicable to Desktop Reserve studies. ⁴ 90th percentile compliance

⁵ ref: *South African Water Quality Guidelines, Volume 1: Domestic Water Use, 2nd Ed.* 1996. Department of Water Affairs and Forestry. Pretoria, South Africa.

⁶ ref: *South African Water Quality Guidelines, Volume 7: Aquatic Ecosystems, 2nd Ed.* 1996. Department of Water Affairs and Forestry. Pretoria, South Africa.

Table 5.37: PES categories and overall site assessment for the Diep River at WQSU 3

RIVER	Diep River	WATER QUALITY MONITORING POINTS		
WQSU	WQSU 3 (start of level II Ecoregion 20.02 to the beginning of the mountain stream zone)	RC	Diep R. @ Woodville Forest Reserve K4H003Q01 ('77 – '80; n = 58)	
EWR SITE	3	PES	K4H003Q01 ('03 – '07; n = 36)	
Confidence assessment		Moderate to high confidence		
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄	8.86	15.87	TEACHA was used for data assessment. Salts = an A category
	Na ₂ SO ₄	0.00	0.00	
	MgCl ₂	14.83	14.52	
	CaCl ₂	7.18	9.89	
	NaCl	102.10*	94.31	
Nutrients (mg/L)	SRP	0.003	0.018	B/C category.
	TIN	0.04	0.07	A category.
Physical Variables	pH (5 th + 95 th %ile)	4.8 + 6.2	4.8 + 7.5	B category.
	Temperature	-	-	No data but no impacts expected.
	Dissolved oxygen	-	-	
	Turbidity (NTU)	-	-	No data. Slight evidence of sedimentation.
	Electrical conductivity (mS/m)	-	18.0	A category
Response variables	Chl a: periphyton	-	21.25	Some nutrient elevations shown by periphyton data (C/D category; n=1)
	Chl a: phytoplankton	-	0.18	
	Macroinvertebrates	ASPT [§] : mean of 6.58	ASPT: 7.3 MIRAI ^{**} : 86.1%	B category for the present state.
	Fish community score	-	FRAI [*] : 86.1%	B category for the present state.
	Diatoms	-	SPI [#] =17.6	High quality water
Toxics		-	-	No data, but some impact expected due to farming-related pesticides and fertilizer use.
OVERALL SITE CLASSIFICATION		B (PAI model)		

*: boundary value recalibrated

*: FRAI = Fish Response Assessment Index

-: no data

**: MIRAI = Macro Invertebrate Response Assessment Index

#: Specific Pollution Index

§: ASPT = Average Score Per Taxon

Table 5.38: Water quality Ecospecs for Diep River (WQSU 3, K40A)

River: Diep		EWR Site: 3	Monitoring site: K4H003Q01
Water quality metrics		ECOSPEC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be ≤ 16 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be ≤ 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be ≤ 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be ≤ 21 mg/L.	
	NaCl	The 95 th percentile of the data must be ≤ 191 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be ≤ 351 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be ≤ 30 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must range from 4.5 to 7.5.	
	Temperature	Small deviation from the natural temperature range.	
	Dissolved oxygen	The 5 th percentile of the data must be ≥ 7.5 mg/L.	
	Turbidity	Vary by a small amount from the natural turbidity range; minor silting of instream habitats acceptable.	
Nutrients	TIN	The 50 th percentile of the data must be ≤ 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be ≤ 0.025 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be <15 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be ≤ 52.5 mg/m ² .**	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

* To be generated using TEACHA when the TPC for EC is exceeded or salt pollution expected.

** Periphyton (21.25 mg/m²) is actually in a C/D category (C = 12 - 21 and D = 21 - 84 mg/m²; DWAF, 2008), so have defined the upper boundary of a C/D as the EcoSpec.

Table 5.39: Water quality TPC's for Diep River (WQSU 3, K40A)

River: Diep		EWR Site: 3	Monitoring site: K4H003Q01
Water quality metrics		TPC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be 13 – 16 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be 16 – 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be 12 – 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be 17 – 21 mg/L.	
	NaCl	The 95 th percentile of the data must be 36 – 45 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be 153 – 191 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be 24 – 30 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must be <4.7 and >7.3.	
	Temperature	Small to moderate deviation from the natural temperature range. Some highly temperature sensitive species in lower abundances and frequency of occurrence than expected for reference.	
	Dissolved oxygen	The 5 th percentile of the data must be 7.8 – 7.5 mg/L.	
	Turbidity	Moderate changes to the catchment land-use resulting in <u>temporary</u> unnaturally high sediment loads and high turbidities.	
Nutrients	TIN	The 50 th percentile of the data must be 0.2 – 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be 0.02 – 0.025 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be 12 – 15 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be 42 – 52.5 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

Table 5.40: PES categories and overall site assessment for the Karatara River at WQSU 5

RIVER	Karatara River		WATER QUALITY MONITORING POINTS	
WQSU	WQSU 5 (source to Swartvlei)		RC	Karatara R. @ Karatara Forest Reserve K4H002Q01 ('76 – '79; n = 115)
EWR SITE	4		PES	K4H002Q01 ('03 – '07; n = 36)
Confidence assessment		Moderate as adequate data to assess reference and present states		
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄	7.62	12.63	TEACHA was used for data assessment. Salts = an A category.
	Na ₂ SO ₄	2.05	3.01	
	MgCl ₂	4.43	4.16	
	CaCl ₂	9.89	9.16	
	NaCl	35.59	36.15	
	CaSO ₄	0.73	0.73	
Nutrients (mg/L)	SRP	0.022 *	0.047	B category.
	TIN	0.07	0.128	A category.
Physical Variables	pH (5 th + 95 th %ile)	3.7 + 7.7	3.8 + 6.6	A category.
	Temperature	-	-	No data but no impacts expected
	Dissolved oxygen	-	-	
	Turbidity (NTU)	-	-	No data. Slight evidence of sedimentation.
	Electrical conductivity (mS/m)	-	7.0	A category.
Response variables	Chl a: periphyton	-	9.91	B category for periphyton.
	Chl a: phytoplankton	-	0.09	
	Macroinvertebrates	ASPT: 7.3	ASPT: 8.1 MIRAI: 92.3%	A/B category for the present state.
	Fish community score	-	FRAI: 82.4%	B category for the present state.
	Diatoms	-	SPI=19.9	High quality water
Toxics		-	-	No data, but some impact from timber processing at Geelhoutvlei.
OVERALL SITE CLASSIFICATION		B for WQSU 5 , although an A category for upstream river stretch (PAI model)		

*: boundary value recalibrated

-: no data

Table 5.41: Water quality Ecospecs for Karatara River at WQSU 5

River: Karatara		EWR Site: 4	Monitoring site: K4H002Q01
Water quality metrics		ECOSPEC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be ≤ 16 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be ≤ 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be ≤ 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be ≤ 21 mg/L.	
	NaCl	The 95 th percentile of the data must be ≤ 191 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be ≤ 351 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be ≤ 30 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must range from 4.5 to 7.5.	
	Temperature	Natural temperature range.	
	Dissolved oxygen	The 5 th percentile of the data must be ≥ 8.0 mg/L.	
	Turbidity	Vary by a small amount from the natural turbidity range; minor silting of instream habitats acceptable.	
Nutrients	TIN	The 50 th percentile of the data must be ≤ 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be ≤ 0.075 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be < 10 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be ≤ 12 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

* To be generated using TEACHA when the TPC for EC is exceeded or salt pollution expected.

Table 5.42: Water quality TPC's for Karatara River at WQSU 5

River: Karatara		EWR Site: 4	Monitoring site: K4H002Q01
Water quality metrics		TPC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be 13 – 16 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be 16 – 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be 12 – 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be 17 – 21 mg/L.	
	NaCl	The 95 th percentile of the data must be 36 – 45 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be 153 – 191 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be 24 – 30 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must be <4.7 and >7.3.	
	Temperature	Small deviation from the natural temperature range.	
	Dissolved oxygen	The 5 th percentile of the data must be 8.2 – 8.0 mg/L.	
	Turbidity	Small to moderate changes to the catchment land-use resulting in minor and temporary effects of silting of habitats.	
Nutrients	TIN	The 50 th percentile of the data must be 0.2 – 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be 0.06 – 0.075 mg/L.**	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be 8 – 10 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be 10 – 12 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

* To be generated using TEACHA when the TPC for EC is exceeded or salt pollution expected.

** Although the upper boundary for the relevant phosphate category is 0.125 mg/L, the TPC has been set at 0.075 mg/L as PES levels for phosphate were 0.046 mg/L.

Table 5.43: PES categories and overall site assessment for the Hoëkraal River at WQSU 4

RIVER	Hoëkraal River		WATER QUALITY MONITORING POINTS	
WQSU	WQSU 4 (source to beginning of lower foothills)		RC	Hoëkraal R. @ Eastbrook K4H001Q01 ('77 – '80; n = 83)
EWR SITE	-		PES	K4H001Q01 ('03 – '07; n = 28)
Confidence assessment	Low - Moderate as limited data for the present state assessment. Lower section in the backup zone from Swartvlei lake. Note that the gauging weir is located on the lower Hoëkraal River.			
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄	35.87	153.20	TEACHA was used for the analysis of WMS data. Results should be disregarded as data shows salt intrusion from the saline Swartvlei lake.
	Na ₂ SO ₄	9.95	159.5	
	MgCl ₂	35.92	100.33	
	CaCl ₂	24.80	45.24	
	NaCl	399.69	1560.41	
	CaSO ₄	0.59	0.73	
Nutrients (mg/L)	SRP	0.014 *	0.034	B category.
	TIN	0.06	0.088	A category.
Physical Variables	pH (5 th + 95 th %ile)	4.4 + 7.2	4.5 + 7.8	A category.
	Temperature	-	-	No data but no impacts expected, although the river is deep and wide above the lower site, which may result in lowering oxygen levels.
	Dissolved oxygen	-	-	
	Turbidity (NTU)	-	-	No data, but some evidence of sedimentation.
	Electrical conductivity (mS/m)		4.2: Upper site	A category
Response variables	Chl a: periphyton	-	4.81: Upper site 152.93: Lower site	Lower site shows some nutrient build-up in the water.
	Chl a: phytoplankton	-	0.14: Upper site 0.47: Lower site	
	Macroinvertebrates	-	-	Not assessed as not an EWR site.
	Fish community score	-	-	
	Diatoms	-	SPI=19.8: Upper river. SPI=16.2: Lower river.	Upper site: High quality water. Lower site: Good quality water.
Toxics		-	-	No data, but no impacts expected.
OVERALL SITE CLASSIFICATION		C Category (PAI model)		

* boundary value recalibrates

Table 5.44: Water quality Ecospecs for Hoëkraal River at WQSU 4

River: Hoëkraal		EWR Site: -	Monitoring site: K4H001Q01, although located in the lower section of the river and salt water intrusion detected due to back-up from Swartvlei Lake.
Water quality metrics		ECOSPEC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be ≤ 16 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be ≤ 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be ≤ 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be ≤ 21 mg/L.	
	NaCl	The 95 th percentile of the data must be ≤ 191 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be ≤ 351 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be ≤ 30 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must range from 4.5 to 7.5.	
	Temperature	Small deviation from the natural temperature range.	
	Dissolved oxygen	The 5 th percentile of the data must be ≥ 7.5 mg/L.	
	Turbidity	Vary by a small amount from the natural turbidity range; minor silting of instream habitats acceptable.	
Nutrients	TIN	The 50 th percentile of the data must be ≤ 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be ≤ 0.125 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be <10 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be ≤ 12 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

* To be generated using TEACHA when the TPC for EC is exceeded or salt pollution expected.

Table 5.45: Water quality TPC's for Hoëkraal River at WQSU 4

River: Hoekraal		EWR Site: -	Monitoring site: K4H001Q01
Water quality metrics		TPC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be 13 – 16 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be 16 – 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be 12 – 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be 17 – 21 mg/L.	
	NaCl	The 95 th percentile of the data must be 36 – 45 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be 153 – 191 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be 24 – 30 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must be <4.7 and >7.3.	
	Temperature	Small to moderate changes to the catchment land-use resulting in minor and temporary effects of silting of habitats.	
	Dissolved oxygen	The 5 th percentile of the data must be 7.8 – 7.5 mg/L.	
	Turbidity	Moderate changes to the catchment land-use resulting in <u>temporary</u> unnaturally high sediment loads and high turbidities.	
Nutrients	TIN	The 50 th percentile of the data must be 0.2 – 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be 0.1 – 0.125 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be 8 – 10 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be 10 – 12 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

Table 5.46: PES categories and overall site assessment for Goukamma River at WQSU 1

RIVER	Goukamma River		WATER QUALITY MONITORING POINTS	
WQSU	WQSU 1 (source to the end of the mountainous area)		RC	Rooi R. @ George K3H002-Q01 ('77 – '79 n = 84)
EWR SITE	None		PES	K3H002-Q01 ('01 – '06 n = 65)
Confidence assessment		Low to medium because no DWAF monitoring station. However unimpacted catchment and likely to be similar to other mountain streams.		
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄			No data
	Na ₂ SO ₄			
	MgCl ₂			
	CaCl ₂			
	NaCl			
	CaSO ₄			
Nutrients (mg/L)	SRP	0.014	≤0.014*	A category.
	TIN	0.04	≤0.25*	A category.
Physical Variables	pH (5 th – 95 th %)		3.7 – 6.7	Naturally acidic.
	Temperature			No data but expected to be natural as little development in upstream catchment.
	Dissolved oxygen			
	Turbidity (NTU)			
	Electrical conductivity (mS/m)	16	17	A category.
Response variables	Chl a: periphyton			No data
	Chl a: phytoplankton			No data
	Macroinvertebrates			No data
	Fish community score			No data
Toxics				No data.
OVERALL SITE CLASSIFICATION		A		

Table 5.47: Water quality Ecospecs and TPCs for the Goukamma River at WQSU 2

RIVER	WATER QUALITY MONITORING POINTS					
	Goukamma River	DWAF WQ WMS	RHP			
WQSU	2	None	Currently a RHP site higher up in catchment			
EWRS SITE	none					
Confidence in PES assessment	Low because extrapolated from another catchment (Karataara R) and WQSU (Goukamma WQSU 2). Biological data (from current project and RHP upper catchment) support inferred PES for water quality.					
Water Quality Constituents	PES Category	WQ Eco specs	Improvement required?			
			TPC			
			Monitoring frequency			
Inorganic salts (mg/L)	MgSO ₄	≤16 mg/L	95 th percentile to be < 16 mg/L	Every 2 months		
	Na ₂ SO ₄	≤20 mg/L	95 th percentile to be < 20 mg/L			
	MgCl ₂	≤15 mg/L	95 th percentile to be < 15 mg/L			
	CaCl ₂	≤21 mg/L	95 th percentile to be < 21 mg/L			
Nutrients (mg/L)	NaCl	≤45 mg/L	95 th percentile to be < 45 mg/L	Every 2 months		
	SRP	≤0.012 mg/L	Yes - B/C			
	TIN	≤0.25 mg/L	No			
Physical Variables	pH	< 6.4	No	95 th percentile to be < 6.4	Every 2 months	
	Temperature	Natural range	N/A	Natural temp. range	Every 2 months	
	Dissolved oxygen	No data. Visual inspection did not reveal a turbidity problem.	N/A	N/A	5 th percentile to be > 8 mg/L	Every 2 months
	Turbidity (NTU)		N/A	N/A	No change allowed	Every 2 months
	Electrical conductivity (mS/m)	A – Stable	≤30 mS/m	No	95 th percentile to be < 30 mS/m	Every 2 months
	Chl a: periphyton	No data. Visual inspection did not reveal a problem.	≤ 1.7 mg/m ² (A category) ≤ 10 µg/L (B category)	N/A	50 th percentile to be < 1.7 mg/m ² 50 th percentile to be < 10 µg/L	Quarterly
Response variables	Chl a: phytoplankton					
	Macroinvertebrates (ASPT)	B (this study)				
	Fish community score	C (this study)				
Toxics	Instream toxicity	No data				
		No data.				
See Ecospecs for fish and invertebrates respectively						
Unlikely to be a problem. Assess only if the biomonitoring results indicate there is a serious problem and the cause is unknown.						

Table 5.48: PES categories and overall site assessment for Goukamma River at WQSU 2

RIVER	Goukamma River		WATER QUALITY MONITORING POINTS		
WQSU	WQSU 2 (edge of mountainous/forested area to the upper limit of tidal influence - at approximately the N2 Bridge)		RC	Karatara R. @ Karatara State Forest K4H002-Q01 ('71 – '76; n = 107)	
EWR SITE	Gou 1		PES	K4H002-Q01 ('01 – '06; n = 51)	
Confidence assessment	Low, the monitoring station is in an adjacent catchment.				
Water Quality Constituents		RC Value	PES Value	Category/Comment	G-power (Confidence)
Inorganic salts (mg/L)	MgSO ₄			No data	
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
Nutrients (mg/L)	SRP	0.018	0.048*	Category = C. Trend = increasing	0.232 (Low)
	TIN	0.11	0.11	Category = A. Trend = stable	
Physical Variables	pH (5 th – 95 th %)		3.8 – 6.4	Naturally acidic waters	0.781 (Medium)
	Temperature			No data. Unlikely to be a problem	
	Dissolved oxygen				
	Turbidity (NTU)			No data. Visual inspection = low turbidity	
	Electrical conductivity (mS/m)	10	12	Category = A. Trend = stable	0.997 (High)
Response variables	Chl a: periphyton			No data. Visual inspection = no excessive algal growth	
	Chl a: phytoplankton				
	Macroinvertebrates			B category; ASPT = 7.2, SASS = 143 (this study). Natural (RHP)	
	Fish community score			C (this study)	
Toxics				No data, but unlikely to be a problem	
OVERALL SITE CLASSIFICATION		A Category (PAI model)			

Table 5.49: Water quality Ecospecs and TPCs for the Goukamma River at WQSU 2

RIVER	WATER QUALITY MONITORING POINTS					
	Goukamma River	DWAF WQ WMS				
WQSU	2	None				
EWRS SITE	Gou 1	Currently a RHP site higher up in catchment				
Confidence in PES assessment	Low - medium because extrapolated from another catchment (Karataru R). Biological data (from current project and RHP upper catchment) support inferred PES for water quality.					
Water Quality Constituents	PES Category	WQ Eco specs	Improvement required?	TPC	Monitoring frequency	
Inorganic salts (mg/L)	MgSO ₄	≤16 mg/L		95 th percentile to be < 16 mg/L		
	Na ₂ SO ₄	≤20 mg/L		95 th percentile to be < 20 mg/L		
	MgCl ₂	≤15 mg/L	N/A	95 th percentile to be < 15 mg/L	Every 2 months	
	CaCl ₂	≤21 mg/L		95 th percentile to be < 21 mg/L		
Nutrients (mg/L)	NaCl	≤45 mg/L		95 th percentile to be < 45 mg/L		
	SRP	≤0.025 mg/L	Yes - B/C	50 th percentile to be < 0.025 mg/L	Every 2 months	
Physical Variables	TIN	≤0.25 mg/L	No	50 th percentile to be < 0.25 mg/L	Every 2 months	
	pH	< 6.4	No	95 th percentile to be < 6.4	Every 2 months	
	Temperature	Natural range	N/A	Natural temp. range	Every 2 months	
	Dissolved oxygen	No data. Visual inspection did not reveal a turbidity problem.	N/A	8 mg/L	5 th percentile to be > 8 mg/L	Every 2 months
	Turbidity (NTU)		N/A	No change	No change allowed	Every 2 months
	Electrical conductivity (mS/m)	A – Stable	≤30 mS/m	No	95 th percentile to be < 30 mS/m	Every 2 months
	Chl a: periphyton	No data. Visual inspection did not reveal a problem.	≤ 1.7 mg/m ² (A category) ≤ 10 µg/L (B category)	N/A	50 th percentile to be < 1.7 mg/m ² 50 th percentile to be < 10 µg/L	Quarterly
Response variables	Chl a: phytoplankton					
	Macroinvertebrates (ASPT)	B (this study)				
	Fish community score	C (this study)				
Toxics	Instream toxicity	No data				
		No data.				
See Ecospecs for fish and invertebrates respectively						
Unlikely to be a problem. Assess only if the biomonitoring results indicate there is a serious problem and the cause is unknown.						

Table 5.50: PES categories and overall site assessment for the Knysna River at WQSU 1

RIVER	Knysna River	WATER QUALITY MONITORING POINTS		
WQSU	WQSU 1	RC	Knysna R. @ Millwood K5H002Q01 ('77 – '80; n = 75)	
EWR SITE	1	PES	K5H002Q01 ('04 – '07; n = 26)	
Confidence assessment	Moderate as adequate data to assess reference and present states			
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄	7.18	7.19	TEACHA was used for data assessment. Salts = A category
	Na ₂ SO ₄	1.49	4.20	
	MgCl ₂	2.60	2.73	
	CaCl ₂	11.50	3.92	
	NaCl	50.83*	39.54	
	CaSO ₄	0.53	0.38	
Nutrients (mg/L)	SRP	0.011*	0.021	B category
	TIN	0.06	0.112	A category
Physical Variables	pH (5 th + 95 th %ile)	4.0 + 6.9	4.5 + 7.2	A category
	Temperature	-	-	No data but no impacts expected, although less shading may result in a slight increase in instream temperature.
	Dissolved oxygen	-	-	No data. Some evidence of slight sedimentation seen.
	Turbidity (NTU)	-	-	No data. Some evidence of slight sedimentation seen.
	Electrical conductivity (mS/m)		9.0	A category.
Response variables	Chl a: periphyton	-	4.08	Some nutrient elevation indicated by periphyton values (B category; n=1).
	Chl a: phytoplankton	-	0.12	
	Macroinvertebrates	ASPT: mean of 8.1	ASPT: 6.9 MIRAI: 86.92%	B category for the present state.
	Fish community score	-	FRAI: 86.4%	B category for the present state.
	Diatoms	-	SPI=18.9	High quality water.
Toxics		-	-	No data but no impacts expected.
OVERALL SITE CLASSIFICATION		A/B for WQSU 1 , although an A category for EWR 1 and upstream of the site (PAI model)		

Table 5.51: Water quality Ecospecs for Knysna River at WQSU 1

River: Knysna		EWR: 1	Monitoring site: K5H002Q01
Water quality metrics		ECOSPEC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be ≤ 16 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be ≤ 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be ≤ 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be ≤ 21 mg/L.	
	NaCl	The 95 th percentile of the data must be ≤ 45 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be ≤ 351 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be ≤ 30 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must range from 4.5 to 7.5.	
	Temperature	Natural temperature range.	
	Dissolved oxygen	The 5 th percentile of the data must be ≥ 8.0 mg/L.	
	Turbidity	Vary by a small amount from the natural turbidity range; minor silting of instream habitats acceptable.	
Nutrients	TIN	The 50 th percentile of the data must be ≤ 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be ≤ 0.025 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be <15 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be ≤ 12 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

* To be generated using TEACHA when the TPC for EC is exceeded or salt pollution expected.

Table 5.52: Water quality TPC's for Knysna River at WQSU 1

River: Knysna		EWR: 1	Monitoring site: K5H002Q01
Water quality metrics		TPC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be 13 – 16 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be 16 – 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be 12 – 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be 17 – 21 mg/L.	
	NaCl	The 95 th percentile of the data must be 36 – 45 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be 280 – 351 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be 24 – 30 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must be <4.7 and >7.3.	
	Temperature	Small deviation from the natural temperature range.	
	Dissolved oxygen	The 5 th percentile of the data must be 8.2 – 8.0 mg/L.	
	Turbidity	Moderate changes to the catchment land-use resulting in <u>temporary</u> unnaturally high sediment loads and high turbidities.	
Nutrients	TIN	The 50 th percentile of the data must be 0.2 – 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be 0.02 – 0.025 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be 12 – 15 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be 10 – 12 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

* To be generated using TEACHA when the TPC for EC is exceeded or salt pollution expected.

Table 5.53: PES categories and overall site assessment for the Gouna River at WQSU 2

RIVER	Gouna River	WATER QUALITY MONITORING POINTS		
WQSU	WQSU 2	RC	Gouna R. @ Gouna Commonage K5H001Q01 ('77 - '80; n = 76)	
EWR SITE	2	PES	K5H001Q01 ('81 - '84; n = 30)	
Confidence assessment	Low confidence due to limited data, particularly for present state			
Water Quality Constituents		RC Value	PES Value	Category/Comment
Inorganic salts (mg/L)	MgSO ₄	14.45	23.19	TEACHA was used for data assessment. Slight elevation in salts seen, i.e. an A/B category
	Na ₂ SO ₄	3.78	0.30	
	MgCl ₂	10.75	12.11	
	CaCl ₂	10.59	9.35	
	NaCl	95.29*	102.20	
	CaSO ₄	0.54	0.72	
Nutrients (mg/L)	SRP	0.009*	0.011	A category.
	TIN	0.070	0.120	A category.
Physical Variables	pH (5 th + 95 th %ile)	4.0 + 6.8	4.0 + 5.6	A category as little change from natural.
	Temperature	-	-	No data but no impacts expected.
	Dissolved oxygen	-	-	
	Turbidity (NTU)	-	-	No data. Sedimentation seen in lower section of the WQSU, i.e. below the pump station.
	Electrical conductivity (mS/m)		15.0	A category.
Response variables	Chl a: periphyton	-	43.70	Nutrient elevations indicated by periphyton value (D category; n=1).
	Chl a: phytoplankton	-	0.09	
	Macroinvertebrates	ASPT: 6.9 – 7.6	ASPT: 7.6 MIRAI: 92.8%	A category for the present state.
	Fish community score	-	FRAI: 93.8%	A category for the present state.
	Diatoms	-	SPI=19.8	High quality water.
Toxics		-	-	No data but no toxics expected.
OVERALL SITE CLASSIFICATION		B for WQSU 2 , although an A category for river stretch upstream of the pump station (PAI model)		

Table 5.54: Water quality EcoSpecs for Gouna River at WQSU 2

River: Gouna		EWR: 2	Monitoring site: K5H001Q01
Water quality metrics		ECOSPEC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be ≤ 23 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be ≤ 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be ≤ 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be ≤ 21 mg/L.	
	NaCl	The 95 th percentile of the data must be ≤ 191 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be ≤ 351 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be ≤ 43 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must range from 4.5 to 7.5.	
	Temperature	Largely natural to small deviation from the natural temperature range.	
	Dissolved oxygen	The 5 th percentile of the data must be ≥ 7.5 mg/L.	
	Turbidity	Vary by a small amount from the natural turbidity range; minor silting of instream habitats acceptable.	
Nutrients	TIN	The 50 th percentile of the data must be ≤ 0.15 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be ≤ 0.025 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be <15 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be ≤ 84 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

* To be generated using TEACHA when the TPC for EC is exceeded or salt pollution expected.

Table 5.55: Water quality TPC's for Gouna River at WQSU 2

River: Gouna		EWR: 2	Monitoring site: K5H001Q01
Water quality metrics		TPC	
Inorganic salts*	MgSO ₄	The 95 th percentile of the data must be 18 – 23 mg/L.	
	Na ₂ SO ₄	The 95 th percentile of the data must be 16 – 20 mg/L.	
	MgCl ₂	The 95 th percentile of the data must be 12 – 15 mg/L.	
	CaCl ₂	The 95 th percentile of the data must be 17 – 21 mg/L.	
	NaCl	The 95 th percentile of the data must be 36 – 45 mg/L.	
	CaSO ₄	The 95 th percentile of the data must be 153 – 191 mg/L.	
Physical variables	EC	The 95 th percentile of the data must be 35 – 43 mS/m.	
	pH	The 5 th and 95 th percentiles of the data must be <4.7 and >7.3.	
	Temperature	Small to moderate deviation from the natural temperature range. Some highly temperature sensitive species in lower abundances and frequency of occurrence than expected for reference.	
	Dissolved oxygen	The 5 th percentile of the data must be 7.8 – 7.5 mg/L.	
	Turbidity	Moderate changes to the catchment land-use resulting in <u>temporary</u> unnaturally high sediment loads and high turbidities.	
Nutrients	TIN	The 50 th percentile of the data must be 0.2 – 0.25 mg/L.	
	PO ₄ -P	The 50 th percentile of the data must be 0.012 – 0.015 mg/L.	
Response variables	Chl-a phytoplankton	The 50 th percentile of the data must be 12 – 15 µg/L.	
	Chl-a periphyton	The 50 th percentile of the data must be 67 – 84 mg/m ² .	
Toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996).		

6. GROUNDWATER - QUANTITY COMPONENT

The groundwater quantity component was determined using values (recharge and groundwater baseflow) obtained during the determination of water resource classes and associated resource quality objectives in the Breede-Gouritz WMA, (DWS 2017), shown in Table 6.1.

Population values were obtained from the Water Services dataset of 2011 for the Breede catchment and from the Gouritz Catchment Intermediate Reserve study for the Gouritz catchment. The Basic Human Needs Reserve provides for the essential needs of individuals served by the water resource in question and includes water for drinking, food preparation and for personal hygiene. A life-line amount of 25 litres per person per day was used.

Table 6.1: Breede-Gouritz Groundwater Reserve

Quaternary Catchment	Recharge (Mm ³ /a)	Population	Basic Human Needs (Mm ³ /a)	Groundwater Baseflow (Mm ³ /a)	Reserve (Mm ³ /a)	Reserve as % of Recharge
G40A	13.06	15963	0.15	3.17	3.32	25.39
G40B	19.19	4612	0.04	5.33	5.37	27.99
G40C	45.16	38379	0.35	6.25	6.60	14.62
G40D	59.72	15963	0.15	14.45	14.60	24.44
G40E	13.19	7670	0.07	4.41	4.48	33.97
G40F	11.28	28422	0.26	2.12	2.38	21.09
G40G	16.02	17112	0.16	3.72	3.88	24.20
G40H	6.53	32767	0.30	1.58	1.88	28.77
G40J	6.92	1574	0.01	2.53	2.54	36.77
G40K	9.13	15963	0.15	4.67	4.82	52.75
G40L	13.96	15963	0.15	1.63	1.78	12.72
G40M	10.57	355	0.003	5.17	5.17	48.94
G50A	7.37	1370	0.01	2.61	2.62	35.58
G50B	6.59	1507	0.01	3.47	3.48	52.86
G50C	8.56	1748	0.02	2.05	2.07	24.13
G50D	5.39	9430	0.09	2.55	2.64	48.91
G50E	4.92	14355	0.13	1.37	1.50	30.51
G50F	6.64	1989	0.02	1.27	1.29	19.40
G50G	2.40	844	0.01	1.43	1.44	59.90
G50H	5.75	736	0.01	3.28	3.29	57.16
G50J	6.07	3325	0.03	1.90	1.93	31.80
G50K	2.72	1748	0.02	0.76	0.78	28.53
H10A	13.15	12494	0.11	0.76	0.87	6.65
H10B	12.20	12494	0.11	0.48	0.59	4.87
H10C	21.28	57300	0.52	2.00	2.52	11.86
H10D	14.89	12494	0.11	2.05	2.16	14.53
H10E	20.35	12494	0.11	3.20	3.31	16.29
H10F	25.24	20720	0.19	1.39	1.58	6.26
H10G	31.82	12494	0.11	0.44	0.55	1.74
H10H	28.48	12494	0.11	2.80	2.91	10.23
H10J	61.45	1035	0.01	7.94	7.95	12.94
H10K	43.17	12494	0.11	7.40	7.51	17.41
H10L	2.76	4268	0.04	0.00	0.04	1.41

Quaternary Catchment	Recharge (Mm ³ /a)	Population	Basic Human Needs (Mm ³ /a)	Groundwater Baseflow (Mm ³ /a)	Reserve (Mm ³ /a)	Reserve as % of Recharge
H20A	2.42	427	0.00	0.47	0.47	19.58
H20B	5.37	17136	0.16	0.17	0.33	6.08
H20C	2.84	1266	0.01	0.05	0.06	2.17
H20D	8.74	1266	0.01	2.11	2.12	24.27
H20E	14.68	1266	0.01	2.01	2.02	13.77
H20F	8.65	875	0.01	0.32	0.33	3.79
H20G	4.83	1266	0.01	0.47	0.48	9.97
H20H	1.56	140420	1.28	0.07	1.35	86.62
H30A	5.17	1102	0.01	0.33	0.34	6.58
H30B	6.04	39573	0.36	0.16	0.52	8.63
H30C	10.59	1317	0.01	0.07	0.08	0.77
H30D	3.18	926	0.01	0.06	0.07	2.15
H30E	2.95	9784	0.09	0.31	0.40	13.53
H40A	3.74	2233	0.02	0.87	0.89	23.81
H40B	12.26	2152	0.02	0.87	0.89	7.26
H40C	4.90	2233	0.02	0.86	0.88	17.97
H40D	4.18	2233	0.02	1.85	1.87	44.75
H40E	10.91	2233	0.02	0.20	0.22	2.02
H40F	1.07	1798	0.02	0.58	0.60	55.74
H40G	3.22	2233	0.02	0.23	0.25	7.78
H40H	4.71	1217	0.01	0.13	0.14	3.00
H40J	4.44	26455	0.24	0.18	0.42	9.49
H40K	2.99	3916	0.04	0.24	0.28	9.22
H40L	2.47	2290	0.02	0.42	0.44	17.85
H50A	1.42	3842	0.04	0.26	0.30	20.78
H50B	5.04	5825	0.05	0.78	0.83	16.53
H60A	30.87	10083	0.09	2.49	2.58	8.36
H60B	42.43	7900	0.07	7.28	7.35	17.33
H60C	30.89	15284	0.14	1.64	1.78	5.76
H60D	14.76	511	0.00	0.95	0.95	6.47
H60E	9.73	10305	0.09	0.71	0.80	8.26
H60F	7.65	3321	0.03	0.66	0.69	9.02
H60G	4.11	10083	0.09	0.64	0.73	17.81
H60H	7.49	10083	0.09	1.14	1.23	16.45
H60J	8.17	10083	0.09	1.31	1.40	17.16
H60K	3.59	10083	0.09	1.04	1.13	31.53
H60L	2.88	10083	0.09	0.87	0.96	33.40
H70A	5.55	4786	0.04	1.47	1.51	27.27
H70B	22.83	19350	0.18	4.17	4.35	19.04
H70C	3.99	4915	0.04	0.23	0.27	6.89
H70D	20.70	4786	0.04	5.53	5.57	26.93
H70E	26.55	6729	0.06	5.16	5.22	19.67
H70F	15.50	2721	0.02	2.31	2.33	15.06

Quaternary Catchment	Recharge (Mm ³ /a)	Population	Basic Human Needs (Mm ³ /a)	Groundwater Baseflow (Mm ³ /a)	Reserve (Mm ³ /a)	Reserve as % of Recharge
H70G	3.92	4786	0.04	1.26	1.30	33.26
H70H	2.80	746	0.01	1.89	1.90	67.74
H70J	3.95	4786	0.04	1.43	1.47	37.31
H70K	3.03	772	0.01	1.21	1.22	40.17
H80A	16.34	0	0.00	7.21	7.21	44.12
H80B	24.01	0	0.00	6.45	6.45	26.86
H80C	5.75	10 110	0.09	0.61	0.70	12.21
H80D	2.57	0	0.00	1.23	1.23	47.86
H80E	7.66	392	0.00	2.11	2.11	27.59
H80F	5.96	585	0.01	2.72	2.73	45.73
H90A	19.62	0	0.00	9.04	9.04	46.08
H90B	12.96	0	0.00	6.02	6.02	46.45
H90C	5.51	18 526	0.17	1.93	2.10	38.10
H90D	10.38	1 697	0.02	3.29	3.31	31.84
H90E	9.70	6 253	0.06	4.88	4.94	50.90
J11A	2.98	0	0.00	0.00	0.00	0.00
J11B	3.11	0	0.00	0.00	0.00	0.00
J11C	0.22	192	0.00	0.00	0.00	0.80
J11D	3.74	48	0.00	0.00	0.00	0.01
J11E	1.40	4 773	0.04	0.00	0.04	3.11
J11F	0.43	734	0.01	0.00	0.01	1.56
J11G	0.12	84	0.00	0.00	0.00	0.64
J11H	4.01	885	0.01	0.00	0.01	0.20
J11J	6.02	1 333	0.01	0.00	0.01	0.20
J11K	2.52	11 732	0.11	0.00	0.11	4.25
J12A	3.15	0	0.00	0.02	0.02	0.63
J12B	1.55	7 857	0.07	0.00	0.07	4.63
J12C	1.59	152	0.00	0.01	0.01	0.72
J12D	6.32	535	0.00	0.02	0.02	0.39
J12E	1.93	507	0.00	0.02	0.02	1.28
J12F	6.15	336	0.00	0.03	0.03	0.54
J12G	5.66	294	0.00	0.01	0.01	0.22
J12H	4.53	0	0.00	0.02	0.02	0.44
J12J	4.59	0	0.00	0.01	0.01	0.22
J12K	2.44	0	0.00	0.01	0.01	0.41
J12L	6.59	681	0.01	0.05	0.06	0.85
J12M	3.04	0	0.00	0.06	0.06	1.97
J13A	4.10	0	0.00	0.02	0.02	0.49
J13B	2.86	660	0.01	0.03	0.04	1.26
J13C	2.91	0	0.00	0.03	0.03	1.03
J21A	4.28	34 661	0.32	0.00	0.32	7.39
J21B	0.56	389	0.00	0.00	0.00	0.63
J21C	0.12	406	0.00	0.00	0.00	3.09

Quaternary Catchment	Recharge (Mm ³ /a)	Population	Basic Human Needs (Mm ³ /a)	Groundwater Baseflow (Mm ³ /a)	Reserve (Mm ³ /a)	Reserve as % of Recharge
J21D	0.24	0	0.00	0.00	0.00	0.00
J21E	0.26	0	0.00	0.00	0.00	0.00
J22A	3.04	0	0.00	0.00	0.00	0.00
J22B	1.12	0	0.00	0.00	0.00	0.00
J22C	1.27	241	0.00	0.00	0.00	0.17
J22D	1.22	133	0.00	0.00	0.00	0.10
J22E	1.31	0	0.00	0.00	0.00	0.00
J22F	0.12	1 237	0.01	0.00	0.01	9.41
J22G	2.92	0	0.00	0.00	0.00	0.00
J22H	4.19	919	0.01	0.00	0.01	0.20
J22J	0.90	0	0.00	0.00	0.00	0.00
J22K	0.35	0	0.00	0.00	0.00	0.00
J23A	0.28	2 080	0.02	0.00	0.02	6.78
J23B	0.50	47	0.00	0.00	0.00	0.09
J23C	0.25	12	0.00	0.00	0.00	0.04
J23D	0.70	12	0.00	0.00	0.00	0.02
J23E	2.03	674	0.01	0.18	0.19	9.17
J23F	1.33	6 901	0.06	0.00	0.06	4.73
J23G	0.00	0	0.00	0.00	0.00	0.00
J23H	1.11	0	0.00	0.00	0.00	0.00
J23J	1.82	0	0.00	0.97	0.97	53.30
J24A	2.58	193	0.00	0.00	0.00	0.07
J24B	0.51	1 521	0.01	0.00	0.01	2.72
J24C	0.21	0	0.00	0.00	0.00	0.00
J24D	0.08	314	0.00	0.00	0.00	3.58
J24E	0.39	0	0.00	0.00	0.00	0.00
J24F	1.37	0	0.00	0.00	0.00	0.00
J25A	2.42	200	0.00	1.02	1.02	42.22
J25B	4.45	4 135	0.04	1.23	1.27	28.49
J25C	1.04	1 342	0.01	0.02	0.03	3.10
J25D	2.94	5 170	0.05	0.61	0.66	22.35
J25E	1.12	402	0.00	0.04	0.04	3.90
J31A	7.88	0	0.00	1.13	1.13	14.34
J31B	1.57	0	0.00	0.48	0.48	30.57
J31C	1.87	0	0.00	0.35	0.35	18.72
J31D	2.07	266	0.00	0.38	0.38	18.47
J32A	0.08	50	0.00	0.00	0.00	0.57
J32B	0.01	881	0.01	0.00	0.01	80.39
J32C	0.01	77	0.00	0.00	0.00	7.03
J32D	0.00	0	0.00	0.00	0.00	0.00
J32E	1.76	719	0.01	0.00	0.01	0.37
J33A	4.81	130	0.00	1.44	1.44	29.96
J33B	8.98	680	0.01	1.47	1.48	16.44

Quaternary Catchment	Recharge (Mm ³ /a)	Population	Basic Human Needs (Mm ³ /a)	Groundwater Baseflow (Mm ³ /a)	Reserve (Mm ³ /a)	Reserve as % of Recharge
J33C	2.83	76	0.00	0.01	0.01	0.38
J33D	3.82	255	0.00	1.24	1.24	32.52
J33E	8.22	13 522	0.12	1.98	2.10	25.59
J33F	4.50	34 154	0.31	2.19	2.50	55.59
J34A	3.08	7 465	0.07	1.48	1.55	50.26
J34B	6.44	1 850	0.02	2.85	2.87	44.52
J34C	9.60	669	0.01	3.51	3.52	36.63
J34D	4.06	1 915	0.02	1.80	1.82	44.77
J34E	2.29	1 321	0.01	1.13	1.14	49.87
J34F	3.44	2 928	0.03	0.47	0.50	14.44
J35A	8.47	31 018	0.28	1.20	1.48	17.51
J35B	8.12	5 831	0.05	1.24	1.29	15.93
J35C	1.98	3 633	0.03	0.88	0.91	46.12
J35D	9.82	5 284	0.05	3.65	3.70	37.66
J35E	1.33	1 224	0.01	0.21	0.22	16.63
J35F	6.67	2 796	0.03	2.02	2.05	30.67
J40A	9.73	970	0.01	5.03	5.04	51.79
J40B	5.45	0	0.00	2.71	2.71	49.72
J40C	15.81	1 076	0.01	6.58	6.59	41.68
J40D	10.21	8 056	0.07	4.20	4.27	41.86
J40E	7.48	1 908	0.02	3.45	3.47	46.36
K10A	2.34	53 970	0.49	1.16	1.65	70.62
K10B	1.96	4 727	0.04	1.20	1.24	63.43
K10C	4.43	0	0.00	2.33	2.33	52.60
K10D	2.53	1 579	0.01	1.10	1.11	44.05
K10E	13.70	4 122	0.04	4.30	4.34	31.66
K10F	2.82	7 002	0.06	0.99	1.05	37.37
K20A	19.85	9 650	0.09	6.15	6.24	31.43
K30A	28.06	6 994	0.06	7.15	7.21	25.71
K30B	21.52	6 334	0.06	5.03	5.09	23.64
K30C	27.80	146 970	1.34	7.83	9.17	32.99
K30D	18.44	9 839	0.09	7.43	7.52	40.78
K40A	8.99	512	0.00	3.79	3.79	42.21
K40B	13.52	0	0.00	4.85	4.85	35.87
K40C	17.00	6 256	0.06	4.32	4.38	25.75
K40D	17.74	20 130	0.18	3.71	3.89	21.95
K40E	26.56	13 515	0.12	10.61	10.73	40.41
K50A	27.43	49	0.00	10.09	10.09	36.79
K50B	24.71	14 745	0.13	8.58	8.71	35.27
K60A	6.43	154	0.00	4.20	4.20	65.34
K60B	8.43	43	0.00	5.70	5.70	67.62
K60C	10.95	1 668	0.02	6.60	6.62	60.41
K60D	23.54	681	0.01	12.43	12.44	52.83

Quaternary Catchment	Recharge (Mm ³ /a)	Population	Basic Human Needs (Mm ³ /a)	Groundwater Baseflow (Mm ³ /a)	Reserve (Mm ³ /a)	Reserve as % of Recharge
K60E	6.39	2 392	0.02	3.95	3.97	62.16
K60F	14.35	10 113	0.09	9.35	9.44	65.80
K60G	11.31	23 053	0.21	5.02	5.23	46.25
K70A	14.30	5 364	0.05	6.84	6.89	48.17
K70B	20.46	852	0.01	4.46	4.47	21.84

PROPOSED GROUNDWATER RESERVE – WATER QUALITY COMPONENT

The groundwater quality of quaternary catchments with available hydrochemistry data was assessed against the domestic water target water quality ranges as shown in Table 6.2. A summary of the results for the groundwater quality classification at quaternary level in the terms of basic human needs requirements is included in the tables that follow.

Table 6.2: Physical and chemical water quality

Parameter	Target Water Quality Ranges ¹⁾				
	Units	Class 0	Class I	Class II	Class III
pH	pH units	6 – 9	5 – 6 & 9 – 9.5	4 – 5 & > 9.5 – 10	<4 & > 10
Electrical Conductivity	mS/m	< 70	70 - 150	150 – 370	> 370
Calcium as Ca	mg/l	< 80	80 - 150	150 – 300	> 300
Magnesium as Mg	mg/l	< 70	70 - 100	100 – 200	> 200
Sodium as Na	mg/l	< 100	100 - 200	200 – 400	> 400
Chloride as Cl	mg/l	< 100	100 - 200	200 – 600	> 600
Sulphate as SO ₄	mg/l	< 200	200 - 400	400 – 600	> 600
Nitrate as NO _x -N	mg/l	< 6	6 - 10	10 – 20	> 20
Fluoride as F	mg/l	<0.7	0.7 – 1.0	1.0 – 1.5	> 1.5

¹⁾ Reference: Classification Systems in terms of – Water Research Commission: Quality of Domestic Water Supplies – Volume 1. Report No. TT 101/98, Second Edition, 1998.

The water quality for the following quaternary catchments were not assessed due to insufficient information (lack of representative groundwater quality data):

- G40B; G40D; G40E; G40G; G40K
- G50A; G50B; G50C; G50D; G50G; G50J
- H10D; H10E; H10J; H10K
- H20C; H20G; H20H
- H30B; H30E
- H40C; H40D; H40E; H40G; H40J; H40K; H40L
- H50A; H50B
- H60A; H60B; H60C; H60D; H60E; H60F; H60G; H60H; H60J; H60K; H60L
- H70A; H70C; H70D; H70E; H70F; H70G; H70H; H70J
- J11A; J11C; J11D; J11F; J11H; J11J; J11K
- J12A; J12C; J12E; J12J; J12M
- J13A; J13C; J22A; J22G; J22K; J23B; J23E; J23J, J24, J25B; J25D; J25E
- J31A; J31B; J31C; J32D; J33A; J33B; J33C; J34C; J34E; J35C; J35E; J35F
- J40A; J40B; J40C
- K10C; K10E; K10F
- K20A
- K30A; K30C; K30D
- K40A; K40B; K40C; K40D; K40E
- K50A
- K60A; K60B; K60C; K60D; K60E; K60F; K60G
- K70A; K70B

Table 6.3. Groundwater Quality Reserve: Breede Gouritz Water Management Area

Chemical Parameter	Unit	Quaternary Catchments G40C, G40F, G40J, G40L													
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾			
		G40C	G40F	G40J	G40L	G40C	G40F	G40J	G40L			G40C	G40F	G40J	G40L
pH		39	61	45	43	7.09	7.26	7.01	8.16	5.0 – 9.5		6.38–7.80	6.53–7.99	6.31–7.71	7.34–8.97
Electrical Conductivity	mS/m	39	61	45	43	20.70	20.90	21.20	208.00	<150		22.77	22.99	23.32	228.80
Calcium as Ca	mg/l	36	57	42	40	7.94	7.67	7.68	105.32	<150		8.73	8.43	8.45	115.85
Magnesium as Mg	mg/l	36	57	42	40	3.30	3.30	3.40	33.40	<100		3.63	3.63	3.74	36.74
Sodium as Na	mg/l	36	55	42	38	18.08	18.40	18.32	253.05	<200		19.89	20.24	20.16	253.05
Chloride as Cl	mg/l	36	58	42	40	31.94	32.69	32.57	391.39	<200		35.13	35.96	35.83	391.39
Sulphate as SO ₄	mg/l	36	58	42	40	5.71	5.35	5.81	78.14	<400		6.28	5.89	6.39	85.95
Nitrate as NO ₃ -N	mg/l	36	57	42	38	0.04	0.05	0.06	5.17	<10		0.04	0.06	0.07	5.68
Fluoride as F	mg/l	36	54	42	35	0.30	0.31	0.29	0.28	<1.0		0.33	0.34	0.32	0.31
Chemical Parameter	Unit	Quaternary Catchments G40M, G50E, G50F, G50H													
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾			
		G40M	G50E	G50F	G50H	G40M	G50E	G50F	G50H			G40M	G50E	G50F	G50H
pH		37	49	45	23	8.17	6.21	8.17	8.02	5.0 – 9.5		7.35–8.99	5.59–6.83	7.35–8.99	7.22–8.82
Electrical Conductivity	mS/m	37	49	45	23	109.10	54.90	91.90	1830.00	<150		120.01	60.39	101.09	1630.00
Calcium as Ca	mg/l	37	44	42	23	88.34	4.37	84.20	182.00	<150		97.18	4.81	92.62	182.00
Magnesium as Mg	mg/l	37	44	42	23	15.03	8.91	12.37	358.00	<100		16.54	9.81	13.61	358.00
Sodium as Na	mg/l	37	42	40	23	113.15	80.39	85.23	3107.18	<200		124.47	88.43	93.76	3107.18
Chloride as Cl	mg/l	37	44	42	23	203.98	143.85	153.44	5302.10	<200		203.98	158.24	168.79	5302.10
Sulphate as SO ₄	mg/l	37	44	41	23	31.52	20.58	25.59	742.40	<400		34.67	22.64	28.15	742.40
Nitrate as NO ₃ -N	mg/l	37	43	40	23	1.70	0.06	0.26	0.13	<10		1.87	0.07	0.29	0.14
Fluoride as F	mg/l	37	40	39	23	0.18	0.11	0.22	1.35	<1.0		0.20	0.12	0.25	1.49
Chemical Parameter	Unit	Quaternary Catchments G50J, H10A, H10B, H10C													
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾			
		G50J	H10A	H10B	H10C	G50J	H10A	H10B	H10C			G50J	H10A	H10B	H10C
pH		9	34	73	60	7.58	7.69	7.88	7.33	5.0 – 9.5		6.82–8.34	6.92–8.46	7.09–8.67	6.60–8.06
Electrical Conductivity	mS/m	9	34	73	60	142.90	180.00	49.00	45.60	<150		157.19	180.00	53.90	50.16
Calcium as Ca	mg/l	9	34	70	60	92.00	65.00	46.98	20.35	<150		101.20	71.50	51.67	22.39
Magnesium as Mg	mg/l	9	34	70	60	23.40	54.25	6.72	12.85	<100		25.74	59.68	7.39	14.14
Sodium as Na	mg/l	9	34	68	60	194.60	219.35	36.21	49.80	<200		200.00	241.29	39.83	54.78
Chloride as Cl	mg/l	9	34	70	60	338.90	380.55	80.09	106.25	<200		338.90	418.61	88.10	116.88
Sulphate as SO ₄	mg/l	9	34	70	60	38.70	153.60	21.75	22.30	<400		42.57	168.96	23.92	24.53
Nitrate as NO ₃ -N	mg/l	9	34	70	60	1.04	0.42	0.17	0.44	<10		1.14	0.46	0.19	0.48
Fluoride as F	mg/l	9	34	68	60	0.16	0.34	0.18	0.17	<1.0		0.18	0.37	0.20	0.19

Chemical Parameter	Unit	Quaternary Catchments H10F, H10G, H10H, H10I, H10L																	
		No. of Samples					Ambient GW quality or median ¹⁾					BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		H10F	H10G	H10H	H10I	H10L	H10F	H10G	H10H	H10I	H10L	H10F	H10G	H10H	H10I	H10L			
pH		15	117	54	82	7.29	7.05	7.59	6.98	6.98	6.56-8.02	6.35-7.76	6.83-8.35	6.28-7.68					
Electrical Conductivity	mS/m	15	117	54	82	24.70	13.40	117.70	9.87	9.87	27.17	14.74	129.47	10.85					
Calcium as Ca	mg/l	15	111	53	78	6.10	7.49	28.50	3.70	3.70	6.71	8.24	31.35	4.07					
Magnesium as Mg	mg/l	15	112	53	78	12.00	2.80	28.30	2.12	2.12	20.74	3.08	31.13	2.34					
Sodium as Na	mg/l	12	107	52	76	18.85	12.10	176.10	8.60	8.60	20.20	13.31	193.71	9.46					
Chloride as Cl	mg/l	15	114	54	79	30.50	18.45	308.75	14.80	14.80	33.55	20.30	308.75	16.28					
Sulphate as SO ₄	mg/l	12	109	52	79	4.15	5.50	53.65	3.00	3.00	4.57	6.05	59.02	3.30					
Nitrate as NO ₃ -N	mg/l	15	112	54	78	0.06	0.25	0.03	0.18	0.18	0.07	0.28	0.03	0.20					
Fluoride as F	mg/l	15	110	54	75	0.31	0.16	0.57	0.12	0.12	0.34	0.17	0.62	0.13					
Chemical Parameter	Unit	Quaternary Catchments H20A, H20B, H20D, H20E																	
		No. of Samples					Ambient GW quality or median ¹⁾					BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		H20A	H20B	H20D	H20E	H20A	H20B	H20D	H20E	H20A	H20B	H20D	H20E	H20A	H20B	H20D	H20E		
pH		85	344	12	15	7.20	7.09	6.77	6.46	6.46	6.48-7.92	6.38-7.79	6.09-7.44	5.81-7.11					
Electrical Conductivity	mS/m	85	344	12	15	50.60	40.75	12.75	2.30	2.30	55.66	44.83	14.03	2.53					
Calcium as Ca	mg/l	85	344	12	15	34.80	23.45	9.70	0.50	0.50	38.28	25.80	10.67	0.55					
Magnesium as Mg	mg/l	85	344	12	15	5.60	9.15	2.15	0.50	0.50	6.16	10.07	2.37	0.55					
Sodium as Na	mg/l	85	344	11	15	25.70	31.80	7.70	2.60	2.60	28.27	34.98	8.47	2.86					
Chloride as Cl	mg/l	85	344	12	15	55.00	50.55	13.90	1.50	1.50	60.50	55.61	15.29	1.65					
Sulphate as SO ₄	mg/l	85	344	11	15	81.20	39.65	4.00	2.00	2.00	89.32	43.62	4.40	2.20					
Nitrate as NO ₃ -N	mg/l	85	344	12	15	0.05	0.90	0.33	0.05	0.05	0.06	0.99	0.36	0.06					
Fluoride as F	mg/l	85	344	12	15	0.11	0.13	0.05	0.05	0.05	0.12	0.14	0.06	0.06					
Chemical Parameter	Unit	Quaternary Catchments H30A, H30C, H30D																	
		No. of Samples					Ambient GW quality or median ¹⁾					BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		H30A	H30C	H30D	H30A	H30C	H30D	H30A	H30C	H30D	H30A	H30C	H30D	H30A	H30C	H30D			
pH		302	9	21	45	7.10	7.95	7.70	7.81	7.81	6.39-7.80	7.16-8.75	6.93-8.47	7.03-8.59					
Electrical Conductivity	mS/m	302	9	21	45	29.00	142.00	41.70	19.50	19.50	31.90	156.20	45.87	21.45					
Calcium as Ca	mg/l	302	9	21	42	22.80	63.80	36.40	17.10	17.10	25.08	70.18	40.04	18.81					
Magnesium as Mg	mg/l	302	9	21	42	4.70	32.50	6.00	3.31	3.31	5.17	35.75	6.60	3.64					
Sodium as Na	mg/l	302	9	21	40	24.90	191.90	37.00	10.94	10.94	27.39	200.00	40.70	12.04					
Chloride as Cl	mg/l	302	9	21	43	28.30	273.50	82.00	18.08	18.08	31.13	300.85	90.20	19.89					
Sulphate as SO ₄	mg/l	302	9	21	43	28.00	121.90	48.00	4.30	4.30	30.80	134.09	52.80	4.73					
Nitrate as NO ₃ -N	mg/l	302	9	21	42	0.11	0.02	0.02	0.13	0.13	0.12	0.02	0.02	0.14					
Fluoride as F	mg/l	302	9	16	40	0.10	0.55	0.23	0.16	0.16	0.11	0.61	0.25	0.18					

Chemical Parameter	Unit	Quaternary Catchments H40A, H40B, H40F, H70B															
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾					
		H40A	H40B	H40F	H70B	H40A	H40B	H40F	H70B	H40A	H40B	H40F	H70B	H40A	H40B	H40F	H70B
pH		54	14	17	9	7.86	7.35	6.54	8.21	5.0 - 9.5	7.07-8.65	6.62-8.09	5.89-7.19	7.39-9.03			
Electrical Conductivity	mS/m	55	14	17	9	112.50	10.88	28.90	925.00	<150	123.75	11.97	31.79	925.00			
Calcium as Ca	mg/l	48	14	17	9	92.38	2.24	14.80	98.9	<150	101.62	2.47	16.28	108.79			
Magnesium as Mg	mg/l	49	14	17	9	44.90	0.93	7.00	259.10	<100	49.39	1.02	7.70	259.10			
Sodium as Na	mg/l	47	12	17	9	88.94	21.43	26.30	1722.90	<200	97.83	23.57	28.93	1722.90			
Chloride as Cl	mg/l	50	14	17	9	127.45	21.15	35.90	2741.20	<200	140.20	23.27	39.49	2741.20			
Sulphate as SO ₄	mg/l	50	14	17	9	229.91	3.50	20.70	338.60	<400	252.90	3.85	22.77	372.46			
Nitrate as NO ₃ -N	mg/l	50	14	17	9	0.06	0.18	4.70	0.02	<10	0.06	0.20	5.27	0.02			
Fluoride as F	mg/l	46	13	17	9	0.43	0.36	0.05	1.75	<1.0	0.47	0.39	0.06	1.75			
Chemical Parameter	Unit	Quaternary Catchments H70K, H90D, H90E, J11B															
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾					
		H70K	H90D	H90E	J11B	H70K	H90D	H90E	J11B	H70K	H90D	H90E	J11B	H70K	H90D	H90E	J11B
pH		29	21	85	37	7.50	8.12	8.09	7.80	5.0 - 9.5	6.75-8.25	7.31-8.94	7.28-8.90	7.02-8.58			
Electrical Conductivity	mS/m	29	21	85	37	121.00	102.70	179.00	81.30	<150	133.10	112.97	179.00	89.43			
Calcium as Ca	mg/l	26	21	81	37	46.20	53.66	87.18	60.50	<150	50.82	59.02	95.89	66.55			
Magnesium as Mg	mg/l	26	21	82	37	19.20	17.40	28.50	18.40	<100	21.12	19.14	31.35	20.24			
Sodium as Na	mg/l	26	21	80	37	166.95	144.96	208.48	79.20	<200	183.65	159.45	208.48	87.12			
Chloride as Cl	mg/l	26	20	81	37	273.10	247.79	358.72	94.70	<200	273.10	247.79	358.72	104.17			
Sulphate as SO ₄	mg/l	26	21	82	37	29.88	33.16	36.65	68.60	<400	32.87	36.47	40.31	75.46			
Nitrate as NO ₃ -N	mg/l	26	20	79	37	0.02	0.83	3.40	1.41	<10	0.02	0.91	3.74	1.55			
Fluoride as F	mg/l	26	20	77	37	0.52	0.14	0.18	0.82	<1.0	0.57	0.15	0.19	0.90			
Chemical Parameter	Unit	Quaternary Catchments J11E, J11G, J12B, J12D															
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾					
		J11E	J11G	J12B	J12D	J11E	J11G	J12B	J12D	J11E	J11G	J12B	J12D	J11E	J11G	J12B	J12D
pH		81	20	17	11	8.19	8.18	7.73	8.23	5.0 - 9.5	7.37-9.01	7.36-9.00	6.96-8.50	7.41-9.05			
Electrical Conductivity	mS/m	81	20	17	11	174.30	164.50	368.00	195.00	<150	174.30	164.50	368.00	195.00			
Calcium as Ca	mg/l	77	20	17	11	98.70	115.05	117.10	68.30	<150	108.57	126.56	128.81	75.13			
Magnesium as Mg	mg/l	77	20	17	11	45.00	44.65	129.30	50.60	<100	49.50	49.12	129.30	55.66			
Sodium as Na	mg/l	73	20	17	11	197.89	190.65	335.70	252.80	<200	217.68	200.00	335.70	252.80			
Chloride as Cl	mg/l	75	20	17	11	250.10	294.70	726.10	415.00	<200	250.10	294.70	726.10	415.00			
Sulphate as SO ₄	mg/l	77	20	17	11	179.50	120.45	144.20	104.80	<400	197.45	132.50	158.62	115.28			
Nitrate as NO ₃ -N	mg/l	71	20	17	11	0.30	1.18	0.06	0.02	<10	0.33	1.29	0.07	0.02			
Fluoride as F	mg/l	67	20	17	11	0.88	0.92	0.80	0.53	<1.0	0.96	0.92	0.88	0.58			

Chemical Parameter	Unit	Quaternary Catchments J12F, J12G, J12H, J12K																			
		No. of Samples						Ambient GW quality or median ¹⁾						BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		J12G		J12H		J12K		J12F	J12G	J12H	J12K					J12F	J12G	J12H	J12K		
pH		10	11	13	10	8.23	8.20	7.75	8.13	5.0 – 9.5	7.41–9.05	7.38–9.02	6.98–8.53	7.32–8.94							
Electrical Conductivity	mS/m	10	11	13	10	306.50	323.00	157.00	462.00	<150	306.50	323.00	157.00	462.00							
Calcium as Ca	mg/l	10	11	13	10	79.65	103.10	121.90	212.95	<150	87.62	113.41	134.09	212.95							
Magnesium as Mg	mg/l	10	11	13	10	81.35	91.80	26.00	112.55	<100	89.49	100.98	28.60	123.81							
Sodium as Na	mg/l	10	11	13	10	406.70	262.30	151.50	703.25	<200	406.70	262.30	166.65	703.25							
Chloride as Cl	mg/l	10	11	13	10	691.30	795.60	258.70	1168.65	<200	691.30	795.60	258.70	1168.65							
Sulphate as SO ₄	mg/l	10	11	13	10	241.45	107.20	156.00	459.80	<400	265.60	117.92	171.60	459.80							
Nitrate as NO ₃ -N	mg/l	10	11	13	10	0.02	0.02	0.02	0.02	<10	0.02	0.02	0.02	0.02							
Fluoride as F	mg/l	10	11	13	10	0.56	0.50	0.39	0.55	<1.0	0.61	0.55	0.43	0.61							
Chemical Parameter	Unit	Quaternary Catchments J12L, J13B, J21A, J21B																			
		No. of Samples						Ambient GW quality or median ¹⁾						BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		J13B		J21A		J21B		J12L	J13B	J21A	J21B					J12L	J13B	J21A	J21B		
pH		57	15	64	56	7.54	7.98	7.97	7.60	5.0 – 9.5	6.79–8.29	7.18–8.78	7.17–8.76	6.84–8.35							
Electrical Conductivity	mS/m	57	15	64	56	25.50	418.00	61.40	100.50	<150	28.05	418.00	67.54	110.55							
Calcium as Ca	mg/l	54	14	60	56	16.11	94.90	58.18	60.10	<150	17.72	104.39	64.00	66.11							
Magnesium as Mg	mg/l	54	15	59	56	2.89	93.80	18.81	19.45	<100	3.18	103.18	20.69	21.40							
Sodium as Na	mg/l	52	15	60	54	21.85	623.60	43.23	116.95	<200	24.03	623.60	47.55	128.65							
Chloride as Cl	mg/l	53	15	58	56	30.71	906.39	18.77	104.70	<200	33.78	906.39	20.64	115.17							
Sulphate as SO ₄	mg/l	54	15	57	54	7.55	253.60	75.14	113.80	<400	8.31	278.96	82.65	125.18							
Nitrate as NO ₃ -N	mg/l	52	15	54	56	0.03	0.35	0.09	2.56	<10	0.03	0.39	0.09	2.81							
Fluoride as F	mg/l	51	15	54	56	0.21	0.72	0.46	0.87	<1.0	0.23	0.79	0.51	0.96							
Chemical Parameter	Unit	Quaternary Catchments J21C, J21D, J21E, J22B																			
		No. of Samples						Ambient GW quality or median ¹⁾						BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		J21C		J21D		J22B		J21C	J21D	J21E	J22B					J21C	J21D	J21E	J22B		
pH		107	40	26	21	7.75	7.66	7.96	8.00	5.0 – 9.5	6.98–8.53	6.89–8.42	7.16–8.76	7.20–8.80							
Electrical Conductivity	mS/m	107	40	26	21	76.40	85.20	78.80	74.20	<150	84.04	93.72	86.68	81.62							
Calcium as Ca	mg/l	107	40	26	21	56.90	43.30	61.00	60.30	<150	62.59	47.63	67.10	66.33							
Magnesium as Mg	mg/l	107	40	26	21	16.40	17.70	16.00	24.80	<100	18.04	19.47	17.60	27.28							
Sodium as Na	mg/l	107	38	26	21	83.80	107.90	83.40	67.90	<200	92.18	118.69	91.74	74.69							
Chloride as Cl	mg/l	107	40	26	21	76.90	102.60	93.00	57.20	<200	84.59	112.86	102.30	62.92							
Sulphate as SO ₄	mg/l	107	38	26	21	83.50	95.70	63.05	63.40	<400	91.85	105.27	58.36	69.74							
Nitrate as NO ₃ -N	mg/l	107	39	26	21	3.13	3.55	2.23	1.95	<10	3.44	3.91	2.45	2.15							
Fluoride as F	mg/l	107	40	26	21	0.68	1.00	0.95	0.91	<1.0	0.75	1.00	0.95	1.00							

Chemical Parameter	Unit	Quaternary Catchments J22C, J22D, J22E, J22F															
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾					
		J22C	J22D	J22E	J22F	J22C	J22D	J22E	J22F	J22C	J22D	J22E	J22F	J22C	J22D	J22E	J22F
pH		26	46	80	24	8.00	7.94	7.79	7.80	5.0 – 9.5	7.20–8.80	7.15–8.73	7.01–8.57	7.02–8.58			
Electrical Conductivity	mS/m	26	46	80	24	97.10	89.85	88.70	103.45	<150	106.81	98.84	97.57	113.80			
Calcium as Ca	mg/l	26	46	80	24	50.15	44.95	61.80	82.10	<150	55.17	49.45	67.98	90.31			
Magnesium as Mg	mg/l	26	46	80	24	27.00	24.80	21.00	18.95	<100	29.70	27.28	23.10	20.85			
Sodium as Na	mg/l	26	46	80	24	102.75	118.25	95.95	86.35	<200	113.03	130.08	105.55	94.99			
Chloride as Cl	mg/l	26	46	80	24	94.00	102.75	92.40	109.40	<200	103.40	113.03	101.64	120.34			
Sulphate as SO ₄	mg/l	26	46	80	24	123.80	121.30	90.80	110.70	<400	136.18	133.43	99.88	121.77			
Nitrate as NO ₃ -N	mg/l	26	46	80	24	1.78	1.47	4.12	3.82	<10	1.95	1.61	4.53	4.20			
Fluoride as F	mg/l	26	46	80	24	1.03	1.10	1.05	0.89	<1.0	1.03	1.10	1.05	0.97			
Chemical Parameter	Unit	Quaternary Catchments J22H, J22J, J23A, J23C															
Chemical Parameter	Unit	No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾					
		J22H	J22J	J23A	J23C	J22H	J22J	J23A	J23C	J22H	J22J	J23A	J23C	J22H	J22J	J23A	J23C
		32	26	39	10	7.79	7.83	7.93	7.78	5.0 – 9.5	7.01–8.56	7.14–8.72	7.00–8.56	7.01–8.56	7.05–8.61	7.14–8.72	7.00–8.56
Electrical Conductivity	mS/m	32	26	39	10	145.05	83.20	238.00	136.00	<150	145.05	91.52	238.00	149.60			
Calcium as Ca	mg/l	32	26	36	10	105.60	50.75	125.25	126.75	<150	116.16	55.83	137.78	139.43			
Magnesium as Mg	mg/l	32	26	36	10	38.25	20.45	16.26	32.70	<100	42.08	22.50	17.89	35.97			
Sodium as Na	mg/l	32	26	35	10	158.05	86.90	342.80	143.70	<200	173.86	95.59	342.80	158.07			
Chloride as Cl	mg/l	32	26	35	10	208.55	82.15	265.46	159.05	<200	229.41	90.37	265.46	174.96			
Sulphate as SO ₄	mg/l	32	26	36	10	225.10	117.50	497.01	112.50	<400	247.61	129.25	497.01	123.75			
Nitrate as NO ₃ -N	mg/l	32	26	35	10	3.31	3.02	0.04	1.10	<10	3.64	3.32	0.04	1.21			
Fluoride as F	mg/l	32	26	33	10	1.09	0.90	2.14	0.69	<1.0	1.19	0.99	2.14	0.76			
Chemical Parameter	Unit	Quaternary Catchments J23D, J23F, J24A, J24B															
Chemical Parameter	Unit	No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾					
		J23D	J23F	J24A	J24B	J23D	J23F	J24A	J24B	J23D	J23F	J24A	J24B	J23D	J23F	J24A	J24B
		26	85	168	220	8.10	7.97	7.77	7.79	5.0 – 9.5	7.29–8.91	7.17–8.76	6.99–8.54	7.01–8.56	7.29–8.91	7.17–8.76	6.99–8.54
Electrical Conductivity	mS/m	26	85	168	220	155.00	126.70	86.30	86.54	<150	155.00	139.37	94.93	95.19			
Calcium as Ca	mg/l	26	83	168	213	98.95	51.07	70.60	76.59	<150	108.85	56.17	77.66	84.24			
Magnesium as Mg	mg/l	26	83	168	214	40.65	28.60	18.55	17.15	<100	44.72	31.46	20.41	18.87			
Sodium as Na	mg/l	26	80	168	210	166.55	120.85	91.25	83.50	<200	183.21	132.94	100.38	91.85			
Chloride as Cl	mg/l	26	83	168	214	208.95	140.10	78.80	78.20	<200	208.95	154.11	86.68	86.02			
Sulphate as SO ₄	mg/l	26	83	168	214	151.40	157.15	103.40	69.00	<400	166.54	172.86	113.74	75.90			
Nitrate as NO ₃ -N	mg/l	26	83	168	212	1.89	0.12	1.85	3.45	<10	2.08	0.13	2.04	3.80			
Fluoride as F	mg/l	26	81	168	209	0.79	0.48	0.92	0.81	<1.0	0.86	0.52	1.01	0.89			

Chemical Parameter	Unit	Quaternary Catchments J24C, J24D, J24E, J25A													
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾			
		J24C	J24D	J24E	J25A	J24C	J24D	J24E	J25A			J24C	J24D	J24E	J25A
pH		112	97	18	33	7.70	7.90	7.94	7.11	5.0 - 9.5	6.93-8.47	7.11-8.69	7.15-8.73	6.40-7.83	
Electrical Conductivity	mS/m	112	97	18	33	96.70	146.70	161.50	50.90	<150	106.37	146.70	161.50	55.99	
Calcium as Ca	mg/l	112	93	18	33	68.45	94.20	106.00	14.80	<150	75.30	103.62	116.60	16.28	
Magnesium as Mg	mg/l	112	93	18	33	18.00	28.80	30.95	10.90	<100	19.80	31.68	34.05	11.99	
Sodium as Na	mg/l	112	86	18	33	103.40	162.40	198.65	50.40	<200	113.74	178.64	198.65	55.44	
Chloride as Cl	mg/l	112	95	18	33	120.20	194.50	213.80	96.50	<200	132.22	194.50	213.80	106.15	
Sulphate as SO ₄	mg/l	112	89	18	33	78.40	114.00	105.70	26.40	<400	86.24	125.40	116.27	29.04	
Nitrate as NO ₃ -N	mg/l	112	95	18	32	3.90	1.76	0.82	0.02	<10	4.28	1.94	0.90	0.02	
Fluoride as F	mg/l	112	93	18	32	0.79	0.81	1.05	0.31	<1.0	0.87	0.89	1.05	0.34	

Chemical Parameter	Unit	Quaternary Catchments J25C, J31D, J32A, J32B													
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾			
		J25C	J31D	J32A	J32B	J25C	J31D	J32A	J32B			J25C	J31D	J32A	J32B
pH		31	48	19	15	7.11	6.54	7.68	7.86	5.0 - 9.5	6.40-7.83	5.89-7.19	6.91-8.45	7.07-8.65	
Electrical Conductivity	mS/m	31	48	19	15	50.90	14.06	165.80	187.00	<150	55.99	15.46	165.80	187.00	
Calcium as Ca	mg/l	31	43	19	15	14.80	4.92	85.60	99.50	<150	16.28	5.41	94.16	109.45	
Magnesium as Mg	mg/l	31	43	19	15	10.90	2.86	42.50	44.50	<100	11.99	3.14	46.75	48.95	
Sodium as Na	mg/l	31	41	19	15	50.40	9.30	172.20	175.10	<200	55.44	10.23	189.42	192.61	
Chloride as Cl	mg/l	31	43	19	15	96.50	23.80	203.20	242.60	<200	106.15	26.18	203.20	242.60	
Sulphate as SO ₄	mg/l	31	43	19	15	26.40	9.90	180.30	170.80	<400	29.04	10.89	198.33	187.88	
Nitrate as NO ₃ -N	mg/l	31	41	19	15	0.02	0.03	3.80	0.88	<10	0.02	0.03	4.18	0.96	
Fluoride as F	mg/l	31	39	19	15	0.31	0.29	1.06	0.99	<1.0	0.34	0.32	1.06	0.99	

Chemical Parameter	Unit	Quaternary Catchments J32C, J32E, J33D, J33E													
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾		Groundwater Quality Reserve ³⁾			
		J32C	J32E	J33D	J33E	J32C	J32E	J33D	J33E			J32C	J32E	J33D	J33E
pH		29	41	40	149	7.76	7.55	7.53	6.64	5.0 - 9.5	6.98-8.54	6.80-8.30	6.78-8.29	6.98-7.30	
Electrical Conductivity	mS/m	29	41	40	149	127.40	19.80	19.80	12.10	<150	140.14	21.78	21.78	13.31	
Calcium as Ca	mg/l	29	40	39	148	80.80	11.07	11.00	2.76	<150	88.88	12.17	12.10	3.04	
Magnesium as Mg	mg/l	29	40	39	148	30.90	3.50	3.50	2.07	<100	33.99	3.85	3.85	2.27	
Sodium as Na	mg/l	29	39	38	146	143.90	12.40	12.39	13.40	<200	158.29	13.64	13.63	14.74	
Chloride as Cl	mg/l	29	40	38	147	170.70	23.87	23.64	22.99	<200	187.77	26.26	26.01	25.29	
Sulphate as SO ₄	mg/l	29	39	38	146	121.00	5.72	5.58	6.34	<400	133.10	6.29	6.14	6.97	
Nitrate as NO ₃ -N	mg/l	29	39	38	147	2.15	0.04	0.04	0.19	<10	2.37	0.04	0.04	0.20	
Fluoride as F	mg/l	29	37	36	146	0.90	0.27	0.27	0.11	<1.0	0.99	0.30	0.30	0.12	

Chemical Parameter	Unit	Quaternary Catchments J33F, J34A, J34B, J34D															
		No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		J33F	J34A	J34B	J34D	J33F	J34A	J34B	J34D	J33F	J34A	J34B	J34D	J33F	J34A	J34B	J34D
pH		70	36	11	22	6.82	6.33	6.77	6.56	5.0 – 9.5	6.14-7.50	5.70-6.96	6.09-7.45	5.90-7.22			
Electrical Conductivity	mS/m	70	36	11	22	44.60	19.55	114.50	12.54	<150	49.06	21.51	125.95	13.79			
Calcium as Ca	mg/l	70	36	11	22	11.95	3.26	46.30	2.39	<150	13.15	3.68	50.93	2.63			
Magnesium as Mg	mg/l	70	36	11	22	7.05	3.44	21.80	1.73	<100	7.76	3.78	23.98	1.91			
Sodium as Na	mg/l	70	36	11	22	34.25	25.95	136.70	15.40	<200	37.68	28.55	150.37	16.94			
Chloride as Cl	mg/l	70	36	11	22	61.74	40.85	252.50	27.36	<200	67.92	44.94	252.50	30.10			
Sulphate as SO ₄	mg/l	70	36	11	22	27.50	5.22	42.20	2.50	<400	30.25	5.74	46.42	2.75			
Nitrate as NO ₃ -N	mg/l	68	36	11	22	0.08	0.17	0.37	0.22	<10	0.09	0.19	0.41	0.24			
Fluoride as F	mg/l	69	36	11	22	0.22	0.11	0.29	0.08	<1.0	0.24	0.12	0.32	0.08			
Chemical Parameter	Unit	Quaternary Catchments J34F, J35A, J35B, J35D															
Chemical Parameter	Unit	No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		J34F	J35A	J35B	J35D	J34F	J35A	J35B	J35D	J34F	J35A	J35B	J35D	J34F	J35A	J35B	J35D
		pH		9	17	49	49	7.66	8.07	7.01	7.64	5.0 – 9.5	6.89-8.43	7.26-8.88	6.31-7.71	6.88-8.40	
Electrical Conductivity	mS/m	9	17	49	49	46.60	42.00	46.20	173.10	<150	51.26	46.20	50.82	173.10			
Calcium as Ca	mg/l	9	17	47	48	6.87	6.270	6.23	86.30	<150	7.55	68.97	6.85	94.93			
Magnesium as Mg	mg/l	9	17	47	48	7.70	6.30	8.70	35.65	<100	8.47	6.93	9.57	39.22			
Sodium as Na	mg/l	9	17	45	48	61.80	11.10	57.30	145.70	<200	67.98	12.21	63.03	160.27			
Chloride as Cl	mg/l	9	17	45	48	101.64	19.10	108.70	239.50	<200	111.81	21.01	119.57	239.50			
Sulphate as SO ₄	mg/l	9	17	47	48	27.90	11.50	4.85	95.95	<400	30.69	12.65	5.34	105.55			
Nitrate as NO ₃ -N	mg/l	9	17	45	48	0.05	0.16	4.51	0.64	<10	0.06	0.18	4.96	0.70			
Fluoride as F	mg/l	9	17	43	47	0.48	0.16	0.10	0.52	<1.0	0.53	0.18	0.11	0.57			
Chemical Parameter	Unit	Quaternary Catchments J40D, J40E, K10A, K10B															
Chemical Parameter	Unit	No. of Samples				Ambient GW quality or median ¹⁾				BHN Reserve ²⁾				Groundwater Quality Reserve ³⁾			
		J40D	J40E	K10A	K10B	J40D	J40E	K10A	K10B	J40D	J40E	K10A	K10B	J40D	J40E	K10A	K10B
		pH		48	63	20	14	7.30	7.52	7.88	7.51	5.0 – 9.5	6.57-8.03	6.77-8.28	7.09-8.67	6.76-8.26	
Electrical Conductivity	mS/m	48	63	20	14	83.15	219.00	276.00	214.50	<150	91.47	219.00	276.00	214.50			
Calcium as Ca	mg/l	48	63	20	14	12.41	59.40	52.50	32.87	<150	13.65	65.34	57.75	36.15			
Magnesium as Mg	mg/l	48	63	20	14	14.72	31.97	44.17	35.74	<100	16.19	35.17	48.58	39.32			
Sodium as Na	mg/l	45	63	20	14	131.41	288.45	399.60	288.65	<200	144.55	288.45	399.60	288.65			
Chloride as Cl	mg/l	48	63	20	13	213.26	492.38	698.45	545.73	<200	213.26	492.38	698.45	545.73			
Sulphate as SO ₄	mg/l	45	63	20	14	31.69	65.21	97.08	59.60	<400	34.86	71.73	106.79	66.56			
Nitrate as NO ₃ -N	mg/l	47	62	20	12	0.09	0.04	0.05	0.27	<10	0.10	0.04	0.06	0.29			
Fluoride as F	mg/l	48	63	20	13	0.14	0.22	0.22	0.25	<1.0	0.16	0.24	0.24	0.28			

Chemical Parameter	Unit	Quaternary Catchments K10D, K30B, K50B											
		No. of Samples			Ambient GW quality or median ¹⁾			BHN Reserve ²⁾	Groundwater Quality Reserve ³⁾				
		K10D	K30B	K50B	K10D	K30B	K50B		K10D	K30B	K50B		
pH		11	47	9	7.85	6.83	7.48	5.0 – 9.5	7.07–8.64	6.15–7.51	6.73–8.22		
Electrical Conductivity	mS/m	11	47	9	257.00	27.72	61.90	<150	257.00	30.49	68.09		
Calcium as Ca	mg/l	11	43	9	30.82	4.27	15.08	<150	33.90	4.70	16.68		
Magnesium as Mg	mg/l	11	43	9	28.10	7.49	9.80	<100	30.91	8.24	10.78		
Sodium as Na	mg/l	11	41	9	426.18	28.35	85.57	<200	426.18	31.19	94.13		
Chloride as Cl	mg/l	11	42	9	533.12	41.92	139.99	<200	533.12	46.11	153.98		
Sulphate as SO ₄	mg/l	11	44	9	66.60	7.95	17.54	<400	73.26	8.74	19.29		
Nitrate as NO ₃ -N	mg/l	10	43	9	0.09	7.99	0.48	<10	0.10	8.79	0.52		
Fluoride as F	mg/l	10	23	9	0.84	0.16	0.34	<1.0	0.93	1.49	0.18		

¹⁾ Based on long term groundwater quality datasets (DWS Water Management System). Minimum number of analyses used for the statistical evaluation is nine (9).
²⁾ Upper limit of Class 1 water quality [Drinking] (WRC et al. 2nd Edition, 1998, Volume 1: Assessment Guide); and
³⁾ Median value plus 10%. Where a difference in the water quality values for the ambient groundwater quality and basic human needs was found, the lesser or more protective value was selected for the groundwater quality Reserve. Where the ambient groundwater quality was selected as the groundwater quality Reserve, the value was scaled up by 10 per cent provided that the value does not exceed the BHN Reserve.

Table 6.4: Summary of the water quality class and parameters of concern

Quaternary catchment	Water quality class (WRC, 1998)	Water quality parameters of concern
G40C	0	None
G40F	0	None
G40J	0	None
G40L	II	Sodium, Chloride
G40M	II	Chloride
G50E	I	Chloride
G50F	I	Chloride
G50H	III	Electrical Conductivity, Magnesium, Sodium, Chloride, Sulphate
G50J	II	Chloride, Sodium, Electrical Conductivity
H10A	II	Sodium, Chloride, Electrical Conductivity
H10B	0	None
H10C	I	Chloride
H10F	0	None
H10G	0	None
H10H	II	Chloride
H10L	0	None
H20A	0	None
H20B	0	None
H20D	0	None
H20E	0	None
H20F	0	None
H30A	II	Chloride
H30C	0	None
H30D	0	None
H40A	I	Electrical Conductivity, Calcium, Chloride
H40B	0	None
H40F	0	None
H70B	III	Electrical Conductivity, Magnesium, Sodium, Chloride
H70K	II	Chloride
H90D	II	Chloride
H90E	II	Electrical Conductivity, Sodium, Chloride
J11B	I	Electrical Conductivity
J11E	II	Electrical Conductivity, Chloride
J11G	II	Electrical Conductivity, Chloride
J12B	III	Electrical Conductivity, Magnesium, Sodium, Chloride
J12D	II	Electrical Conductivity, Sodium, Chloride
J12F	III	Electrical Conductivity, Sodium, Chloride
J12G	III	Electrical Conductivity, Sodium, Chloride
J12H	II	Electrical Conductivity, Chloride
J12K	III	Electrical Conductivity, Calcium, Sodium, Chloride, Sulphate
J12L	0	None
J13B	III	Electrical Conductivity, Sodium, Chloride
J21A	0	None
J21B	I	Electrical Conductivity, Sodium, Chloride
J21C	I	Electrical Conductivity
J21D	I	Electrical Conductivity, Sodium, Chloride
J21E	I	Electrical Conductivity
J22B	I	Electrical Conductivity
J22C	I	Electrical Conductivity, Sodium
J22D	I	Electrical Conductivity, Sodium, Chloride
J22E	I	Electrical Conductivity
J22F	I	Electrical Conductivity, Calcium, Chloride
J22H	II	Electrical Conductivity, Calcium, Sodium, Chloride
J22J	I	Electrical Conductivity
J23A	II	Electrical Conductivity, Sodium, Chloride, Sulphate
J23C	I	Electrical Conductivity, Calcium, Sodium, Chloride
J23D	II	Electrical Conductivity, Chloride
J23F	I	Electrical Conductivity, Sodium, Chloride
J24A	I	Electrical Conductivity
J24B	I	Electrical Conductivity
J24C	I	Electrical Conductivity, Sodium, Chloride
J24D	II	Electrical Conductivity, Chloride

Quaternary catchment	Water quality class (WRC, 1998)	Water quality parameters of concern
J24E	II	Electrical Conductivity, Chloride
J25A	0	None
J25C	0	None
J31D	0	None
J32A	II	Electrical Conductivity, Chloride
J32B	II	Electrical Conductivity, Chloride
J32C	I	Electrical Conductivity, Calcium, Sodium, Chloride
J32E	0	None
J33D	0	None
J33E	0	None
J33F	0	None
J34A	0	None
J34B	II	Chloride
J34D	0	None
J34F	I	Chloride
J35A	0	None
J35B	I	Chloride
J35D	II	Electrical Conductivity, Chloride
J40D	II	Chloride, Sodium
J40E	II	Electrical Conductivity, Sodium, Chloride
K10A	III	Electrical Conductivity, Sodium, Chloride
K10B	II	Electrical Conductivity, Sodium, Chloride
K10D	III	Electrical Conductivity, Sodium, Chloride
K30B	0	None
K50B	I	Chloride

CONTINUES ON PAGE 258 OF BOOK 3

Printed by and obtainable from the Government Printer, Bosman Street, Private Bag X85, Pretoria, 0001
Contact Centre Tel: 012-748 6200. eMail: info.egazette@gpw.gov.za
Publications: Tel: (012) 748 6053, 748 6061, 748 6065



Government Gazette Staatskoerant

REPUBLIC OF SOUTH AFRICA
REPUBLIEK VAN SUID AFRIKA

Vol. 686

26

August
Augustus

2022

No. 46798

PART 3 OF 4

N.B. The Government Printing Works will not be held responsible for the quality of "Hard Copies" or "Electronic Files" submitted for publication purposes

ISSN 1682-5845



9 771682 584003



AIDS HELPLINE: 0800-0123-22 Prevention is the cure

7. ESTUARIES (WATER QUANTITY COMPONENT)

Table 7.1 Water Quantity

Quaternary Catchment	Estuary name	PES	REC	nMAR (MCM)	EWR nMAR) (MCM)
G40B	Rooiels	B	B	*9.44	n/a
G40D	Palmiet	C	B	*177.94	n/a
G40G	Bot/Kleinmond	C	B	*77.67	n/a
G40H	Onrus	E	D	*4.74	n/a
G40L	Klein	C	B	*51.21	n/a
G40M	Uilkraals	D	C	*6.82	n/a
G40F	Heuningnes	C	A	29.53	n/a
H70K	Breede	B	B	1785.00	954.00
H80E	Duiwenhoks	B	A	89.29	73.01
H90C	Goukou	C	B	115.95	91.73
J40B	Gouritz	C	B	623.52	377.23
K10A	Blinde	C	C	n/a	n/a
K10B	Hartenbos	D	C	n/a	n/a
K10F	Klein Brak	C	C	50.67	37.66
K20A	Groot Brak	D	C	36.79	11.11
K30A	Maalgate	B	C	41.51	24.41
K30B	Gwaing	B	C	35.07	21.7
K30C	Kaaimans	B	B	53.6	41.3

Quaternary Catchment	Estuary name	PES	REC	nMAR (MCM)	EWR nMAR (MCM)
K30D	Wilderness system: Touws	B	A	29.66	25.15
K40D	Swartvlei	B	B	83.4	56.6
K40E	Goukamma	B	A	57.5	48.8
K50B	Kynsna	B	B	83.2	63.4
K60 E and F	Keurbooms	A/B	A/B	232	214.10
K60G	Noetsie	B	B	4.8	n/a
K60G	Piesang	D	B/C	n/a	n/a
K70A	Groot (Wes)	B	A	n/a	n/a
K70A	Matjie	B	B	5.10	n/a
K70A	Sout	A	A	11.22	n/a
K70B	Bloukrans	A	A	n/a	n/a

*Present Day MAR

ESTUARY WATER QUALITY COMPONENT

Table 7.2: EcoSpecs and Thresholds of Potential Concern for the Blinde Estuary

Ecological component	EcoSpecs	Thresholds of Potential Concern
Hydrology	Maintain flow regime (small system needs most flows)	MAR does not vary by more than 10% from present Floods (indicated by 1:10 year event) do not reduce by more than 5% from present Base flows do not differ by more than 5% from present
Hydrodynamics	Maintain mouth state to create the require habitat for birds, fish, macrophytes, microalgae and water quality	Closed mouth state increase/decrease by 10% from present Presence of semi-closed mouth state with continuous outflow to sea. Average water depth <0.5 m (to be confirmed by monitoring) Rate of change in water level > 30% from present
Water quality	Salinity distribution not to cause exceedance of TPCs for fish, invertebrates, macrophytes and microalgae Turbidity and dissolved oxygen not to cause exceedance of TPCs for biota Dissolved inorganic (DIN)/dissolved inorganic phosphate (DIP) concentrations not to cause exceedance of TPCs for macrophytes and microalgae Toxic substances not to cause exceedance of TPCs for biota	Salinity > 20 (expected range 5-15) Dissolved oxygen (DO) < 5 mg/l in estuary Turbidity > 10 NTU in low flow Secchi depth: to bottom DIN > 100 µg/l (average) DIP > 20 µg/l (average) Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) Concentrations in sediment exceed target values as per Western Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Sediment dynamics	Flood regime to maintain the sediment distribution patterns and aquatic habitat (in stream physical habitat) so as not to exceed TPCs for biota Changes in sediment grain-size distribution patterns not to cause exceedance of TPCs in benthic invertebrates Change in average sediment composition and characteristics Change in average bathymetry	Average sediment composition (% fractions) along estuary change from baseline (to be measured) by 30% (per survey) Average depth along main channel change from 30% of baseline (to be determined) (system expected to experience significant fluctuation in bathymetry between flood and extended closed periods)
Microalgae	Maintain low/median phytoplankton/benthic microalgae biomass Prevent formation of phytoplankton blooms	Phytoplankton > 3.5 µg/l (median) Benthic microalgae > 23 mg/m ² (median) Phytoplankton > 20 µg/l and/or cell density > 10 000 cells/ml (once-off)
Macrophytes	Maintain distribution of macrophyte habitats Prevent the spread of reeds into open water Prevent an increase in nutrients and macro-algal blooms Prevent the spread of invasive trees (e.g. <i>Acacia</i> spp.) in the riparian zone	20% change in the macrophyte area. (Reeds currently cover 0.04 ha.) Reeds occupy > 0.5 ha Macro-algal blooms cover > 50% of the open water area Presence of invasive aquatic macrophytes e.g. <i>Azolla</i> , water hyacinth etc. Invasive trees cover > 50% of riparian zone
Invertebrates	Establish presence/absence of sand prawn <i>Callinectes kraussi</i> on sand banks in lower estuary Establish presence/absence of the copepod <i>Pseudodiaptomus hessei</i> or estuarine congeneric in the zooplankton of the estuary	If present populations deviate from average baselines (as determined in first three visits) by more 30%
Fish	Fish assemblage should comprise the five estuarine association categories in similar proportions (diversity and abundance) to that under the reference. Numerically, assemblage should comprise: Ia estuarine residents (50-80% of total abundance)	Ia estuarine residents < 50% Ib marine and estuarine breeders < 10% IIa obligate estuarine-dependent < 10% IIb estuarine associated species < 5% IIc marine opportunists < 20% III marine vagrants > 5% IV indigenous fish < 1%

Ecological component	EcoSpecs	Thresholds of Potential Concern
	<p>Ib marine and estuarine breeders (5-20%) IIa obligate estuarine-dependent (10-20%) IIb estuarine associated species (5-15%), IIc marine opportunists (20-80%) III marine vagrants (not more than 5%) IV indigenous fish (1-5%) V catadromous species (1-5%)</p> <p>Category Ia species should contain viable populations of at least two species (e.g. <i>G.aestuaria</i>, & <i>Hyporamphus capensis</i>).</p> <p>Category IIa obligate dependents should be well represented by at least two large exploited species (i.e. <i>L. lithognathus</i>, <i>Lichia amia</i>).</p> <p>REI (River Estuary Interface) species dominated by both <i>Myxus capensis</i> and <i>G. aestuaria</i>.</p>	V catadromous species <1%
Birds	Maintain population of original groups of birds present on the estuary	Number of birds in any group, other than species that are increasing regionally such as Egyptian geese, drops below the baseline median (determined by past data and/or initial surveys) number of species and/or birds counted for three consecutive summer or winter counts

Table 7.3: EcoSpecs and Thresholds of Potential Concern for the Hartenbos Estuary

Ecological component	EcoSpecs	Thresholds of Potential Concern
Hydrology	Maintain at least present day base flows (to be confirmed)	MAR does not vary by more than 10% Floods (indicated by 1:10 year event) do not reduce by more than 5% from present Base flows do not increase by more than 50% from present
Hydrodynamics	Maintain mouth state to create the required habitat for birds, fish, macrophytes, microalgae and water quality	Closed mouth state does not decrease by 10% from present Average water level in system > 10% from present Tidal amplitude (when open) < 20%
Water quality	Salinity distribution not to cause exceedance of TPCs for fish, invertebrates, macrophytes and microalgae Turbidity and dissolved oxygen not to cause exceedance of TPCs for biota DIN/DIP concentrations not to cause exceedance of TPCs for macrophytes and microalgae Toxic substances not to cause exceedance of TPCs for biota	Average salinity along estuary decreases by 5 below baseline average (to be determined) DO < 5 mg/l in estuary Turbidity > 20 NTU in low flow Secchi in fresher part: <0.5 m DIN >200 µg/l average (to be confirmed) DIP > 50 µg/l average (to be confirmed) Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAf, 1995) Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Sediment dynamics	Flood regime to maintain the sediment distribution patterns and aquatic habitat (instream physical habitat) so as not to exceed TPCs for biota Changes in sediment grain-size distribution patterns not to cause exceedance of TPCs in benthic invertebrates Change in average sediment composition and characteristics Change in average bathymetry	Average sediment composition (% fractions) along estuary change from baseline (to be measured) by 30% (per survey) Average depth along main channel changes from 30% of baseline (to be determined) (system expected to experience significant fluctuation in bathymetry between flood and extended closed periods)

Ecological component	EcoSpecs	Thresholds of Potential Concern
Microalgae	Maintain median phytoplankton/benthic microalgae biomass Prevent formation of phytoplankton blooms	Phytoplankton >8 µg/l (median) Benthic microalgae >42 mg/m ² (median) Phytoplankton >20 µg/l and/or cell density >10 000 cells/ml (once-off) Dinoflagellates, chlorophytes and/or cyanobacteria >10% of relative abundance
Macrophytes	Maintain distribution of macrophyte habitats Prevent the spread of reeds into open water Prevent an increase in nutrients and macroalgal blooms Prevent the spread of invasive trees (e.g. <i>Acacia</i> spp.) in the riparian zone. Maintain integrity of salt marsh	20% change in macrophyte area (Reeds currently cover 9 ha, saltmarsh 47 ha.) Macroalgal blooms cover > 50% of the open water area Presence of invasive aquatic macrophytes e.g. <i>Azolla</i> , water hyacinth Invasive plants cover > 10% of flood plain Increase in bare areas in salt marsh because of decrease in moisture and increase in salinity >30% of salt marsh
Invertebrates	Establish presence/absence of sand prawn <i>Callinectes kraussi</i> on sand banks in lower estuary Establish abundance of the copepod <i>Pseudodiaptomus hessei</i> or estuarine congeneric in the zooplankton of the estuary	If present populations deviate from average baselines (as determined in first three visits) by more 30%
Fish	Fish assemblage should comprise the five estuarine association categories in similar proportions (diversity and abundance) to that under the reference. Numerically, assemblage should comprise: Ia estuarine residents (20-60%) Ib marine and estuarine breeders (10-30%) IIa obligate estuarine-dependent (20-40%) IIb estuarine associated species (5-20%) IIc marine opportunists (20-80%) IV indigenous fish (1-5%) V catadromous species (1-5%) Category Ia species should contain viable populations of at least two species (e.g. <i>G.aestuaria</i> , <i>Hyporhamphus capensis</i> , <i>Omobranchus woodii</i>). Category IIa obligate dependents should be well represented by large exploited species (i.e. <i>A. japonicus</i> , <i>L. lithognathus</i> , <i>P. commersonii</i> , <i>Lichia amia</i>). REI species dominated by both <i>Myxus capensis</i> and <i>G. aestuaria</i> .	Ia estuarine residents <20% Ib marine and estuarine breeders < 10% IIa obligate estuarine-dependent <20% IIb estuarine associated species <5% IIc marine opportunists < 20% IV indigenous fish <1% V catadromous species <1% Ia represented only by <i>G. aestuaria</i> . IIa exploited species in very low numbers or absent REI species represented only by <i>G. aestuaria</i> , <i>Myxus capensis</i> absent
Birds	Maintain population of original groups of birds present on the estuary	Number of birds in any group, other than species that are increasing regionally such as Egyptian geese, drops below the baseline median (determined by past data and/or initial surveys) number of species and/or birds counted for three consecutive summer or winter counts

Table 7.4: Eco-Specs and Thresholds of Potential Concern for the Piesang Estuary

Ecological component	EcoSpecs	Thresholds of Potential Concern
Hydrology	Maintain present day base flow as a minimum (to be confirmed)	MAR does not vary by more than 10% Floods (indicated by 1:10 year event) do not reduce by more than 5% from present. Base flows do not increase by more than 50% from present
Hydrodynamics	Maintain mouth state to create the required habitat for birds, fish, macrophytes, microalgae and water quality	Closed mouth state increase by 10% from present Average water level in system > 10% from present Tidal amplitude (when open) < 20%
Water quality	Salinity distribution not to cause exceedance of TPCs for fish, invertebrates, macrophytes and microalgae Turbidity and dissolved oxygen not to cause exceedance of TPCs for biota DIN/DIP concentrations not to cause exceedance of TPCs for macro-phytes and microalgae Toxic substances not to cause exceedance of TPCs for biota	Salinity > 20 (expected range 10-20) Salinity < 5 (expected range 10-20) DO < 5 mg/l in estuary Turbidity > 10 NTU in low flow Secchi: to bottom DIN > 100 µg/l/once-off DIP > 20 µg/l/once-off Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Sediment dynamics	Flood regime to maintain the sediment distribution patterns and aquatic habitat (instream physical habitat) so as not to exceed TPCs for biota Changes in sediment grain-size distribution patterns not to cause exceedance of TPCs in benthic invertebrates Change in average sediment composition and characteristics Change in average bathymetry	Average sediment composition (% fractions) along estuary changes from baseline (to be measured) by 30% (per survey) Average depth along main channel changes from 30% of baseline (to be determine) (system expected to experience significant fluctuation in bathymetry between flood and extended closed periods)
Microalgae	Maintain median phytoplankton/benthic microalgae biomass Prevent formation of phytoplankton blooms	Phytoplankton > 3.5 µg/l (median) Benthic microalgae > 11 mg/m ² (median) Phytoplankton > 20 µg/l and/or cell density > 10 000 cells/ml (once-off)
Macrophytes	Maintain distribution of macrophyte habitats Prevent an increase in nutrient input leading to macroalgal blooms Control the spread of invasive plants in the riparian zone	Greater than 20 % change in the area covered by macrophytes (reeds and sedges currently cover 3.14 ha, submerged macrophytes and salt marsh present) Macroalgal blooms cover > 50% of the open water area during closed mouth conditions Invasive plants cover > 5% of total habitat
Invertebrates	Maintain presence of sand prawn <i>Callichirus kraussi</i> on sand banks in lower estuary Maintain presence of the copepod <i>Pseudodiaptomus hessei</i> or estuarine congeneric in the zooplankton of the estuary	Populations deviate from average baselines (as determined in first three visits) by more 30%

Ecological component	EcoSpecs	Thresholds of Potential Concern
Fish	<p>Fish assemblage should comprise the five estuarine association categories in similar proportions (diversity and abundance) to that under the reference. Numerically, assemblage should comprise:</p> <p>Ia estuarine residents (50-80% of total abundance)</p> <p>Ib marine and estuarine breeders (5-20%)</p> <p>Ila obligate estuarine-dependent (10-20%)</p> <p>Ilb estuarine associated species (5-15%),</p> <p>Ilc marine opportunists (20-80%)</p> <p>III marine vagrants (not more than 5%)</p> <p>IV indigenous fish (1-5%)</p> <p>V catadromous species (1-5%)</p> <p>Category Ia species should contain viable populations of at least two species (e.g. <i>G.aestuaria</i>, & <i>Hyporamphus capensis</i>).</p> <p>Category Ila obligate dependents should be well represented by at least two large exploited species (i.e. <i>L. lithognathus</i>, <i>Lichia amia</i>).</p> <p>REI species dominated by both <i>Myxus capensis</i> and <i>G. aestuaria</i>.</p>	<p>Ia estuarine residents <50%</p> <p>Ib marine and estuarine breeders <10%</p> <p>Ila obligate estuarine-dependent <10%</p> <p>Ilb estuarine associated species <5%</p> <p>Ilc marine opportunists < 20%</p> <p>III marine vagrants > 5%</p> <p>IV indigenous fish <1%</p> <p>V catadromous species <1%</p>
Birds	Maintain population of original groups of birds present on the estuary	Number of birds in any group, other than species that are increasing regionally such as Egyptian geese, drops below the baseline median (determined by past data and or initial surveys) number of species and/or birds counted for three consecutive summer or winter counts

Table 7.5: EcoSpecs for the Groot (Wes) Estuary

Ecological component	EcoSpecs	Thresholds of Potential Concern
Hydrology	Maintain present day base flow as a minimum (to be confirmed)	MAR does not vary by more than 10% Floods (indicated by 1:10 year event) do not reduce by more than 5% from present. Base flows do not increase by more than 50% from present
Hydrodynamics	Maintain mouth state to create the required habitat for birds, fish, macrophytes, microalgae and water quality	Closed mouth state increases by 10% from present Average water level in system > 10% from present Tidal amplitude (when open) < 20%
Water quality	<p>Salinity distribution not to cause exceedance of TPCs for fish, invertebrates, macrophytes and microalgae</p> <p>Turbidity and dissolved oxygen not to cause exceedance of TPCs for biota</p> <p>DIN/DIP concentrations not to cause exceedance of TPCs for macrophytes and microalgae</p> <p>Toxic substances not to cause exceedance of TPCs for biota</p>	<p>Average salinity along estuary decreases by 5 below baseline average (to be determined)</p> <p>Average salinity < 10at the head of the estuary (expected average range 5-10 for most of the system)</p> <p>DO < 5 mg/l in estuary</p> <p>Turbidity > 10 NTU in low flow</p> <p>Secchi: to bottom</p> <p>DIN > 100 µg/l once-off</p> <p>DIP > 20 µg/l once-off</p> <p>Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995)</p> <p>Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)</p>

Ecological component	EcoSpecs	Thresholds of Potential Concern
Sediment dynamics	Flood regime to maintain the sediment distribution patterns and aquatic habitat (instream physical habitat) so as not to exceed TPCs for biota Changes in sediment grain-size distribution patterns not to cause exceedance of TPCs in benthic invertebrates Change in average sediment composition and characteristics Change in average bathymetry	Average sediment composition (% fractions) along estuary changes from baseline (to be measured) by 30% (per survey) Average depth along main channel changes from 30% of baseline (to be determined) (system expected to experience significant fluctuation in bathymetry between flood and extended closed periods)
Microalgae	Maintain median phytoplankton/benthic microalgae biomass Prevent formation of phytoplankton blooms	Phytoplankton >3.5 µg/l (median) Benthic microalgae >11 mg/m ² (median) Phytoplankton >20 µg/l and/or cell density >10 000 cells/ml (once-off)
Macrophytes	Maintain distribution of macro-phyte habitats. Prevent an increase in nutrient input leading to macroalgal blooms. Control the spread of invasive plants in the riparian zone	Greater than 20 % change in the area covered by macro-phytes (reeds and sedges currently cover 2.54 ha salt marsh 0.76 ha) Macro-algal blooms cover > 50% of the open water area during closed mouth conditions. Invasive plants cover >5% of total habitat
Invertebrates	Establish presence/absence of sand prawn <i>Callichirus kraussi</i> on sand banks in lower estuary Establish presence/absence of the copepod <i>Pseudodiaptomus hessei</i> or estuarine congeneric in the zooplankton of the estuary	If present populations deviate from average baselines (as determined in first three visits) by more 30%
Fish	Fish assemblage should comprise the five estuarine association categories in similar proportions (diversity and abundance) to that under the reference. Numerically, assemblage should comprise: Ia estuarine residents (50-80% of total abundance) Ib marine and estuarine breeders (5-20%) IIa obligate estuarine-dependent (10-20%) IIb estuarine associated species (5-15%), IIc marine opportunists (20-80%) III marine vagrants (not more than 5%) IV indigenous fish (1-5%) V catadromous species (1-5%) Category Ia species should contain viable populations of at least two species (e.g. <i>G.aestuaria</i> , & <i>Hyporamphus capensis</i>). Category IIa obligate dependents should be well represented by at least two large exploited species (i.e. <i>L. lithognathus</i> , <i>Lichia amia</i>). REI species dominated by both <i>Myxus capensis</i> and <i>G. aestuaria</i> .	Ia estuarine residents <50% Ib marine and estuarine breeders < 10% IIa obligate estuarine-dependent <10% IIb estuarine associated species <5% IIc marine opportunists < 20% III marine vagrants > 5% IV indigenous fish <1% V catadromous species <1%
Birds	Maintain population of original groups of birds present on the estuary	Number of birds in any group, other than species that are increasing regionally such as Egyptian geese, drops below the baseline median (determined by past data and or initial surveys) number of species and/or birds counted for three consecutive summer or winter counts

Table 7.6: EcoSpecs and Thresholds of Potential Concern for the Bloukrans Estuary

Ecological component	EcoSpecs	Thresholds of Potential Concern
Hydrology	Maintain present flow regime	Varies more than 10% of MAR
Hydrodynamics	Maintain mouth state to create the required habitat for birds, fish, macro-phytes, microalgae and water quality	Estuary mouth closes
Water quality	Salinity distribution not to cause exceedance of TPCs for fish, invertebrates, macrophytes and microalgae Turbidity and dissolved oxygen not to cause exceedance of TPCs for biota DIN/DIP concentrations not to cause exceedance of TPCs for macrophytes and microalgae Toxic substances not to cause exceedance of TPCs for biota	Average salinity < 10 at the head of the estuary (expected average range >30 for most of the system) DO < 5 mg/l in estuary Turbidity > 10 NTU in low flow Secchi: to bottom DIN > 100 µg/l once-off DIP > 20 µg/l once-off Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Sediment dynamics	Flood regime to maintain the sediment distribution patterns and aquatic habitat (instream physical habitat) so as not to exceed TPCs for biota Changes in sediment grain-size distribution patterns not to cause exceedance of TPCs in benthic invertebrates Change in average sediment composition and characteristics Change in average bathymetry	Average sediment composition (% fractions) along estuary change from baseline (to be measured) by 30% (per survey) Average depth along main channel change from 30% of baseline (to be determine) (system expected to significant fluctuation in bathymetry between flood and extended closed periods)
Microalgae	Maintain median phytoplankton/benthic microalgae biomass Prevent formation of phytoplankton blooms	Phytoplankton > 1.0 µg/l (median) Benthic microalgae > 11 mg/m ² (median) Phytoplankton > 20 µg/l and/or cell density > 10 000 cells/ml (once-off)
Macrophytes	The estuary habitats only consists of sand/mud banks (0.63 ha) and channel (2.88 ha), no macrophytes	N/A
Invertebrates	Establish presence/absence of sand prawn <i>Callinectes kraussi</i> on sand banks in lower estuary Establish presence/absence of the copepod <i>Pseudodiaptomus hessei</i> or estuarine congeneric in the zooplankton of the estuary	If present populations deviate from average baselines (as determined in first three visits) by more 30%

Ecological component	EcoSpecs	Thresholds of Potential Concern
Fish	<p>Fish assemblage should comprise the five estuarine association categories in similar proportions (diversity and abundance) to that under the reference. Numerically, assemblage should comprise:</p> <p>Ia estuarine residents (50-80% of total abundance) Ib marine and estuarine breeders (10-20%) IIa obligate estuarine-dependent (10-20%) IIb estuarine associated species (5-15%), IIc marine opportunists (20-80%) III marine vagrants (not more than 5%) IV indigenous fish (1-5%) V catadromous species (1-5%)</p> <p>Category Ia species should contain viable populations of at least 4 species (<i>G.aestuarina</i>, <i>Hyporhamphus capensis</i>, <i>Omobranchus woodii</i>).</p> <p>Category IIa obligate dependents should be well represented by large exploited species (<i>A. japonicus</i>, <i>L. lithognathus</i>, <i>P. commersonii</i>, <i>Lichia amia</i>).</p> <p>REI species dominated by both <i>Myxus capensis</i> and <i>G. aestuarina</i>.</p>	<p>Ia estuarine residents <50% Ib marine and estuarine breeders < 10% IIa obligate estuarine-dependent < 10% IIb estuarine associated species < 5% IIc marine opportunists < 20% III marine vagrants > 5% IV indigenous fish < 1% V catadromous species <1%</p>
Birds	Maintain population of original groups of birds present on the estuary	Number of birds in any group, other than species that are increasing regionally such as Egyptian geese, drops below the baseline median (determined by past data and or initial surveys) number of species and/or birds counted for three consecutive summer or winter counts

Table 7.7: EcoSpecs and TPCs for the Goukou Estuary

EcoSpecs	TPC
Water quality	
Salinity distribution not to cause exceedance of TPCs for biota.	<ul style="list-style-type: none"> ▪ Salinity > 0 at head of estuary. ▪ Average salinity in Zone D > 5. ▪ Average salinity in Zone C > 20. ▪ Average salinity 5 km upstream from mouth > 20 more than three months of the year.
System variables (pH, dissolved oxygen and turbidity) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ 6.0 < pH > 7.5. ▪ DO < 5 mg/L. ▪ Suspended solids >5 mg/L (low flow). <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Average turbidity >10 Nephelometric Turbidity Units (NTU) (low flow). ▪ Average 6.0 < pH > 8.5 (increasing with increase in salinity). ▪ Average DO < 5 mg/L.
Inorganic nutrient concentrations (NO ₃ -N, NH ₃ -N and PO ₄ -P) not to cause in exceedance of TPCs for macrophytes and microalgae.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ NO_x-N >150 µg/L over two consecutive months. ▪ NH₃-N > 20 µg/L over two consecutive months. ▪ PO₄-PP > 20 µg/L over two consecutive months. <p>Estuary (except during upwelling or floods):</p> <ul style="list-style-type: none"> ▪ Average NO_x-N 150 µg/L single concentration > 200 µg/L.

EcoSpecs	TPC
	<ul style="list-style-type: none"> ▪ Average NH₃-N > 20 µg/L during survey, single concentration > 100 µg/L. ▪ Average PO₄-P > 20 µg/L during survey, single concentration > 50 µg/L.
Presence of toxic substances (e.g. trace metals and pesticides/herbicides) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ Trace metals (to be refined and confirmed through future monitoring). ▪ Pesticides/herbicides (to be refined and confirmed through future monitoring). <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Concentrations in water column exceed target values as per SA Water Quality Guidelines for Coastal Marine Waters (DWAF, 1995). ▪ Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

Table 7.8: Water Quality EcoSpecs and TPCs for the Gouritz Estuary

Component	EcoSpecs	Thresholds of Potential Concern
Water Quality	Salinity distribution not to cause exceedance of TPCs for biota.	<ul style="list-style-type: none"> ▪ Salinity > 0 at head of estuary ▪ Average salinity in Site 11, 1 km upstream of bridge > 5 ▪ Average salinity in Zone C > 20 ▪ Average salinity 11 km upstream from mouth > 20 more than three months of the year ▪ Salinity > 40 in saltmarsh sediments (linked to decrease in moisture and drying of floodplain habitat).
	System variables (pH, dissolved oxygen and turbidity) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ 7.0 < pH > 8.3 ▪ DO < 5 mg/l ▪ Suspended solids > 5 mg/l (low flow) <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Average turbidity > 10 NTU (low, calm condition flow, wind mixing can increase turbidity to 20-40 NTU) ▪ Average 7.0 < pH > 8.5 (increasing with increase in salinity) ▪ Average DO < 5 mg/l
	Inorganic nutrient concentrations (NO ₃ -N, NH ₃ -N and PO ₄ -P) not to cause exceedance of TPCs for macrophytes and microalgae.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ NO_x-N > 100 µg/l over two consecutive months ▪ NH₃-N > 20 µg/l over two consecutive months ▪ PO₄-P > 20 µg/l over two consecutive months <p>Estuary (except during upwelling or floods):</p> <ul style="list-style-type: none"> ▪ Average NO_x-N > 100 µg/l single concentration > 150 µg/l ▪ Average NH₃-N > 20 µg/l during survey, single concentration > 100 µg/l ▪ Average PO₄-P > 20 µg/l during survey, single concentration > 50 µg/l
	Presence of toxic substances (e.g. trace metals and pesticides/herbicides) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ Trace metals (to be confirmed) ▪ Pesticides/herbicides (to be confirmed) <p>Estuary</p> <ul style="list-style-type: none"> ▪ Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) <p>Concentrations in sediment exceed target values as per Western Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)</p>

Table 7.9: Eco Specs and TPCs for the Klein Brak Estuary

Component	EcoSpecs	Thresholds of Potential Concern
Water quality	Salinity distribution not to cause exceedance of TPCs for biota.	<ul style="list-style-type: none"> ▪ No salinity gradient in the upper reaches of the estuary (Zone D and F) ▪ No REI in the upper reaches of the estuary (Zone D and F) ▪ Salinity > 35
	System variables (pH, dissolved oxygen and turbidity) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ 7.0 < pH > 8.5 ▪ DO < 5 mg/l ▪ Suspended solids > 5 mg/ l (low flow) <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Average turbidity >10 NTU (low flow) ▪ Average 7.0 < pH > 8.5 (increasing with increase in salinity) ▪ Average DO < 5 mg/l
	Inorganic nutrient concentrations (NO ₃ -N, NH ₃ -N and PO ₄ -P) not to cause in exceedance of TPCs for macro-phytes and microalgae.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ NO_x-N >150 µg/l over two consecutive months ▪ NH₃-N > 20 µg/l over two consecutive months ▪ PO₄-P > 20 µg/l over two consecutive months <p>Estuary (except during upwelling or floods):</p> <ul style="list-style-type: none"> ▪ Average NO_x-N > 150 µg/l during survey, single concentration > 200 µg/l ▪ Average NH₃-N > 20 µg/l during survey, single concentration > 100 µg/l ▪ Average PO₄-P > 20 µg/l during survey, single concentration > 50 µg/l
	Presence of toxic substances (e.g. trace metals and pesticides/herbicides) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ Trace metals (to be confirmed) ▪ Pesticides/herbicides (to be confirmed) <p>Estuary</p> <ul style="list-style-type: none"> ▪ Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

Table 7.10: EcoSpecs and TPCs for the Wilderness System

Component	EcoSpecs	Thresholds of Potential Concern
Water quality	Salinity	<p>Estuary in the closed state:</p> <ul style="list-style-type: none"> ▪ Average salinity in Zone A < 12, ▪ Average salinity in Zone B: < 10 ▪ Average salinity in Zone C < 5 <p>Lakes average salinity +2 from baseline (2013) and variability do not increase as below:</p> <ul style="list-style-type: none"> ▪ Serpentine: 12 ± 10 ▪ Eilandvlei: 8 ± 5 ▪ Langvlei: 10 ± 4 ▪ Rondevlei: 10 ± 5

	System variables (pH, dissolved oxygen and turbidity) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ 6.0 < pH > 7.0 (Touw) ▪ 7.0 < pH > 8.0 (Duiwe) ▪ DO < 5 mg/l ▪ Suspended solids > 5 mg/ l (low flow) <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Average turbidity > 5 NTU (low flow) ▪ Average 6.0 < pH > 8.5 (increasing with increase in salinity) ▪ Average DO < 5 mg/l <p>Lakes:</p> <ul style="list-style-type: none"> ▪ Average turbidity > 5 NTU ▪ Average 7.0 < pH > 8.5 ▪ Average DO < 5 mg/l
	Inorganic nutrient concentrations (NO ₃ -N, NH ₃ -N and PO ₄ -P) not to cause exceedance of TPCs for macro-phytes and microalgae.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ NO_x-N > 50 µg/l over two consecutive months ▪ NH₃-N > 10 µg/l over two consecutive months ▪ PO₄-P > 10 µg/l over two consecutive months <p>Estuary (except during upwelling or floods):</p> <ul style="list-style-type: none"> ▪ Average NO_x-N > 50 µg/l single concentration > 100 µg/l ▪ Average NH₃-N > 10 µg/l during survey, single concentration > 100 µg/l ▪ Average PO₄-P > 10 µg/l during survey, single concentration > 50 µg/l <p>Lakes:</p> <ul style="list-style-type: none"> ▪ Average NO_x-N > 50 µg/l during survey, single concentration > 100 µg/l ▪ Average NH₃-N > 20 µg/l during survey (to be confirmed) ▪ Average PO₄-P > 20 µg/l during survey (to be confirmed)
	Presence of toxic substances (e.g. trace metals and pesticides/herbicides) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ Trace metals (to be confirmed) ▪ Pesticides/herbicides (to be confirmed) <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

Table 7.11: Water quality present state assessment for H8DUIW-EWR1

Water Quality Constituents	PES Value	Category/Comment
Inorganic salt ions (mg/l)		
Sulphate as SO ₄	N/A	-
Sodium as Na	382.2	Exceeds the ≤ 70 mg/L (TWQR) for Agricultural Use: Irrigation.
Magnesium as Mg	67.4	No guideline.
Calcium as Ca	55.0	No guideline.
Chloride as Cl	805.4	Exceeds the ≤ 100 mg/L (TWQR) for Agricultural Use: Irrigation.
Potassium as K	9.25	No guideline.
Electrical conductivity (mS/m)		
	272	E/F: RC = 80 mS/m.
Nutrients (mg/l)		
SRP	0.014	A
TIN	0.118	A

Water Quality Constituents	PES Value	Category/Comment
Physical variables		
pH (5 th + 95 th %ile)	6.6 and 8.1	B
Temperature (°C)	N/A	A/B. Impacts expected at low flows.
Dissolved oxygen (mg/L)	N/A	B. Impacts expected at low flows.
Turbidity (NTU)	N/A	B. Changes in turbidity appear to be largely related to natural with minor man-made modifications, e.g. gravel mining upstream
Response variables		
Chl-a: phytoplankton (ug/L)	N/A	N/A
Macroinvertebrate score (MIRAI) SASS score ASPT score	50.7% 78 56	D
Diatoms	11.1	C/D (n = 1, Jan 2014)
Fish score (FRAI)	51.6%	D (all estuarine spp. that moved into the freshwater zone and aliens).
Toxics		
Ammonia (as N)	0.003	A
Fluoride (as F)	0.33	A
OVERALL SITE CLASSIFICATION (PAI model)		C (73.2%)

(a) N/A- No data were available for this assessment.

Table 7.12: H8DUIW-EWR1: Water quality (C category) EcoSpecs and TPCs

Metrics	EcoSpecs	TPCs
Inorganic salt ions		
Sulphate as SO ₄	N/A	N/A
Sodium as Na	The 95 th percentile of the data must be ≤ 380 mg/L.	The 95 th percentile of the data must be 300 - 380 mg/L.
Magnesium as Mg	The 95 th percentile of the data must be ≤ 67 mg/L.	The 95 th percentile of the data must be 53.5 - 67 mg/L.
Calcium as Ca	The 95 th percentile of the data must be ≤ 55 mg/L.	The 95 th percentile of the data must be 44 - 55 mg/L.
Chloride as Cl	The 95 th percentile of the data must be ≤ 800 mg/L.	The 95 th percentile of the data must be 640 - 800 mg/L.
Potassium as K	The 95 th percentile of the data must be ≤ 9 mg/L.	The 95 th percentile of the data must be 7 - 9 mg/L.
Physical Variables		
Electrical conductivity (mS/m)	The 95 th percentile of the data must be ≤ 270 mS/m.	The 95 th percentile of the data must be 210 - 270 mS/m.
pH	The 5 th percentile of the data must be 6.5 - 8.0, and the 95 th percentile 8.0 - 8.8.	The 5 th percentile of the data is ≤ 6.3 and the 95 th percentile is ≥ 8.6.
Temperature ^(a)	Natural temperature range.	Initiate baseline monitoring for this variable.
Dissolved oxygen ^(a) (DO)	The 5 th percentile of the data must be ≥ 7.0 mg/L.	The 5 th percentile of the data must be 7.2 - 7.0 mg/L. Initiate baseline monitoring for this variable.
Turbidity ^(a)	Changes in turbidity are related to minor man-made modifications (e.g. gravel mining upstream). Some silting of habitats is expected.	Initiate baseline monitoring for this variable.
Nutrients		

Metrics	EcoSpecs	TPCs
TIN	The 50 th percentile of the data must be ≤ 0.25 mg/L.	The 50 th percentile of the data must be 0.2 - 0.25 mg/L.
PO ₄ -P	The 50 th percentile of the data must be ≤ 0.015 mg/L.	The 50 th percentile of the data must be 0.012 - 0.015 mg/L.
Response variables		
Chl-a phytoplankton	The 50 th percentile of the data must be < 15 µg/L.	The 50 th percentile of the data must be 12 - 15 µg/L.
Chl-a periphyton	The 50 th percentile of the data must be ≤ 12 mg/m ² .	The 50 th percentile of the data must be 10 - 12 mg/m ² .
Toxics		
Fluoride	The 50 th percentile of the data must be ≤ 1.5 mg/L.	The 50 th percentile of the data must be 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile of the data must be ≤ 0.015 mg/L.	The 50 th percentile of the data must be 0.012 - 0.015 mg/L.
Other toxics	The 95 th percentile of the data must be within the Target Water Quality Range (TWQR) as stated in DWAF (1996) or the A Category boundary as stated in DWAF (2008).	An impact is expected if the 95 th percentile of the data exceeds the TWQR as stated in DWAF (1996) or the upper limit of the A Category boundary as stated in DWAF (2008).

(a) N/A- No data were available for this assessment

Table 7.13: EcoSpecs and TPCs for the Duiwenhoks Estuary

Component	EcoSpecs	Thresholds of Potential Concern
Water quality	Salinity distribution not to cause exceedance of TPCs for biota.	<ul style="list-style-type: none"> ▪ Salinity > 0 at head of estuary ▪ Average salinity in Zone D > 5 ▪ Average salinity in Zone C > 20 ▪ Average salinity 5 km upstream from mouth > 20 more than three months of the year
	System variables (pH, dissolved oxygen and turbidity) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ 6.0 < pH > 7.5 ▪ DO < 5 mg/l ▪ Suspended solids > 5 mg/l (low flow) <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Average turbidity > 10 NTU (low flow) ▪ Average 6.0 < pH > 8.5 (increasing with increase in salinity) ▪ Average DO < 5 mg/l
	Inorganic nutrient concentrations (NO ₃ -N, NH ₃ -N and PO ₄ -P) not to cause an exceedance of TPCs for macrophytes and microalgae.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ NO_x-N > 150 µg/l over 2 consecutive months ▪ NH₃-N > 20 µg/l over 2 consecutive months ▪ PO₄-P > 20 µg/l over 2 consecutive months <p>Estuary (except during upwelling or floods):</p> <ul style="list-style-type: none"> ▪ Average NO_x-N > 150 µg/l single concentration > 200 µg/l ▪ Average NH₃-N > 20 µg/l during survey, single concentration > 100 µg/l ▪ Average PO₄-P > 20 µg/l during survey, single concentration > 50 µg/l
	Presence of toxic substances (e.g. trace metals and pesticides/herbicides) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ Trace metals (to be confirmed) ▪ Pesticides/herbicides (to be confirmed) <p>Estuary</p> <ul style="list-style-type: none"> ▪ Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

Table 7.14: EcoSpecs and TPCs for the Goukou Estuary

EcoSpecs	TPC
Water quality	
Salinity distribution not to cause exceedance of TPCs for biota.	<ul style="list-style-type: none"> ▪ Salinity > 0 at head of estuary. ▪ Average salinity in Zone D > 5. ▪ Average salinity in Zone C > 20. ▪ Average salinity 5 km upstream from mouth > 20 more than three months of the year.
System variables (pH, dissolved oxygen and turbidity) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ 6.0 < pH < 7.5. ▪ DO < 5 mg/L. ▪ Suspended solids >5 mg/L (low flow). <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Average turbidity >10 Nephelometric Turbidity Units (NTU) (low flow). ▪ Average 6.0 < pH < 8.5 (increasing with increase in salinity). ▪ Average DO < 5 mg/L.
Inorganic nutrient concentrations (NO ₃ -N, NH ₃ -N and PO ₄ -P) not to cause in exceedance of TPCs for macrophytes and microalgae.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ NO_x-N >150 µg/L over two consecutive months. ▪ NH₃-N > 20 µg/L over two consecutive months. ▪ PO₄-PP > 20 µg/L over two consecutive months. <p>Estuary (except during upwelling or floods):</p> <ul style="list-style-type: none"> ▪ Average NO_x-N 150 µg/L single concentration > 200 µg/L. ▪ Average NH₃-N > 20 µg/L during survey, single concentration > 100 µg/L. ▪ Average PO₄-P > 20 µg/L during survey, single concentration > 50 µg/L.
Presence of toxic substances (e.g. trace metals and pesticides/herbicides) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ Trace metals (to be refined and confirmed through future monitoring). ▪ Pesticides/herbicides (to be refined and confirmed through future monitoring). <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Concentrations in water column exceed target values as per SA Water Quality Guidelines for Coastal Marine Waters (DWAF, 1995). ▪ Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

Table 7.15: Water Quality EcoSpecs and TPCs for the Gouritz Estuary

Component	EcoSpecs	Thresholds of Potential Concern
Water Quality	Salinity distribution not to cause exceedance of TPCs for biota.	<ul style="list-style-type: none"> ▪ Salinity > 0 at head of estuary ▪ Average salinity in Site 11, 1 km upstream of bridge > 5 ▪ Average salinity in Zone C > 20 ▪ Average salinity 11 km upstream from mouth > 20 more than three months of the year ▪ Salinity > 40 in saltmarsh sediments (linked to decrease in moisture and drying of floodplain habitat).
	System variables (pH, dissolved oxygen and turbidity) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ 7.0 < pH > 8.3 ▪ DO < 5 mg/l ▪ Suspended solids > 5 mg/l (low flow) <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Average turbidity > 10 NTU (low, calm condition flow, wind mixing can increase turbidity to 20-40 NTU) ▪ Average 7.0 < pH > 8.5 (increasing with increase in salinity) ▪ Average DO < 5 mg/l
	Inorganic nutrient concentrations (NO ₃ -N, NH ₃ -N and PO ₄ -P) not to cause exceedance of TPCs for macrophytes and microalgae.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ NO_x-N > 100 µg/l over two consecutive months ▪ NH₃-N > 20 µg/l over two consecutive months ▪ PO₄-P > 20 µg/l over two consecutive months <p>Estuary (except during upwelling or floods):</p> <ul style="list-style-type: none"> ▪ Average NO_x-N > 100 µg/l single concentration > 150 µg/l ▪ Average NH₃-N > 20 µg/l during survey, single concentration > 100 µg/l ▪ Average PO₄-P > 20 µg/l during survey, single concentration > 50 µg/l
	Presence of toxic substances (e.g. trace metals and pesticides/herbicides) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ Trace metals (to be confirmed) ▪ Pesticides/herbicides (to be confirmed) <p>Estuary</p> <ul style="list-style-type: none"> ▪ Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAf, 1995) ▪ Concentrations in sediment exceed target values as per Western Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

Table 7.16: EcoSpecs and TPCs for the Wilderness System

Component	EcoSpecs	Thresholds of Potential Concern
Water quality	Salinity	<p>Estuary in the closed state:</p> <ul style="list-style-type: none"> ▪ Average salinity in Zone A < 12, ▪ Average salinity in Zone B: < 10 ▪ Average salinity in Zone C < 5 <p>Lakes average salinity +2 from baseline (2013) and variability do not increase as below:</p> <ul style="list-style-type: none"> ▪ Serpentine: 12 ± 10 ▪ Eilandvlei: 8 ± 5 ▪ Langvlei: 10 ± 4 ▪ Rondevlei: 10 ± 5
	System variables (pH, dissolved oxygen and turbidity) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ 6.0 < pH > 7.0 (Touw) ▪ 7.0 < pH > 8.0 (Duiwe) ▪ DO < 5 mg/l ▪ Suspended solids > 5 mg/l (low flow) <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Average turbidity > 5 NTU (low flow) ▪ Average 6.0 < pH > 8.5 (increasing with increase in salinity) ▪ Average DO < 5 mg/l <p>Lakes:</p> <ul style="list-style-type: none"> ▪ Average turbidity > 5 NTU ▪ Average 7.0 < pH > 8.5 ▪ Average DO < 5 mg/l
	Inorganic nutrient concentrations (NO ₃ -N, NH ₃ -N and PO ₄ -P) not to cause exceedance of TPCs for macro-phytes and microalgae.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ NO_x-N > 50 µg/l over two consecutive months ▪ NH₃-N > 10 µg/l over two consecutive months ▪ PO₄-P > 10 µg/l over two consecutive months <p>Estuary (except during upwelling or floods):</p> <ul style="list-style-type: none"> ▪ Average NO_x-N > 50 µg/l single concentration > 100 µg/l ▪ Average NH₃-N > 10 µg/l during survey, single concentration > 100 µg/l ▪ Average PO₄-P > 10 µg/l during survey, single concentration > 50 µg/l <p>Lakes:</p> <ul style="list-style-type: none"> ▪ Average NO_x-N > 50 µg/l during survey, single concentration > 100 µg/l ▪ Average NH₃-N > 20 µg/l during survey (to be confirmed) ▪ Average PO₄-P > 20 µg/l during survey (to be confirmed)
	Presence of toxic substances (e.g. trace metals and pesticides/herbicides) not to cause exceedance of TPCs for biota.	<p>River inflow:</p> <ul style="list-style-type: none"> ▪ Trace metals (to be confirmed) ▪ Pesticides/herbicides (to be confirmed) <p>Estuary:</p> <ul style="list-style-type: none"> ▪ Concentrations in water column exceed target values as per SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Concentrations in sediment exceed target values as per WIO Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

8. WETLANDS

Table 8.1: Wetlands - Quaternary Protection Specification

Quaternary catchment	EIS	PES	REC	How to achieve the REC
K10A	Moderate	C	C	Control invasive alien vegetation, erosion and land-use encroachment.
K10B	Moderate	C	C	
K10C	Moderate	B/C	B/C	
K10D	Moderate	B/C	B/C	
K10E	Moderate	B/C	B/C	
K10F	Moderate	C	C	
K20A	Moderate	C	C	
K30A	High	C	C	Buffers in urban and agricultural areas, manage water quality, erosion and invasive vegetation.
K30B	High	D	C/D	
K30C	Moderate	D	D	Control invasive alien vegetation, erosion and land-use encroachment.
K30D	Very High	B	B	
K40A	Moderate	D	D	
K40B	Moderate	C	C	
K40C	Moderate	C	C	
K40D	Very High	B	B	
K40E	Moderate	B/C	B/C	
K50A	Moderate	B/C	B/C	Protect and improve the condition of remaining wetland patches, control invasive vegetation.
K50B	High	C/D	C	
K60A	Moderate	B	B	Control invasive alien vegetation, erosion and land-use encroachment.
K60B	Moderate	B	B	
K60C	Moderate	B	B	
K60D	High	A	A	
K60E	High	C	C	
K60F	High	C	C	
K60G	Moderate	C	C	
K70A	Moderate	C	C	
K70B	Low	A	A	
H80A	HIGH	C/D	C	
H80B	Moderate	C	C	
H80C	Moderate	D	D	
H80D	Moderate	D	D	
H80E	Moderate	C/D	C/D	
H90A	Moderate	C	C	

Quaternary catchment	EIS	PES	REC	How to achieve the REC
H90B	Moderate	D	D	Control invasive alien vegetation, erosion and land-use encroachment.
H90C	Moderate	D	D	
H90D	Moderate	C	C	
H90E	Moderate	C/D	C/D	
J11D	Low	C	C	
J11F	Moderate	C	C	
J11G	Moderate	B	B	
J12A	Moderate	B	B	
J12B	Moderate	B	B	
J12J	Moderate	B	B	
J12K	Moderate	B	B	
J12L	Moderate	C	C	
J21A	Moderate	B/C	B/C	
J21B	Moderate	B	B	
J22B	Moderate	B	B	
J22G	Moderate	B	B	
J22K	Low	B/C	B/C	
J23E	Low	C	C	
J23J	Moderate	B	B	
J24F	Low	C	C	
J25A	Low	B	B	
J33B	Low	C	C	
J33E	Low	C	C	
J34C	Low	C	C	
J34D	Low	C	C	
J34E	Low	C/D	C/D	
J34F	Low	D	D	
J40B	Low	B	B	
J40C	Moderate	C/D	C/D	
J40D	Moderate	D	D	
J40E	High	C	C	




Figure 1: Locations of Gouritz region river/estuary nodes and EWR sites




Figure 2: Locations of Breede-Overberg region river/estuary nodes and EWR sites

ISAZISO SIKARHULUMENTE

ISAZISO _____ SIK2022**ISEBE LEZAMANZI NOGUTYULO****UMTHETHO WEZAMANZI WESIZWE, KA1998
(UMTHETHO NO. 36 KA1998)****UHLELO LWEMIJELO YAMANZI KWIMILAMBO YOMMANDLA WOLAWULO
LWAMANZI I- BREEDE-GOURITZ**

Mna, Senzo Mchunu, kwisikhundla sam njengoMphathiswa weSebe lezaManzi noGutyulo, ndigunyaziswa yimiqathango yesiqendu 16 (1) soMthetho wezaManzi weSizwe, ka1998 (Umthetho No. 36 ka1998), ukuba ndibhengeze amahlelo emijelo yamanzi ekuMmandla woLawulo lwaManzi i-Breede-Gouritz njengoko kubonisiwe kwiShedyuli yesi Saziso.

Silungiswe nguMphathi: weCandelo lokuHlelwa kweMijelo yaManzi
Makubhalelwe ku: Mnu Yakeen Atwaru
WeSebe lezaManzi noGutyulo
Ndinaye Building 185 Francis Baard Street
Private Bag X313
Pretoria
0001
Email: atwaruy@dws.gov.za



NISIPHATHELWA NGUSENZO MCHUNU (MP)
UMPHATHISWA WESEBE LEZAMANZI NOGUTYULO
UMHLA: 15/08/22

UHLELO LWEMIJELO YAMANZI KWIMILAMBO YOMMANDLA WOLOWULO LWAMANZI I-BREED-GOURITZ NGOKWEMIQATHANGO YESIQENDU 6(1) NO (2) SOMTHETHO WEZAMANZI WESIZWE, KA1998 (UMTHETHO NO. 36 KA1998)

ISHEDULI

1. INKCAZO NGOMJELO WAMANZI

- 1.1 Xa sithetha ngomjelo sixela yonke into engumlambo ekuMmandla woLawulo lwaManzi i Breede-Gouritz, ngolu hlobo lubonisiweyo apha ngezantsi:

UMmandla woLawulo lwaManzi: iBreede-Gouritz
 IiNgingqi zoFunxo: INgingqi ePhakamileyo yoFunxo enguG40-G50, uH10- H90, uJ11-J40, noK10-K70

Imilambo: EkuMmandla iBreede Overberg: Umlambo iBreede, umlambo iRiviersonderend, umlambo i-Overberg, neminye ke imilanjana eselunxwemeni.
 EkuMmandla wonxweme iGouritz: Umlambo iGouritz, umlambo iBuffels, umlambo iTouws, umlambo iGroot, umlambo iGamka, umlambo i-Olifants, umlambo iKammanassie, neminye ke imilanjana eselunxwemeni.

- 1.2 UMphathiswa, ngokwesiqendu 12 soMthetho waManzi weSizwe, ka1998 (UMthetho No.36 ka 1998) ("uMthetho"), uxele indlela yokuhlela imijelo yamanzi ngokuthi akhuphe iSaziso sikaRhulumente esingu- No. R. 810, esithe sabhengezwa kwincwadi yeZaziso zikaRhulumente engu- No. 33541 neshicilelwe ngomhla we-17 kweyoMsintsi ku2010. Ngokwemiqathango yesiqendu 6(1) soMthetho, uMphathiswa – kanye nje emva kokuba ihlelo lawo onke amanqanaba (okanye lezo ndawana nje) z/omjelo wamanzi ngamnye liqingqiwe – makabhengenze ezo mpawu ziqingqiweyo zaloo Mjelo uthile kwiSaziso sikaRhulumente.

- 1.3 uMphathiswa, ngokwemiqathango isiqendu 16(1) soMthetho, sokuba kuqatshelwe ezi mpawu zilandelayo ngeMijelo ekuMmandla woLawulo lwaManzi iBreede-Gouritz.

2. NAZI IIPAWU ZOMJELO EZIPHAKANYISWAYO NEZIQINGQWE NGOKWEMIQATHANGO YESIQENDU 16(1) NO- (2) SOMTHETHO WEZAMANZI WESIZWE, KA1998

- 2.1 Iimpawu zoMjelo eziphakanyiswayo – nezibandakanya iiMfuno zaManzi zeNdalo (ii-Ecological Water Requirements ngelasemzini, ii- EWRs ngamafutshane) neMijelo yeeMfuno ezisiSiseko zoLuntu (ii-Basic Human Needs Reserve ngelasemzini, iiBHN ngamafutshane) kwimilambo ekwizikhundla ze-EWR nekwezo ndawo zikhethiweyo zendalo kuMmandla woLawulo lwaManzi iBreede-Gouritz – zibonisiwe **Sahluko 4**. IiNdawo zoBoniselo ngaManzi iGouritz neBreede-Overberg nezikhundla ze-EWR zibonisiwe phaya **kwiMizobo 1 no- 2 ngokulandelelanayo**.

- 2.2 Ibakala elifana neKwaliti yaManzi (phakathi kwezo mpawu zoMjelo ziphakanyiswayo ngaloo milambo ikwizikhundla ze-EWR kuMmandla woLawulo lwaManzi i-Breede-Gouritz ngokwemiqathango yesiqendu 16(1) soMthetho) libonisiwe phaya **kwiSahluko 5**.

- 2.3 Iimpawu eziphakanyiswayo zoMjelo wokuqokelela amanzi angaphantsi komhlaba ngokoMthamo neKwaliti yamanzi (njengoko kuxeliwe kwimiqathango yesiqendu 16(1) soMthetho malunga noMmandla woLawulo lwaManzi Breede-Gouritz) zibonisiwe phaya **kwiSahluko 6**.

2.4 Iimpawu zoMjelo oliChweba eziphakanyiswayo ngokwemiqathango yesiqendu 16(1) soMthetho kuMmandla woLawulo lwaManzi i-Breede-Gouritz zibonisiwe phaya **kwiSahluko 7**.

2.5 Iimpawu zoMjelo onguMwonyo eziphakanyisiweyo kuMmandla woLawulo lwaManzi i-Breede-Gouritz ngokwemiqathango yesiqendu 16(1) soMthetho zibonisiwe phaya **kwiSahluko 8**.

2.6 Ezi mpawu ziqingqiweyo ngoMjelo ngamnye ziza kuqala ukusebenza ukusukela ngomhla ezityikitywe ngawo njengoko isitsho imiqathango yesiqendu 16(1) somthetho, ngaphandle kokuba uMphathiswa ubone ngenye indlela.

3. IZISHUNQULELO ZAMAGAMA NEENKCAZO

3.1 Izishunqulelo

Isishunqulelo	Inkcazo
BHN	Basic Human Needs (iiMfuno ezisiSiseko zoLuntu)
EC	Ecological Category (Inqanaba lendalo)
EcoSpecs	Ecological Specifications (Imiqathango yendalo)
EIS	Ecological Importance and Sensitivity (Undoqo novakalelo lwendalo)
ER	Ecological Reserve (Umjelo wendalo)
EWR	Ecological Water Requirement (IMfuno yaManzi yeNdalo)
MAR	Mean Annual Runoff (Ubuncikane bamanzi emvula ngonyaka)
MCM	ii-Million Cubic Metres
PES	Present Ecological Status (isimo sangoku sendalo)
RC	Reference conditions (iimeko ekubhekiswa kuzo)
REC	Recommended Ecological Category (Icandelo lendalo elindululwayo/elixhaswayo)
TPCs	Thresholds of Potential Concern (Imilinganiselo yenkathazo enokuvela)
WQSU	Water quality sub-unit (icandelwana lekwaliti yamanzi)

3.2 Inkcazo

Inketshenkeshana yamanzi/ ulwelo: Ngalaa manzana asalayo emilanjeni ngexesha leemeko ezifana nemozulu engumqwebedu okanye leyo ingacacanga, kodwa ke ingengawo lawo asuka phantsi komhlaba. La manzana aquka lawo angena ngolibaziseko oluthile emilanjeni nalawo ke atsitisa nje ngaphantsi komhlaba.

Undoqo novakalelo lwendalo (i-EIS): Ezi ke zizalathisi/ziimpawu eziphambili ezijongwayo xa kuhlelwa imijelo yamanzi yendalo. Undoqo wendalo ubhekisa kwiimpawu ezifana nobukho, ubumeli nokwahluka kwemigqeku ye- biota nendawo yokuphilisana. Uvakalelo lwendalo lubhekisa kubungozi obunokwenzeka kwindawo yokuphilisana okanye kwi-biota xa kuthe kwenzeka inguqu kwindlela ahamba ngayo amanzi, kumanqanaba amanzi kanti nakwezo meko zibangelwa bubukho bekhemikhali kuloo ndawo ithile.

Iimfuno zamanzi zendalo (i-EWR): lipethini zokuhamba kwamanzi (oko kukuthi ubungakanani, uhlalutyo lwexesha nelixa lobukho) nekwaliti yamanzi edingekayo ukuze kugcinwe impilo yasemanzini intle phantsi kweemeko ezithile. Esi sigama sisetyenziswa kakhulu xa kubhekiswa kwizinto ezifana namabakala omthamo nekwaliti.

Izikhundla zeemfuno zamanzi zendalo (ze-EWR): La ngamanqanaba omlambo athi aqingqwe ngaloo maxa kukhethwa isikhundla. Isikhundla se-EWR senziwa bubude bomlambo obunokuquka iindawana zokudlula kwezinto zasemanzini nezinye izinto zendalo. Ezi zikhundla zinezalathisi ezaneleyo zokuvavanya indlela ahamba ngayo amanzi zibuye zihlole namabakala ezifundo ezifunekayo ngentshukumo yamanzi (i-hydrology), ngokwakhekha komhlaba nangefuthe lobukho beekhemikhali nelohlobo oluthile lwendalo (umzekelo kwiintlanzi, kwizinto eziphilayo ezingenamathambo nakutyani waselunxwemeni).

Isimo sangoku sendalo (i-PES): Eli bakala libonisa impilo nemfezeko yangoku yeempawu ezininzi ezibonakalayo kuloo mjelo uthile wamanzi, ngakumbi xa uzithelekisa neemeko ezibhekiselele kwezo zendalo. Iziphumo zohlolo lwesimo sangoku ziye ziboniswe ngokusebenzisa amaBakala eNdalo (ii-Ecological Categories ngelasemzini – ii-ECs ngamafutshane) asukela ku-A (ochaza ukuba umjelo usekwisimo sendalo) ze aye kutsho ku-F (ochaza ukuba isimo somjelo siguquke ngokupheleleyo) xa kuthethwa nge-PES.

Uvuselelo: Oku kuxela ukuba kongezwa amanzi apho ebesele ephela khona – nokuba kungokujikwa kwendlela yokunyuka komphunga ukuze ujlolise ezantsi okanye ngamanzi angaphezu komhlaba okanye ke kufuduswe/kuthuthwe amanzi kwii-akhwifa (iindawo ezigcina amanzi emvula) ezimelene nendawo leyo iphelelwe ngamanzi.

IBakala leNdalo eliNdululwayo/elixhaswayo (i-REC): Ibakala lendalo elibonisa ibinzana elingqaliweyo lolawulo lwendalo kuloo mjelo uthile wamanzi oko kuxhomekeke kwihlelo lendalo elimisiweyo. La mahlelo asukela kwiBakala A (obonisa ukuba akukho nguqu yenzekileyo okanye isimo somjelo iseseso sendalo) ukuya kwiBakala D (obonisa ukuba isimo somjelo siguquke ngokupheleleyo).

Umjelo: Ngumthamo nekwaliti yamanzi afunekayo ukuze kwaneliswe iimfuno ezisisiseko zoluntu ngokuthi kugcinwe amanzi aya kwaneza loo mfuno isisiseko, kukhuselwe nempilo yasemanzini ngeenjongo zokufezekisa uphuhliso lwendalo oluya kuzinza nokuqinisekisa ukuba kusetyenziswa umjelo wamanzi ofanelekileyo.

Ummandla womlambo (ummandla wendalo): La ngamanqanaba aziziboniso nangumfuziselo wemijelo ekumphezulu wonxweme, okanye loo mimandla ithile apho kuphilisana izinto zasemanzini khona (njengemilambo, imiwonyo, amachweba namanzi angaphantsi komhlaba) apho kuqhuba intlalo yokuphilisana).

Imimandla yobonisele ngamanzi yeminyaka emine (yekota): Ngamacandelwana amahle emimandla yobonisele ngamanzi yeminyaka emine (imimandla yobonisele ngamanzi ekwimisebe yemilambo emikhulu).

4. UDONGA LWEBAKALA LEKWALITI YAMANZI KWIMILAMBO EKWIZIKHUNDLA NEENDAWO EZIKHETHIWEYO ZE-EWR

Umjelo ngamnye uzizintlu ezimbini – uMjelo weeMfuno ezisiSiseko zoLuntu (i-BHN) noMjelo weNdalo (i-ER). I-BHN ibonelela ngeemfuno eziphambili zabantu abathile abancedwa nguloo mjelo uthile futhi yona iquka amanzi okusela, okupheka nalawo okuhlamba. I- ER yona inxulumene namanzi afunelwa imicimbi yokhuselo lwempilo yasemanzini kuloo mjelo uthile. Eli gama umjelo libhekisa **kumthamo nekwaliti** yamanzi akuloo mlambo uthile, futhi wona uguquguquka ngokuxhomekeke kwihlelo lomlambo lowo (umzekelo iHlelo I, II no-III).

UMMANDLA IBREEDE-OVERBERG (UMJELO WENDALO)

UTafle 4.1: Isishwankathelo seenkcukacha zeendawo nezikhundla ze-EWR. Izikhundla ze-EWR zibonise mnyama kakhulu.

Ummandla wobonisele ngamanzi yeminyaka emine	Indawo/isi khundla se-EWR	Umjelo wamanzi	I-PES	I-EIS	i-REC	i-nMAR (MCM)	(i- nMAR)	(%)
G40C	Piii1	Palmiet	B	Phezulu	B	39.9		19.1
G40C	Piv10	Witklippiesskl oof	D	Phezulu	D	15.1		21.5
G40C	Piv9	Palmiet	D	Phezulu	B	78.8		21.5
G40C	Piv8	Klipdrift	D	Phezulu	D	13.6		21.5
G40D	Piv4	Klein-Palmiet	D	Phezulu	D	13.7		21.5
G40D	Piv7	Krom/Ribbok	D	Phezulu kakhulu	A	27.5		21.5
G40D	Piii2	Palmiet	C	Phezulu kakhulu	B/C	206.6		31.2
G40D	Piv12	Dwars/Louws	C	Phezulu kakhulu	C	25.2		100.0
G40D	Piii3	Palmiet	C	Phezulu kakhulu	B	250.4		34.5
G40G	Niii5	Bot	C	Phezulu kakhulu	A	31.9		21.3

Ummandla woboniselw ngamanzi weminyaka emine	Indawo/isi khundla se-EWR	Umjelo wamanzi	I-PES	I-EIS	i-REC	i-nMAR MCM)	(i- i-EWR nMAR)	(%
G40H	Nx6	Onrus	E	Phezulu	B	5.1	13.4	
G40F	Niv43	Swart	E	Phezulu	B	42.1	13.3	
G40K	Niv45	Steenbok	E	Phezulu kakhulu	A	10.8	12.2	
G40J	Nii4	Hartebees	D	Phezulu kakhulu	B	18.4	12.5	
G40K	Nv23	Klein	D	Phakathi	C	43.0	19.3	
G40M	Nx8	Uilkraal	C	Phezulu kakhulu	A	2.4	19.2	
G50B	Ni4	Nuwejaar	D	Phakathi	D	12.5	13.0	
G50C	Nvii15	Heuninges	D	-	D	17.8	13.1	
G50C	Niv44	Heuninges	D	Phezulu kakhulu	B	18.8	13.1	
G50C	Nii5	Kars	E	Phezulu kakhulu	B	21.6	20.4	
G50E	Nv24	Kars	C	Phakathi	B	15.4	30.3	
G50H	Nii7	DeHoop Vlei	B	Phezulu	B	27.1	30.0	
G50H	Nii6	Sout	D	-	B	4.2	12.6	
H10B	Nvii3	Rooikloof	B	Phezulu	B	6.807	37.95	
H10B	Niv3	Titus	C	Phakathi	C	26.2	22.0	
H10C	Niv1	Koekedou	D	Phezulu kakhulu	A	18.8	14.2	
H10C	Niv2	Dwars	C	Phezulu	B	74.9	22.0	
H10D	Nvi4	Breede	D	Phakathi	D	175.509	17.51	
H10D	Niv4	Witels	A	Phezulu kakhulu	A	84.3	43.3	
H10D	Nvi3	Breede	C	Phezulu	B	252.8	31.7	
H10E	Nvi2	Wit	A	Phezulu kakhulu	A	42.6	46.6	
H10F	Niv6	Wabooms	D	Phezulu	B	7.4	14.4	
H10F	Nviii1/ EWR1	Breede	D/E	Phezulu	D	434.90	31.7	
H10G	Niv7	Slanghoek	D	Phezulu	B	32.6	14.5	
H10G	Niii1	Breede	D	Phezulu	B	497.6	25.4	
H10J	Niv40	Elands	B	Phezulu kakhulu	A	58.1	50.8	
H10J	Niv41	Krom	B	Phezulu kakhulu	A	9.0	50.8	
H10J	Nvii2/ EWR2	Molenaars	C	Phezulu kakhulu	B	105.6	35.0	
H10J	Niv42	Molenaars (Smalblaar)	E	Phezulu	B	191.2	17.4	
H10K	Niv12	Holsloot	C	Phezulu	B	119.5	35.0	
H10H	Nvii6	Hartbees	D	Phezulu kakhulu	A	4.0	14.4	
H10H	Niv9	Hartbees/ de Wetskloof	D	Phezulu kakhulu	A	10.2	14.4	
H10L	Nv3	Breede	C	Phezulu	B	850.9	31.7	
H20G	Nvii7	Hex	C	Phakathi	C	102.8	22.3	
H20H	Niv10	Hex	D	Phezulu	B	107.1	22.3	
H40B	Nvii5	Koo	D	Phezulu	B	0.9	13.1	
H40C	Niv11	Nuy	E	Phezulu	B	29.4	13.2	
H40D	Niv13	Doring	E	Phezulu	B	47.4	12.9	
H40F	Nvii8/ EWR3	Breede	C/D	Phakathi	C/D	1042.8	45.5	
H40G	Nvii11	Poesnells	D	Phezulu	B	16.1	12.8	

Ummandla woboniselongamanzi weminyaka emine	Indawo/isi khundla se-EWR	Umjelo wamanzi	I-PES	I-EIS	i-REC	i-nMAR (MCM)	(i- i-EWR nMAR)	(%)
H40K	Niv14	Keisers	D	Phezulu kakhulu	A	12.6		12.5
H40H	Niv15	Vink	D	Phezulu kakhulu	A	15.6		12.4
H30C	Niv20	Pietersfontein	D	Phakathi	C	17.3		12.0
H30B	Niv18	Kingna	D	Phezulu	B	27.1		12.3
H30D	Nvii9	Keisie	D	Phezulu	B	21.5		11.9
H30E	Nii2	Kogmanskloof	D	Phezulu kakhulu	B	52.0		18.9
H50B	Ni2	Breede	D	Phezulu	B	1170.1		17.3
H60B	Nvii10	Du Toits	B	Phezulu kakhulu	A	43.9		50.8
H60D	Nv7	Riviersonderend	C	Phezulu kakhulu	A	370.2		30.1
H60E	Niv28/ EWR6	Baviaans	B	Phezulu	B	7.9		70.90
H60E	Niv29	Sersants	D	Phezulu	B	4.6		29.9
H60F	Niv30	Gobos	C	Phezulu kakhulu	A	12.4		48.1
H60F	Nv9/EWR5	Riviersonderend	D	Phezulu	D	413.7		24.5
H60G	Niv31	Kwartel	D	Phezulu	B	10.7		13.4
H60H	Niv33	Soetmelksvlei	D	Phezulu kakhulu	A	4.0		29.9
H60H	Niv34	Slang	D	Phezulu kakhulu	A	2.1		29.9
H60H	Nv10	Riviersonderend	D	Phezulu kakhulu	A	442.9		24.5
H60K	Niv35	Kwassadie	E	Phezulu kakhulu	A	5.9		17.3
H60L	Ni3	Riviersonderend	D	Phezulu	B	483.8		24.5
H70A	Niv24	Leeu	E	Phezulu kakhulu	A	5.8		12.6
H70B	Nv2	Breede	C	Phezulu	B	1701.4		26.4
H70D	Nii3	Tradouw	B	Phezulu kakhulu	A	19.4		29.9
H70F	Niv25	Buffeljags	E	Phezulu	B	119.4		14.1
H70G	Niii4/ EWR4	Breede	C	Phezulu kakhulu	B/C	1832.7		40.1
H70J	Niv26	Slang	E	Phezulu	B	10.0		14.2

UMMANDLA WONXWEME IGOURITZ (UMJELO WENDALO)

UTafle 4.2: Isishwankathelo seenkcukacha seendawo nezikhundla ze-EWR sites. Izikhundla zeEWR ziboniswe mnyama kakhulu.

Umandla woboniso ngamanzi weminyaka emine	Indawo/isikhundla se-EWR	Umjelo wamanzi	I-PES	I-EIS	i-REC	i-nMAR (i-MCM)	i-EWR (% nMAR)
J11C	giv34	Buffels	B	Phezulu	B	13.1	26.5
J11F	gv25	Buffels	B	Phezulu	B	24.2	17.8
J11H	J1BUFF-EWR5	Buffels	C	Phakathi	C	27.4	17.9
J11K	giv32	Groot	D	Phezulu	B	30.5	17.9
J12D	giv28	Touws	D	Phezulu	B	16.4	11.3
J12H	giv27	Touws	B	Phakathi	C	26.4	26.8
J12K	giv26	Brak	C	Phezulu	B	2.9	17.7
J12L	J1DORI-EWR7	Doring	C/D	Phantsi	C/D	2.9	12.0
J12L	J12L Modelled	Huis	D		D	1.56	40.3
J12M	J1TOUW-EWR3	Touws	B/C	High	B/C	33.5	17.8
J13B	gv7	Groot	C	Phezulu	B	72.7	18.0
J13C	gii3	Groot	B	Phezulu	B	78.1	27.0
J21D	giv3	Gamka	B	Phezulu	A	31.9	27.1
J22F	giv1	Koekemoers	C	Phezulu kakhulu	A	7.4	17.9
J22K	giv2	Leeu	C	Phezulu kakhulu	A	17.1	17.9
J23F	gv17	Gamka	B	Phezulu	B	58.1	27.0
J23J	gv27	Gamka	C	Phezulu	B	69.6	18.3
J24E	gv14	Dwyka	A	Phezulu	B	4.0	39.1
J25A	J2GAMK-EWR4	Gamka	C/D	Phezulu	C	79.8	14.9
J25E	gii2	Gamka	C	Phezulu	B	111.8	15.2
J31D	J3OLIF-EWR9	Olifants	C	Phakathi	C	11.8	17.8
J32E	giv15	Traka	C	Phezulu	C	2.7	17.9
J33B	gv33	Olifants	D	Phezulu	B	25.0	11.9
J33E	gv21	Meirings	C	Phezulu kakhulu	A	21.4	19.1
J33F	giv11	Olifants	E	Phezulu	B	80.0	12.4
J34C	J3KAMM-EWR10	Kammanasie	C/D	Phantsi	C/D	41.2	15.3
J34F	giv10	Leeu	E	Phezulu kakhulu	A	59.2	12.1
J35E	gv19	Olifants	E	Phezulu	B	224.5	12.9
J35F	giv17	Olifants	D	Phezulu	B	253.4	12.9
J40B	J4GOUR-EWR6	Gouritz	C	Phakathi	C	489.1	14.8
J40E	gv9	Gouritz	C	Phezulu	B	571.8	14.8
H80B	giii5	Duiwenhoks	E	Phezulu kakhulu	A	62.5	20.1
H80E	H8DUIW-EWR1	Duiwenhoks	D	Phantsi	D	83.2	20.9
H90C	giv27	Korinte	D	Phezulu	B	34.1	14.5
H90C	H9GOUK-EWR2	Goukou	C/D	Phakathi	C/D	50.9	24.2
H90E	gv41	Goukou	C	Phezulu	B	105.0	28.2

Ummandla woboniselomngamanzi weminyakamemine	Indawo/isikhundla se-EWR	Umjelo wamanzi	I-PES	I-EIS	I-REC	i-nMAR (i-MCM)	i-EWR (% nMAR)
K10D	giv25	Brandwag	D	Phezulu	B	17.9	9.9
K20A	gvii7	Groot-Brak	B/C	Phezulu kakhulu	A	27.0	26.5
K20A	gviii2- EWR GB1-BC	Groot-Brak	B/C	Phezulu kakhulu	B/C	15.3	26.5
K20A	gviii3-EWR Var 3	Varing	D	Phezulu	C/D	8.4	20.9
K20A	gviii12- EWR Var2	Varing	D	Phezulu	C/D	6.0	20.9
K30A	gviii4-EWR	Maalgate	B	Phezulu kakhulu	A	15.3	46.0
K30A	gvii8	Maalgate	B	Phezulu	D	30.1	16.4
K30B	gvii9	Malgas	C	Phezulu kakhulu	C	17.3	31.6
K30B	gviii6 EWR Gwa1 -D	Gwaing	E	Phezulu	D	34.1	16.4
K30C	gviii7 EWR Sw1 - D	Swart	D	Phezulu	D	16.1	14.5
K30C	gvii11 EWR Ka1 - D	Kaaimans	B	Phezulu	B	18.6	50.2
K30C	gviii8 EWR Si1 -B	Silver	B	Phezulu kakhulu	B	14.9	50.2
K30D	gvii12	Touws	B	Phezulu kakhulu	A	16.7	30.3
K30D	gx8	Klein Keurbooms	D	Phezulu kakhulu	B	2.5	14.1
K40A	giii10 EWR 2 Diep -B	Diep	B	Phezulu kakhulu	B	12.4	30.3
K40B	giii13	Hoekraal	B	Phezulu	A	27.9	30.3
K40C	gviii13 EWR 4 Karatara-AB	Karatara	B	Phezulu kakhulu	A/B	11.2	40.2
K40C	giii11	Karatara	A/B	Phezulu kakhulu	A	33.8	40,2
K40E	Gou 1	Goukamma	B/C	Phezulu kakhulu	B/C	30.4	38.5
K50A	EWR 1	Knysna	B	Phezulu	B	26.5	32.1
K50A	Kny 2	Knysna	B	-	B	46.5	32.1
K50B	EWR 2	Gouna	A/B	Phezulu kakhulu	A/B	27.6	53.4
K60C	K6KEUR-EWR8	Keurbooms	C	Phezulu kakhulu	B/C	46.1	34.9
K60D	giv5	Palmiet	A	Phezulu kakhulu	A	42.1	48.3
K60F	giv4	Bitou	C	Phezulu kakhulu	A	23.6	22.8
K60G	Noe 1	Noetsie	B	Phezulu kakhulu	A/B	4.8	63.4
K60G	gx3	Piesang	D	Phezulu kakhulu	A	7.3	28.5
K60E	gx9	Keurbooms	C	Phezulu kakhulu	A	91.3	34.9
K70A	gx4	Buffels	B	Phezulu kakhulu	B	1.8	34.3
K70A	gx5	Sout	B	Phezulu kakhulu	B	3.8	34.3
K70B	gvii15	Bloukrans	B	Phezulu kakhulu	B	31.2	33.9

UTafale 4.3: IIMfuno ezisiSiseko zoLuntu kwiBreede-Gouritz WMA

Ummandla woboniselo ngamanzi weminyaka emine	Umjelo wamanzi	i-BHN (%NMAR)	Ummandla woboniselo ngamanzi weminyaka emine	Umjelo wamanzi	i-BHN (%NMAR)
G40C	Palmiet	0.008	J11C	Buffels	0.02
G40D	Palmiet	0	J11F	Buffels	0.03
G40G	Bot	0.50	J11H	Buffels	0.03
G40H	Onrus	5.88	J11K	Groot	0.36
G40F	Swart	0.17	J12D	Touws	0.03
G40K	Steenbok	0	J12H	Touws	0
G40J	Hartebees	0.08	J12K	Brak	0
G40K	Klein	0	J12L	Doring	0.21
G40M	Uilkraal	0.125	J12M	Touws	0
G50B	Nuwejaar	1.12	J13B	Groot	0.01
G50C	Heuninges	0	J13C	Groot	0
G50E	Kars	0.84	J21D	Gamka	0
G50G	DeHoop Vlei	0.03	J22F	Koekemoers	0.14
G50H	Sout	0.04	J22K	Leeu	0
H10B	Rookloof	0	J23F	Gamka	0.10
H10C	Dwars	0.70	J23J	Gamka	0
H10D	Breede	0	J24E	Dwyka	0
H10E	Wit	0	J25A	Gamka	0.003
H10F	Breede	0.04	J25E	Gamka	0.03
H10G	Slanghoek	0	J31D	Olifants	0.02
H10J	Elands	0.02	J32E	Traka	0.03
H10K	Holsloot	0	J33B	Olifants	0.02
H10H	Breede	0	J33E	Meirings	0.56
H10L	Breede	0.005	J34C	Olifants	0.01
H20G	Hex	0.01	J34F	Kammanassie	0.05
H20H	Hex	1.20	J35E	Leeu	0.004
H40B	Koo	2.22	J35F	Olifants	0.01
H40C	Nuy	0.07	J40B	Olifants	0
H40D	Doring	0	J40E	Gouritz	0.003
H40F	Breede	0.002	H80B	Gouritz	0
H40G	Poesnels	0	H80E	Duiwenhoks	0.005
H40K	Keisers	0.32	H90C	Goukou	0.33
H40H	Vink	0.06	H90E	Goukou	0.06
H30C	Pietersfontein	0.07	K10D	Brandwag	0.06
H30B	Kingna	1.33	K20A	Varing	1.50
H30D	Keisie	0.04	K30A	Maalgate	0.39
H30E	Kogmanskloof	0.17	K30B	Malgas	0.18
H50B	Breede	0.004	K30C	Swart	8.99
H60B	Du Toits	0	K30D	Touws	0.54
H60D	Riviersonderend	0.001	K40A	Diep	0.04
H60E	Baviaans	1.14	K40B	Hoekraal	0
H60F	Gobos	0.007	K40C	Karatara	0.54
H60G	Kwartel	0	K40E	Goukamma	0.39
H60H	Soetmelksvlei	0	K50A	Krystna	0.002
H60K	Kwassadie	0	K50B	Gouna	0.47
H60L	Riviersonderend	0	K60C	Keurbooms	0.03
H70A	Leeu	0	K60D	Palmiet	25.26
H70B	Breede	0.01	K60F	Bitou	0.38
H70D	Tradouw	0	K60G	Piesang	4.34
H70F	Buffeljags	0.02	K60E	Keurbooms	0.02
H70G	Breede	0	K70A	Buffels	1.32
H70J	Slang	0.40	K70B	Bloukrans	0.03

5. IBAKALA LEKWALITI YAMANZI ANGAPHEZU KOMHLABA KWIMILAMBO EKWIZIKHUNDLA ZE-EWR

I-BREDE-OVERBERG

UTafle 5.1: Amabakala e-PES novavanyo lulonke lwesikhundla kumlambo iBreede kuNode Nviii1 (omelwe siSikhundla 1 seEWR)

UMLAMBO	Umlambo iBreede	AMANQANABA OHLOLO LWEKWALITI YAMANZI		
I-WQRU	1(Ekudibaneni komphezulu womlambo iBreede nomlambo iWit)	I-RC	Isitishi sohlolo sikaDWA @ Koekedou kuMlambo ose-Ceres (H1H013Q01) (1998 -2002, n=38)	
ISIKHUNDLA SE-EWR	Isikhundla 1 se-EWR	I-PES	Isitishi sohlolo sikaDWA @ Witbrug (H1H006Q01) (1998 -2002, n=143)	
Uvavanyo lokuba nakho		Phakathi. Isikhundla seEWR sithe qelele ukuya phaya kumazantsi onxweme lwesitishi sohlolo sikaDWA.		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES Value	IBakala/Amagqabantshintshi
Iityuwa ezingezozendalo (mg/L) 1:95 th Amaxabiso ePercentile	MgSO ₄	7	21	IBakala B
	Na ₂ SO ₄	8	4	IBakala A
	MgCl ₂	5	6	IBakala A
	CaCl ₂	12	24	IBakala B
	NaCl	6	27	IBakala A
	CaSO ₄	0	0	IBakala A
Izondlo (mg/L)	SRP	-	-	Azikho iinkcukacha
	TIN	2.8011	0.318	IBakala B
	TDS	21.8-50.6	< 45	IBakala B. I-TDS iyonyuka ngeenyanga zaseBusika
	TSS	-	< 5	Yamkelekile
Ukwahluka kwendalo	PO ₄	0.024	0.042	IBakala C
	pH (5 th – 95 th %)	6.2-7.5	6.9 – 7.8	IBakala C
	Ubushushu	-	-	Azikho iinkcukacha kodwa akukho ngxaki iboniweyo ngemixube ye-DO
	I-oksijini enyibilikisiweyo	-	-	
	Ubukho bodaka (NTU)	-	2	Azikho iinkcukacha
Ukutsala umbane (mS/m)	-	-	Azikho iinkcukacha	
Ukwahluka ngokwemvakalelo	Chl a: periphyton	-	-	Azikho iinkcukacha
	Chl a: phytoplankton	-	-	Azikho iinkcukacha
	Ubukhulu bezinto eziphilayo ezingenamathambo	SASS score = >110 and an ASPT score >7	SASS score = 69 and ASPT score = 5.3	IBakala D/E. Uguquke kakhulu. Kude kwaphela nommandla wokuphilisana ngenxa yokuzaliswa kwamanzi.
	Inqaku lemigqeku yeentlanzi			U-D/E. kwaziswa ngemigqeku yeentlanzi, oko kukuthi i-bass, i-trout ne-blue gills.
Iityhefu		-	Azikho iinkcukacha kodwa iintsalela zezibulali zinambuzane zithande ukuba yingxaki ngenxa yemisebenzi yezolimo eCeres	
LULONKE UHLELO LWESIKHUNDLA		IBakala B		

UTafle 5.2: Amabakala e-PES novavanyo lwesikhundla lunonke kumlambo iMolenaars kwiNode Nvii2 (emelwe siSikhundla 2 seEWR)

UMLAMBO		AMANQANABA OHLOLO LWEKWALITI YAMANZI		
I-WQRU	2 (Umlambo opheleleyo iMolenaars)	I-RC	Umlambo iMolenaars @ kumjelo wehlathi iHawequas (H1H018Q01) (1998 -1992, n=93)	
ISIKHUNDLA SE-EWR	Isikhundla 2 se-EWR	I-PES	Umlambo iMolenaars @ kumjelo wehlathi iHawequas (H1H018Q01) (1998 -2002, n=141)	
Uhlolo lokubanakho		Phezulu. Indawo yohlolo ikufuphi kakhulu kwisikhundla se-EWR. Iinkcukacha zembali azibonisi simbo sithile, yiloo nto iinkcukacha ze-PES ziye zasetyenziselwa ukubonisa iimeko ekubhekiswa kuzo.		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	IBakala/Amagqabantshintshi
lityuwa ezingezozendalo (mg/L) 1:95 th Amaxabiso %	MgSO ₄	6	6	IBakala A
	Na ₂ SO ₄	9	9	IBakala A
	MgCl ₂	4	4	IBakala A
	CaCl ₂	6	6	IBakala A
	NaCl	9	9	IBakala A
	CaSO ₄	0	0	IBakala A
Izondlo (mg/L)	SRP		-	
	TIN (1:50 th %)	0.151	0.151	IBakala A
	TDS	22	22	IBakala A
	TSS	< 5	< 5	IBakala A
	PO ₄ (1:50 th %)	0.025	0.025	IBakala C
Ukwahluka kwendalo	NH ₃ -N	<0.015.	<0.015.	Kobo buthuba besimo esaziwayo
	pH (5 th – 95 th %)	5.5 – 7.3	5.5 – 7.3	<i>IBakala A.</i> Imeko eqhelekileyo yobumuncu bamanzi omlambo waseNtshona Koloni
	Ubushushu			
	I-oksijini enyibilikisiweyo	80 - 120% uhluzo	80 - 120% uhluzo	Kobo buthuba besimo esaziwayo, iBakala A
	Ubukho bodaka (NTU)	-	0	
Ukwahluka ngokwemvakalelo	Ukutsala umbane (mS/m)	-	-	Azikho iinkcukacha
	Chl a: periphyton	-	-	Azikho iinkcukacha
	Chl a: phytoplankton	-	-	Azikho iinkcukacha
	Ubukhulu bezinto eziphilayo ezingenamathambo	Inqaku le-SASS ≥140 nenqaku le-ASPT >8	Inqaku le-SASS score = 175 nenqaku le-ASPT = 7.9	<i>IBakala A/B.</i> Catchment is relatively pristine
lityhefu	Inqaku lemigqeku yeentlanzi			<i>IBakala E.</i> Azikho iintlanzi zomthonyama eziye zabhaliswa ngethuba lophando
		-	-	Akukho zingxaki zeetyhefu ziye zavela
LULONKE UHLELO LWESIKHUNDLA		IBakala A		

UTafale 5.3: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iBreede kuNode Nvii8 (emelwe siSikhundla 3 se-EWR)

UMLAMBO	Umlambo iBreede	IINDAWO ZOHLULO LWEKWALITI YAMANZI		
I-WQRU	3 (kuMbindi womlambo iBreede Ukusuka ekudibaneni kweMolenaars ne Kogmanskloof	I-RC	Ayikho. Asikho isitishi sohlulo lwe-WQ esiyi sanika iinkcukacha ngeli candelo loMjelo.	
ISIKHUNDLA SE-EWR	ISikhundla 3 se-EWR Site 3	I-PES	Umlambo iBreede e-Le Chasseur (H4H017Q01) (1995 -1999, n=214)	
Uvavanyo lokuba nakho		Phakathi		
Izithako zekwaliti yamanzi		Ixabiso I-RC	Ixabiso le-PES Value	IBakala/amagqabantsshintshi
lityuwa ezingezozendalo (mg/L) (1:95 th %)	MgSO ₄	-	39	Category D
	Na ₂ SO ₄	-	5	Category A
	MgCl ₂	-	12	Category A
	CaCl ₂	-	32	Category B
	NaCl	-	95	Category B
	CaSO ₄	-	0	Category A
Izondlo (mg/L)	SRP	-		
	TIN (1:50 th %)	-	0.242	Category A
	TDS	-	< 45	B Category. High TDS loads. Irrigation return flows in tributaries and main stem between Brandvlei Dam and EWR Site 3.
	TSS	-	< 5	High TSS loads. Releases from Brandvlei to alleviate high TDS.
Ukwahluka ngendalo	PO ₄ (1:50 th %)	-	0.032	C Category
	pH (5 th – 95 th %)	-	6.9 – 7.8	A Category
	Temperature (°C)	-	22.4	Summer temperature
	Dissolved oxygen	-	-	No observed data
	Turbidity (NTU)	-	-	No data. Low sediment production area
	Electrical conductivity (mS/m)	-	-	No data
Ukwahluka ngemvakalelo	Chl a: periphyton	-	-	No data
	Chl a: phytoplankton	-	-	No data
	Ubukhulu bezinto eziphilayo ezingenamathambo	Inqaku le-SASS ≥110 nengaku le-ASPT >7	Inqaku le-SASS = 91 nengaku le-ASPT = 7	IBakala A. lithetha ukuthi akukho futhe lingako kwisakhiwo sasekuhlaleni.
	Inqaku lemigqeku yeentlanzi	-		IBakala D. mi-2 kuphela kwesi-7 imigqeku yeentlanzi zamanzi ahlaziyekileyo ekulindeleke ukuba zibekho apho kuthathwa khona iisampulu.
lityhefu		-	-	Azikho iinkcukacha kodwa ke izibulala zinambuzane zixakile ngenxa yefuthe lemisebenzi yezolimo
LULONKE UHLELO LWESIKHUNDLA		IBakala D. ikakhulu ifuthe libakho ngenxa yeemvula ezingephi zasehlotyeni neengqumba zeentlunge ezandileyo.		

UTafale 5.4: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iBreede kuNode Nvii4 (emelwe siSikhundla 4 se-EWR)

MLAMBO	Umlambo iBreede	IINDAWO ZOHLOLO LWEKWALITI YAMANZI		
I-WQRU	6 (umlambo iBreede ukusuka kumlambo i- Buffelsjags ukuya kutsho eChwebeni)	I-RC	Azikho. Asikho isitishi sohlolo lwe-WQ esiye sanika iinkcukacha ngeli candelo loMjelo.	
ISIKHUNDLA SE-EWR	Isikhundla 4 se-EWR	I-PES	Kumazantsi omlambo iBreede @ eSwellendam (H7H006Q01) (1995 -1999, n=214)	
Uhlolo lokuba nakho		Phakathi (Akukhange kubekho kubekho zinkcukacha zezamanzi kulo mmandla womlambo, futhi amanzi ebemaninzi kakhulu ukuba inani losetyenziso lingathatyathwa; indawo yohlolo ibikude lee ukuya kumphezulu womjelo.		
Izithako zekwaliti yamanzi		Ixabiso le-RC	Ixabiso le-PES	Ibakala/Amagqabantshintshi
lityuwa ezingezizo zendalo (mg/L) (1:95 th %)	MgSO ₄	-	83	IBakala E/F
	Na ₂ SO ₄	-	3	IBakala A
	MgCl ₂	-	49	IBakala D
	CaCl ₂	-	62	IBakala C
	NaCl	-	318	IBakala D
	CaSO ₄	-	0	IBakala A
Izondlo (mg/L)	SRP	-	-	Azikho iinkcukacha
	TIN (1:50 th %)	-	0.23	IBakala A
	TDS	-	-	IBakala C. iingqumba zeTDS ziphezulu ngenxa yamanzi abuya kwezonkcnkeshelo kumasebe emilambo nakuleyo mikhulu
	TSS	-	-	Azikho iinkcukacha
	P0 ₄ (1:50 th %)	-	0.024	IBakala C
	NH ₃ -N	-	0.3 (Phakathi)	Azikho iinkcukacha zembali
Ukwahluka kwendalo	NO ₂ NO ₃ -N	-	<1	Kundululwa ubuninzi bazo ebusika kodwa ehlotyeni zingakumaqondo aphantsi kangange- < 0.3 mg/l
	pH (5 th – 95 th %)	-	6.9 – 8.1	iBakala A/B. Akukho nguqu ingako.
	Ubushushu (°C)	-	-	Akukho zinkcukacha ziboniweyo. Kubonwe ingxaki yokwenyuka kusehla kwamanqanaba e- DO.
	I-oksijini enyibilikisiweyo	-	-	
	Ubukho bodaka (NTU)	-	-	
Ukutsala umbane (mS/m)	-	-	Azikho iinkcukacha	
Ukwahluka ngemvakalelo	Chl a: periphyton	-	-	Azikho iinkcukacha
	Chl a: phytoplankton	-	-	Azikho iinkcukacha
	Ubukhulu bezinto eziphilayo ezingenamathambo	Inqaku le-SASS ≥110 nenqaku le-ASPT >7	Inqaku le-SASS = 87 nenqaku le-ASPT = 6.9	IBakala B. <i>chaphazeleke ngokuphakathi</i>
	Inqaku lemigqeku yeentlanzi	10 imigqeku yeentlanzi zomthonyama		IBakala C. <i>ubukho be-carp ne-bass.</i>
lityhefu		-		Azikho iinkcukacha kodwa ke izibulala zinambuzane zixakile ngenxa yefuthe lemisebenzi yezolimo
LULONKE UHLELO LWESIKHUNDLA		IBakala D. <i>oko kuxhomekeka ekungeneni kwamanzi anekwaliti asuka kumlambo iBuffelsjags khonukuze kuqinisekiswa ukuba i-WQ kumazantsi ecandelo lomlambo nasechwebeni yamkelekile</i>		

UTafle 5.5: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iRiviersonderend kuNode Nv7 (emelwe siSikhundla 5 se-EWR)

UMLAMBO	Umlambo iRiviersonderend	IINDAWO ZOHOLO LWEKWALITI YAMANZI		
I-WQRU	7 (kumbindi womlambo iRiviersonderend ukusuka kwidama i-Theewaterskloof ukuya kumlambo i-Bok)	RC	iRiviersonderend kuMjelo weHlathi okumlambo iSwart River/Nuweberg (H6H008Q01) (1990 -1992, n=34)	
ISIKHUNDLA SE-EWR	Isikhundla se-EWR 5	PES	iRiviersonderend kwidama iTheewaterskloof (H6H012Q01) (1998 -2002, n=39) iRiviersonderend eReenen (H6H009Q01) (1995 -1999, n=56)	
Uhlolo lokuba nakho		Phezulu		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	Ibakala/amagqabantsintshi
lityuwa ezingezozendalo (mg/L) 1:95 th % ixabiso	MgSO ₄	7	12	IBakala A
	Na ₂ SO ₄	7	13	IBakala A
	MgCl ₂	4	14	IBakala A
	CaCl ₂	5	14	IBakala A
	NaCl	11	99	IBakala B
	CaSO ₄	0	0	IBakala A
Izondlo (mg/L)	SRP	-	-	Azikho iinkcukacha
	TIN (1:50 th %)	0.068	0.154	IBakala A.
	TDS	13.6-32	Ehlotyeni: <100 Ebusika : < 150	IBakala B. Iingqumba eziphezulu zeTDS ngenxa yamanzi asuka kwezolimo ebusika
	TSS	< 5	-	Azikho iinkcukacha
	P0 ₄ (1:50 th %)	<0.0165	0.013	IBakala B. Phezulwana.
Ukwahluka kwendalo	NO ₂ NO ₃ -N	0.02	<0.25	Kundululwa ubuninzi bazo ebusika kodwa ehlotyeni zingakumaqondo aphantsi kangange < 0.2 mg/l
	pH (5 th – 95 th %)	6.5 – 7.5	6.4 - 7.4	IBakala A/B.
	Temperature (°C)	+4 °C	-	
	I-oksijini enyibilikisiweyo	80 - 120% Uhluzo	-	Azikho iinkcukacha
	Ubukho bodaka (NTU)	-	4	
Ukutsala umbane (mS/m)	-	-	Azikho iinkcukacha	
Ukwahluka ngokwemvakalelo	Chl a: periphyton	-	-	Azikho iinkcukacha
	Chl a: phytoplankton	-	-	Azikho iinkcukacha
	Ubukhulu bezinto eziphilayo ezingenamathambo	Inqaku le-SASS ≥110 nenqaku le-ASPT >7	Inqaku le-SASS = 95 nenqaku le-ASPT = 6.8	IBakala C/D. Iindawo ezininzi zinekwali entle yamanzi kodwa ke ikwali yendawo yokuphilisana yona imbi kakhulu
	Inqaku lemigqeku yeentlanzi	Imigqeku emithandathu yeentlanzi		IBakala E. Zigqunyelelwa zii-bass ezinimilomo emikhulu nemincinci. Linciphile inani lemigqeku yeentlanzi zomthonyama.
lityhefu		-	-	Azikho iinkcukacha kodwa iintsalela zezibulala zinambuzane ziyinxaki ngenxa yezolimo kumphezulu womjelo wesikhundla i- EWR

LULONKE UHLELO LWESIKHUNDLA	IBAKALA A/B (Ubukho bedama iTheewaterskloof kuthande ukuyinciphisa ikwaliti yamanzi alaa mlambo usemazantsi onxweme)
------------------------------------	---

UTafle 5.6: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iBaviaans kuNode Niv28 (emelwe siSikhundla 6 se-EWR)

UMLAMBO		Baviaans River	WATER QUALITY MONITORING POINTS	
I-WQRU	9 (umlambo iBaviaans uphela)	I-RC	Umlambo iBaviaans kwisitishi somsebenzi saseGenadendal (H6H005Q01) (1972 -1994, n=346)	
ISIKHUNDLA SE-EWR	Isikhundla 6 se-EWR Site 6	I-PES	Umlambo iBaviaans kwisitishi somsebenzi saseGenadendal (H6H005Q01) (1998 -2002, n=42)	
Uvavanyo lokubanakho		Phezulu. Iinkcukacha ezingenileyo zintle futhi ziya kunceda kuvavanyo lobhekiselelo nolwe-PES. Iinkcukacha zembali azibonisi zimbo zithile, yiloo nto iinkcukhacha ze PES ziye zasetye data was used for Reference conditions		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	IBakala/Amagqabantshintshi
lityuwa ezingezozendalo (mg/L) 1:95 th % Ixabiso	MgSO ₄	9	9	IBakala A
	Na ₂ SO ₄	9	9	IBakala A
	MgCl ₂	4	4	IBakala A
	CaCl ₂	10	10	IBakala A
	NaCl	25	25	IBakala A
	CaSO ₄	0	0	IBakala A
Izondlo (mg/L)	SRP	-	-	Azikho iinkcukacha
	TIN (1:50 th % value)	0.04	0.040	IBakala A
	TDS	41	41	IBakala A
	TSS	-	-	Azikho iinkcukacha
	PO ₄ (1:50 th % value)	0.022	0.022	IBakala A
Ukwahluka kwendalo	pH (5 th – 95 th %)	4.7 – 7.1	4.7 – 7.0	IBakala A. Lithathwa njengendalo kwimlambo ene pH esezantsi yaseNtshona Koloni
	Ubushushu (°C)	-	-	
	I-oksijini enyibilikisiweyo	-	-	Azikho iinkcukacha kodwa ke akukho ngxaki yeDO evelayo
	Ubukho bodaka (NTU)	-	-	
	Ukutsala umbane (mS/m)	-	-	Azikho iinkcukacha
Ukwahluka ngokwemvakalelo	Chl a: periphyton	-	-	Azikho iinkcukacha
	Chl a: phytoplankton	-	-	Azikho iinkcukacha
	Ubukhulu bezinto eziphilayo ezingenamathambo	Inqaku le-SASS ≥140 nenqaku leASPT >8	Inqaku le-SASS = 109 nenqaku le-ASPT = 8.38	IBakala A. indawo yokuphilisana inciphile.
	Inqaku lemigqeku yeentlanzi	Imigqeku emithathu yeentlanzi zomthonyama		IBakala A/B. Yonke imigqeku ekulindeleke ukuba iphile apha ibhalisiwe
lityhefu		-	-	Azikho iinkcukacha kodwa ke kodwa ke ayikho ingxaki yeetyhefu ebonakalayo
LULONKE UHLELO LWESIKHUNDLA		IBakala		

IGOURITZ

UTafale 5.7: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iDuiwenhoks kwiSikhundla 1 se-H8DUIW-EWR

UMLAMBO		Umlambo iDuiwenhoks	IINDAWO ZOHOLO LWEKWALITI YAMANZI	
I-WQRU			I-RC	Udonga olunqamlezileyo lokuthatha umlinganiselo lukaDWS H8H001Q01 (1967 – 1979; inani leesampulu (n) = 66 - 71, Ukutsala umbane : n = 110).
ISIKHUNDLA SE-EWR		I-H8DUIW-EWR1	I-PES	Udonga olunqamlezileyo lokuthatha umlinganiselo lukaDWS H8H001Q01 (2007 – 2013; n = 69, Fluorine (F) = 48).
Uvavanyo lwesikhundla		Ukuba nakho: 3.5		
IZithako zeKwaliti yaManzi		Ixabiso le-PES	IBakala/Amagqabantsintshi	
lityuwa ezingezozendalo (mg/L)	SO ₄	N/A	-	
	Na	382.2	Zibetha ngaphaya kwe ≤ 70 mg/L (TWQR) zineda kwezolimo : Kunkcenkeshelo	
	Mg	67.4	Asikho isikhokelo.	
	Ca	55.0	Asikho isikhokelo.	
	Cl	805.4	Zibetha ngaphaya kwe ≤ 100 mg/L (TWQR) zineda kwezolimo : Kunkcenkeshelo	
	K	9.25	Asikho isikhokelo	
Izondlo (mg/L)	SRP	0.014	A	
	TIN	0.118	A	
Ukwahluka kwendalo	pH (5 th – 95 th %)	6.6 and 8.1	B	
	Ubushushu (°C)	N/A	A/B. Ifuthe libonakala xa amanzi emancinci.	
	I-oksijini enyibilikisiweyo	N/A	B. Ifuthe libonakala xa amanzi emancinci.	
	Ubukho bodaka (NTU)	N/A	B. Inguqu ebonakala eludakeni ibonakala ibangelwa ikakhulu ziinguqu zendalo ukuya kwezo zenziwe ngabantu, umzekelo ukukhiwa kwegraveli phaya kumphezulu womjelo	
	Ukutsala umbane (mS/m)	272	80 mS/m	
Ukwahluka ngokwemvakalelo	Chl a: phytoplankton	N/A	N/A	
	Izinto eziphilayo ezingenamathambo	50.7% Inqaku le-SASS = 78 Nenqaku le-ASPT = 56	D	
	ii-Diatoms	11.1	C/D (n = 1, Jan 2014)	
	Inqaku leyeentlanzi	51.6%	D (ezifuduselwe emachwebeni kummandla wamanzi ahlaziyekileyo nowezo zingezizo ezomthonyama).	
lityhefu	i-Ammonia (njengo-N)	0.003	A	
	i-Fluoride (njengo-F)	0.33	A	
LULONKE UHLELO LWESIKHUNDLA		IBakala C		

UTafale 5.8: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala C) kumlambo iDuiwenhoks kwi- H8DUIW-EWR1

li-Metrics	li-EcoSpecs	Nee-TPCs
ii-ions zeetyuwa ezingezozendalo		
Sulphate as SO ₄	N/A	N/A
Sodium as Na	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 380 mg/L.	i-95 th percentile yeenkcukacha mayibe ngu 300 - 380 mg/L.
Magnesium as Mg	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 67 mg/L.	i- 95 th percentile yeenkcukacha mayibe ngu 53.5 - 67 mg/L.
Calcium as Ca	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 55 mg/L.	i- 95 th percentile yeenkcukacha mayibe ngu 44 - 55 mg/L.
Chloride as Cl	i-95 th percentile yeenkcukacha mayibe ngu ≤ 800 mg/L.	i- 95 th percentile yeenkcukacha mayibe ngu 640 - 800 mg/L.
Potassium as K	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 9 mg/L.	i- 95 th percentile yeenkcukacha mayibe ngu 7 - 9 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 270 mS/m.	i- 95 th percentile yeenkcukacha mayibe ngu 210 - 270 mS/m.
pH	i- 5 th percentile yeenkcukacha mayibe ngu 6.5. – 8.0, ze i- 95 th percentile ibe ngu 8.0 - 8.8.	i- 5 th percentile yeenkcukacha mayibe ngu ≤ 6.3 ze i-95 th percentile ibe ngu ≥ 8.6.
Ubushushu ^(a)	Ubushushu bendalo obuqhelekileyo.	Qalisa uhlolo olusisiseko loku kwahluka.
I-oksijini enyibilikisiweyo ^(a) (DO)	i-5 th percentile yeenkcukacha mayibe ngu ≥ 7.0 mg/L.	i- 5 th percentile yeenkcukacha mayibe ngu 7.2 - 7.0 mg/L. Qalisa uhlolo olusisiseko loku kwahluka
Ubukho bodaka ^(a)	Inguqu ebonakalayo yodaka ibangelwa ziiinguqulelo ezingephi ezenziwa ngabantu (umzekelo, ukukhiwa kwegraveli kumphezulu womjelo). Kulindeleke ukuba kubekho iintlenge kwindawo yokuphilisana.	Qalisa uhlolo olusisiseko loku kwahluka.
Izondlo		
TIN	i-50 th percentile yeenkcukacha mayibe ngu ≤ 0.25 mg/L.	i- 50 th percentile yeenkcukacha mayibe ngu 0.2 - 0.25 mg/L.
PO ₄ -P	i- 50 th percentile yeenkcukacha mayibe ngu ≤ 0.015 mg/L.	i-50 th percentile yeenkcukacha mayibe ngu 0.012 - 0.015 mg/L.
Ukwahluka ngokwemvakalelo		
Chl-a phytoplankton	i-50 th percentile yeenkcukacha mayibe ngu < 15 µg/L.	i- 50 th percentile yeenkcukacha mayibe ngu 12 - 15 µg/L.
Chl-a periphyton	i-50 th percentile yeenkcukacha mayibe ngu ≤ 12 mg/m ² .	i- 50 th percentile yeenkcukacha mayibe ngu 10 - 12 mg/m ² .
Iityhefu		
Fluoride	i-50 th percentile yeenkcukacha mayibe ngu ≤ 1.5 mg/L.	i- 50 th percentile yeenkcukacha mayibe ngu 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	i-50 th percentile yeenkcukacha mayibe ngu ≤ 0.015 mg/L.	i-50 th percentile yeenkcukacha mayibe ngu 0.012 - 0.015 mg/L.
Ezinye iityhefu	i- 95 th percentile yeenkcukacha mayibe phakathi kweqondo lekwaliti yamanzi eliqhelekileyo (iTarget Water Quality Range ngelasemzini (i-TWQR)njengoko kwaxelwayo nguDWAF (ngo1996) okanye ke kumda weBakala A njengoko kwaxelwayo nguDWAF (ngo2008).	Lilindelekile ifuthe ukuba i- 95 th percentile yeenkcukacha ibetha ngaphaya kwe-TWQR eyaxelwayo nguDWAF (ngo1996) okanye ke kumda ophezulu weBakala A ngengoko kwaxelwayo nguDWAF (ngo2008).

(a) N/A- khange kubekho zinkcukacha kolu uvavanyo.

UTafle 5.9: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iGoukou kwi-H9GOUK-EWR2

UMLAMBO		Umlambo iGoukou	WATER QUALITY MONITORING POINTS	
I-WQRU			RC	Reference Condition (RC) was represented by the A Category benchmark tables in DWAF (2008), as no other data were available to describe natural state.
ISIKHUNDLA EWR SITE		SE- H9GOUK-EWR2	PES	DWS gauging weir H9H005Q01 (2007 – 2014; n = 63 - 71, F = 52).
UVAVANYO NAKHO		LOKUBA	Inqanaba lokuba nakho: 3	
IZithako zeKwaliti yaManzi		Ixabiso le-PES		IBakala/Amagqabantshintshi
lityuwa ezingezozendalo (mg/L)	SO ₄	N/A		N/A
	Na	650.4		Zibetha ngaphaya kwe ≤ 70 mg/L (TWQR) zineda kwezolimo: kunkcenkeshelo.
	Mg	79.0		Asikho isikhokelo.
	Ca	57.1		Asikho isikhokelo.
	Cl	1081.3		Zibetha ngaphaya kwe ≤ 100 mg/L (TWQR) zineda kwezolimo: kunkcenkeshelo.
Izondlo (mg/L)	K	20.4		Asikho isikhokelo.
	SRP	0.085		D
Ukwahluka kwendalo	TIN	0.055		A
	pH (5 th – 95 th %)	6.6 and 8.35		B
	Ubushushu (°C)	N/A		A/B. Ifuthe lilindeleke emanzini amancinci.
	i-oksijini enyibilikisiweyo	N/A		B. ifuthe lilindeleke emanzini amancinci.
	Ubukho bodaka (NTU)	N/A		A/B. Inguqu yodaka ibonakala ibangelwa ikakhulu yindalo.
	Ukutsala umbane (mS/m)	408.4		E/F
Ukwahluka ngokwemvakalelo	Chl a: phytoplankton	N/A		N/A
	Ubukhulu bezinto eziphilayo ezingenamathambo	51.2% Inqaku le-SASS =113 Inqaku le-ASPT = 6.6		D
	ii-Diatoms	14.4 and 11.0		C/D (n = 2; eyoMqungu neyeKhala ku2014)
	Inqaku yeentlanzi lemigqeku	47.4%		D
iityhefu	Ammonia (as N)	0.01		A
	Fluoride (as F)	0.59		A
LULONKE UHLELO LWESIKHUNDLA			IBakala C/D	

UTafle 5.10: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala C/D) kumlambo iGoukou kwi- GOUK-EWR2

ii-Metrics	ii-EcoSpecs	ii-TPCs
II-ions zeetyuwa ezingezozendalo		
Sulphate as SO ₄	N/A	N/A
Sodium as Na	i-95 th percentile yeenkukacha iphakathi kwe ≤ 650 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 520 - 650 mg/L.
Magnesium as Mg	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 80 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 64 - 80 mg/L.
Calcium as Ca	i-95 th percentile yeenkukacha iphakathi kwe ≤ 55 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 44 - 55 mg/L.
Chloride as Cl	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 1 000 mg/L.	i-95 th percentile yeenkukacha iphakathi kwe 800 - 1 000 mg/L.
Potassium as K	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 20 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 16 - 20 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 400 mS/m.	i- 95 th percentile yeenkukacha iphakathi kwe 320 - 400 mS/m.
pH	i- 5 th percentile yeenkukacha mayibe ngu 6.5. – 8.0, ze i- 95 th percentile ibe ngu 8.0 - 8.8.	i- 5 th percentile yeenkukacha ngu ≤ 6.3 ze i- 95 th percentile ibe ngu ≥ 8.6.
Ubushushu ^(a)	Uluhlu lobushushu bendalo.	Qalisa uhlolo olusisiseko loku kwahluka.
I-oksijini enyibilikisiweyo ^(a)	i- 5 th percentile yeenkukacha iphakathi kwe ≥ 7.0 mg/L.	i-5 th percentile yeenkukacha iphakathi kwe 7.2 - 7.0 mg/L. Qalisa uhlolo olusisiseko loku kwahluka.
Ubukho bodaka ^(a)	Inguqu ebonakalayo yodaka ibangelwa ziiinguqulelo ezingephi ezenziwa ngabantu. Kulindeleke ukuba kubekho iintlenge kwindawo yokuphilisana	Qalisa uhlolo olusisiseko loku kwahluka.
Izondlo		
TIN-N	i-50 th percentile yeenkukacha iphakathi kwe ≤ 0.25 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.2 - 0.25 mg/L.
PO ₄ -P	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 0.125 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.1 - 0.125 mg/L.
Ukwahluka ngokwemvakalelo^(a)		
Chl-a phytoplankton	i-50 th percentile yeenkukacha iphakathi kwe < 15 µg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 12 - 15 µg/L.
Chl-a periphyton	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 21 mg/m ² .	i- 50 th percentile yeenkukacha iphakathi kwe 17 - 21 mg/m ² .
Iityhefu		
Fluoride	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 1.5 mg/L.	i-50 th percentile yeenkukacha iphakathi kwe 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	The 50 th percentile yeenkukacha iphakathi kwe ≤ 0.015 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe- 0.012 - 0.015 mg/L.
Ezinye iityhefu	i- 95 th percentile yeenkukacha ikwi- TWQR eyaxelwayo nguDWAF (ngo1996) okanye ke kumda weBakala A owaxelwayo nguDWAF (ngo2008).	Likho ifuthe elilindelekileyo ukuba ngaba i- 95 th percentile yeenkukacga ibetha ngaphaya kwe-TWQR njengoko kwaxelwayo nguDWAF (ngo1996) okanye ke kumda ophezulu weBakala A njengoko kwaxelwayo nguDWAF (ngo2008).

(a) N/A: Bezingekho iinkukacha zolu vavanyo.

UTafle 5.11: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iTouws kwi - J1TOUW-EWR3

UMLAMBO		Umlambo iTouws	IINDAWO ZOHLOLO LWEKWALITI YAMANZI	
I-WQRU			I-RC	N/A
ISIKHUNDLA SE-EWR		J1TOUW-EWR3	I-PES	J1H018Q01 (Ikhawudi yoBugcisa boLawulo lwaManzi (i-WMS) ngu-102147), ume kumphezulu womjelo weSikhundla se- EWR. (iinkcukacha zibhaliswe ngo: 2000 – 2014; inani leesampulu (n) = ± 128).
Uvavanyo lokuba nakho		2.5		
iZithako zeKwaliti yaManzi		Ixabiso le-PES	IBakala/Amagqabantshintshi	
lityuwa ezingezozendalo (mg/L)	SO ₄	N/A	Zisetyenziswe zonke izikhokelo (zade zaphela) ngenxa yeqondo eliphezulu letyuwa emanzini kumhlaba okuloo mmandla. Kuthelakelelwa ukuba iqondo letyuwa emanzini liya konyuka ngenxa yamanzi abuya angene emjelweni emva konkcenkceshelo. Akukho maziko makhulu eedolophu akhoyo kulo mmandla.	
	Na	2 016.9		
	Mg	370.1		
	Ca	258.2		
	Cl	3 494.6		
Izondlo (mg/L)	K	37.06		
	SRP	0.033	D	
	TIN	0.079	A	
Ukwahluka kwendalo	pH (5 th – 95 th %)	7.6 no-8.6	B	
	Ubushushu (°C)	N/A	B. Ifuthe layo lilindeleke ukuba libe phantsi.	
	I-oksijini enyibilikisiweyo	N/A	B. Ifuthe layo lilindelekile ukuba libe phantsi, nangona iinkcukacha zesikhundla zisabonisa amanqanaba aphezulu ayo nje.	
	Ubukho bodaka (NTU)	N/A	B. Inguqu yodaka ikhangeleka ikakhulu iyeyendalo, noxa zikho nezo nguqulelo zincinci zibangelwa ngabantu.	
	Ukutsala umbane (mS/m)	1181.8		
Ukwahluka kwemvakalelo	Chl a: phytoplankton	N/A	N/A	
	Ubukhulu bezinto eziphilayo ezingenamathambo	74.0%	C	
	iiDiatoms	8.6 (i-avareji)	D	
	Inqaku lomgqeku weentlanzi	56.8%	D	
iityhefu	Ammonia (as N)	0.034	A	
	Fluoride (as F)	0.43	A	
LULONKE UHLELO LWESIKHUNDLA			IBakala B/C	

Qaphela:

I-RC: Lulwazi olusezandleni sencutshe yekwaliti yamanzi olungeemeko zekwaliti yamanzi. Bekusakusetyenziswa ulwazi losetyenziso lomhlaba mandulo kuba zazingekho iinkcukacha ze-RC ngoko, futhi ke zona iitafle zolingiso zeBakala A ekwathethwa ngazo nguDWAF (ngo2008) zazisakucingelwa njengezingafanelekanga ngenxa yobukho betyuwa emanzini obabubangelwa luhlobo oluthile kuloo mmandla.

UTafle 5.12: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala B/C) kumlambo ITouws kwi-J1TOUW-EWR3

ii-Metrics	ii-EcoSpecs	ii-TPCs
Ii-ion zeetyuwa ezingezozendalo		
Sulphate as SO ₄	N/A	
Sodium as Na	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 2000 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 1600 - 2000 mg/L.
Magnesium as Mg	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 370 mg/L.	i-95 th percentile yeenkcukacha iphakathi kwe 300 - 370 mg/L.
Calcium as Ca	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 260 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 200 - 260 mg/L.
Chloride as Cl	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 3500 mg/L.	i-95 th percentile yeenkcukacha iphakathi kwe 2800 - 3500 mg/L.
Potassium as K	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 37 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 30 - 37 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 1100 mS/m.	i- 95 th percentile yeenkcukacha iphakathi kwe 880 - 1100 mS/m.
pH	i- 5 th percentile yeenkcukacha ngu 6.5 – 8.0, ze i-95 th percentile ibe ngu 8.0 - 8.8.	i- 5 th percentile yeenkcukacha ngu ≤ 6.3 ze i- 95 th percentile ibe ngu ≥ 8.6.
Ubushushu	Uluhlu lobushushu bendalo.	Qalisa uhlolo olusisiseko loku kwahluka.
i-oksijini enyibilikisiweyo	i-5 th percentile yeenkcukacha iphakathi kwe ≥ 7.0 mg/L.	i-5 th percentile yeenkcukacha iphakathi kwe 7.2 - 7.0 mg/L. Qalisa uhlolo olusisiseko loku kwahluka.
Ubukho bodaka	Inguqu ebonakalayo yodaka ibangelwa ziilinguqulelo ezingephi ezenziwa ngabantu. Kulindeleke ukuba kubekho iintenge	Qalisa uhlolo olusisiseko loku kwahluka.
Izondlo		
TIN-N	i-50 th percentile yeenkcukacha iphakathi kwe ≤ 0.25 mg/L.	i-50 th percentile yeenkcukacha iphakathi kwe 0.2 - 0.25 mg/L.
PO ₄ -P	i- 50 th percentile yeenkcukacha iphakathi kwe ≤ 0.075 mg/L.	i- 50 th percentile yeenkcukacha iphakathi kwe 0.06 - 0.075 mg/L.
Ukwahluka ngokwemvakalelo		
Chl-a phytoplankton	i- 50 th percentile yeenkcukacha iphakathi kwe < 15 µg/L.	i-50 th percentile yeenkcukacha iphakathi kwe 12 - 15 µg/L.
Chl-a periphyton	i-50 th percentile yeenkcukacha iphakathi kwe ≤ 21 mg/m ² .	i- 50 th percentile yeenkcukacha iphakathi kwe 17 - 21 mg/m ² .
Iityhefu		
Fluoride	i- 50 th percentile yeenkcukacha iphakathi kwe ≤ 1.5 mg/L.	i- 50 th percentile yeenkcukacha iphakathi kwe 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	i- 50 th percentile yeenkcukacha iphakathi kwe ≤ 0.015 mg/L.	i- 50 th percentile yeenkcukacha iphakathi kwe 0.012 - 0.015 mg/L.
Ezinye iityhefu	i- 95 th percentile yeenkcukacha ikwi - TWQR ekuthethwa ngayo kuDWAF (1996) okanye kumda weBakala A Category ekuthethwa ngayo kuDWAF (2008).	Likho ifuthe elilindelekileyo ukuba ngaba i- 95 th percentile yeenkcukacga ibetha ngaphaya kwe-TWQR njengoko kwaxelwayo nguDWAF (ngo1996) okanye ke kumda ophezulu weBakala A njengoko kwaxelwayo nguDWAF (ngo2008).

(a) N/A: Azikhange zibekho iinkcukacha kolu hlobo.

UTafle 5.13: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iGamka J2GAMK-EWR4

UMLAMBO	Umlambo iGamka	IINDAWO ZOHLOLO LWEKWALITI YAMANZI	
I-WQRU		I-RC	Kusetyenziswe iitafle zolingiso zeBakala A ezisuka kuDWAF (2008).
ISIKHUNDLA SE-EWR	J2GAMK-EWR4	I-PES	Iinkcukacha zifunyenwe kwidonga elinqamlezileyo lokuthatha umlinganiselo lika DWS elingu-J2H016Q01 (ikhowudi ye-WMS engu102173), elime kumazantsi onxweme lwedama iGamkapoort, nakumphezulu womjelo wesikhundla seEWR site. (iinkcukacha ezibhalisiweyo: 2007 – 2014; n = 127).
Uuvavanyo lokuba nakho		3.0	
IZithako zeKwaliti yaManzi		Ixabiso le-PES	Ibakala/Amagqabantshintshi
lityuwa ezingezozendalo (mg/L)	SO ₄	N/A	
	Na	114.0	Ibetha ngaphaya kwe-70mg/l (TWQR) isetyenziswa kwezolimo: kunkcenkeshelo
	Mg	20.5	Asikho isikhokelo
	Ca	57.6	Asikho isikhokelo
	Cl	155.5	Ibetha ngaphaya kwe-100mg/l (TWQR) isetyenziswa kwezolimo: kunkcenkeshelo
	K	7.9	Asikho isikhokelo
Izondlo (mg/L)	SRP	0.07	D
	TIN	0.523	B
Ukwahluka kwendalo	pH (5 th – 95 th %)	7.4 and 8.6	
	Ubushushu (°C)	N/A	Ifuthe lilindelekile njengoko isikhundla sikumazantsi omjelo wedama iGamkapoort (constructed in 1970)
	i-oksijini enyibilikisiweyo	N/A	
	Ubukho bodaka (NTU)	N/A	
Ukutsala umbane (mS/m)	97.5	C. ubukho betyuwa bendalo bulindeleke ukuba bube ngaphezulu kwe- 30Ms/m elixabiso lolingiso leBakala A ekuthethwa ngalo kuDWAF (2008)	
Ukwahluka ngokwemvakalelo	Chl a: phytoplankton	N/A	N/A
	Ubukhulu bezinto eziphilayo ezingenamathambo	61.4%	C/D
	iiDiatoms	9.9	D
	Inqaku lemigqeku yeentlanzi	60.4%	C/D
lityhefu	Ammonia (as N)	0.015	A
	Fluoride (as F)	0.53	A
LULONKE UHLELO LWESIKHUNDLA		IBakala B/C	

UTafle 5.14: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala B/C) kumlambo iGamka River kwi- J2GAMK-EWR4

iiMetrics	iiEcoSpecs: i-PES	i-TPCs: i-PES
ii-ion zeetyuwa ezingezozendalo		
Sulphate as SO ₄	N/A	N/A
Sodium as Na	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 114 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 90 - 114 mg/L.
Magnesium as Mg	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 20 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 16 - 20 mg/L.
Calcium as Ca	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 58 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 47 - 58 mg/L.
Chloride as Cl	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 155 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 124 - 155 mg/L.
Potassium as K	i-95 th percentile yeenkukacha iphakathi kwe ≤ 8 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 6.5 - 8.0 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	i-95 th percentile yeenkukacha ngu ≤ 100 mS/m.	i- 95 th percentile yeenkukacha iphakathi kwe 80 - 100 mS/m.
pH	i- 5 th percentile yeenkukacha ngu 5.9 - 6.5, ze i- 95 th percentile ibe ngu 8.0 - 8.8.	i- 5 th percentile yeenkukacha ngu ≤ 5.7 ze i- 95 th percentile ibe ngu ≥ 8.6.
Ubushushu ^(a)	Ikho inguqu ephakathi eyenzekayo ngenxa yedama iGamkapoort kumphezulu womjelo.	Qalisa uhlolo olusisiseko loku kwahluka.
i-oksijini enyibilikisiweyo ^(a)	i- 5 th percentile yeenkukacha iphakathi kwe ≥ 7.0 mg/L. Nangona likho ifuthe elilindelekileyo ngenxa yedama iGamkapoort kumphezulu womjelo nje, ubukhulu bomlambo buya kunceda ekulithomalaliseni eli futhe.	i- 5 th percentile of the data is between 7.2 - 7.0 mg/L. Qalisa uhlolo olusisiseko loku kwahluka
Ubukho bodaka ^(a)	Inguqu ebonakalayo yodaka ibangelwa ziiinguqulelo ezingephi ezenziwa ngabantu. Kulindeleke ukuba kubekho iintenge	Qalisa uhlolo olusisiseko loku kwahluka.
Izondlo		
TIN-N	i-50 th percentile yeenkukacha iphakathi kwe ≤ 0.7 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.56 - 0.7 mg/L.
PO ₄ -P	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 0.125 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.1 - 0.125 mg/L.
Ukwahluka kwendalo		
Chl-a phytoplankton	i- 50 th percentile yeenkukacha iphakathi kwe < 15 µg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 12 - 15 µg/L.
Chl-a periphyton	i- 50 th percentile yeenkukacha ngu ≤ 21 mg/m ² .	i- 50 th percentile yeenkukacha iphakathi kwe 17 - 21 mg/m ² .
Toxics		
Fluoride	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 1.5 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 0.015 mg/L.	i-50 th percentile yeenkukacha iphakathi kwe 0.012 - 0.015 mg/L.
Ezinye iityhefu	i- 95 th percentile yeenkukacha ikwi - TWQR ekuthethwa ngayo kuDWAF (1996) okanye kumda weBakala A	Likho ifuthe elilindelekileyo ukuba ngaba i- 95 th percentile yeenkukacga ibetha ngaphaya kwe-TWQR njengoko kwaxelwayo nguDWAF (ngo1996) okanye ke kumda

iiMetrics	iiEcoSpecs: i-PES	i-TPCs: i-PES
	Category ekuthethwa ngayo kuDWAF (2008).	ophezulu weBakala A njengoko kwaxelwayo nguDWAF (ngo2008).

(a) N/A: Khange zibekho iinkcukacha kolu hlolo.

UTafle 5.15: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iBuffels kwi-J1BUFF-EWR5

UMLAMBO		Umlambo iBuffels	IINDAWO ZOHOLOLO LWEKWALITI YAMANZI	
I-WQRU			I-RC	linkcukacha zifunyenwe kudonga olunqamlezileyo lokuthatha umlinganiselo luka DWS olungu-J1H028Q01 (ikhowudi ye-WMS engu-102152), olumi kumazantsi onxweme lwedama i- Floriskraal, nakumphezulu womjelo wesikhundla se-EWR. Qaphela ukuba indawo yohlolo ayikho kwingingqi enye yeNqanaba II nesikhundla se- EWR; Kodwa ke, le ibiyindawo yeenkcukacha ephakathi kwedama nesikhundla eso. (linkcukacha ezibhalisiweyo: 1972 – 1977; n = 54, ukutsala: n = 33).
ISIKHUNDLA EWR	SE-	J1BUFF-EWR5	PES	linkcukacha zifunyenwe kudonga olunqamlezileyo lokuthatha umlinganiselo likaDWS elingu-J1H028Q01 (ikhowudi yeWMS engu-102152) (Inkcukacha ezibhalisiweyoData record: 2010 – 2014; n = 44).
Uvavanyo lokuba nakho		Inqanaba lokuba nakho: 2.5		
IZithako zeKwaliti yaManzi		Ixabiso le-PES	IBakala/Amagqabantshintshi	
lityuwa ezingezozendalo (mg/L)	SO ₄	61.42	Asikho isikhokelo	
	Na	81.44	Zibetha ngaphaya kwe ≤ 70 mg/L (TWQR) zisetyenziswa kwezolimo: kunkcenkeshelo.	
	Mg	25.2	Asikho isikhokelo	
	Ca	48.68	Asikho isikhokelo.	
	Cl	124.0	Zibetha ngaphaya kwe ≤ 100 mg/L (TWQR) zisetyenziswa kwezolimo: kunkcenkeshelo.	
	K	6.11	Asikho isikhokelo.	
Izondlo (mg/L)	SRP	0.015	B. Amanqana ancipha ukusukela kwii-1970s.	
	TIN	0.26	A/B. Ayikho inguqu ukusukela kwii-1970s.	
Ukwahluka kwendalo	pH (5 th – 95 th %)	7.5 no- 8.5	B. Ayikho inguqu ukusukela kwii-1970s.	
	Ubushushu (°C)	N/A	Likho ifuthe elindelekileyo njengoba isikhundla sikumazantsi omjelo wedama elikhulu i-Floriskraal (elokhiwa ngo-1965).	
	i-oksijini enyibilikisiweyo	N/A		
	Ubukho bodaka (NTU)	N/A		
Ukutsala umbane (mS/m)	78.1	Ayikho inguqu ukusukela kwii-1970s.		
Ukwahluka ngokwemvakalelo	Chl a: phytoplankton	N/A	N/A	
	Ubukhulu bezinto eziphilayo ezingenamathambo	72.0%	C	
	iiDiatoms	11.2 (average)	C/D	
	Inqaku lemigqeku yeentlazi	83.7%	B	
lityhefu	Ammonia (as N)	0.017	A	
	Fluoride (as F)	0.66	A. Ayikho inguqu ukusukela kwii-1970s.	
LULONKE UHLELO LWESIKHUNDLA			IBakala B/C	

UTafle 5.16: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala B/C) kumlambo iBuffels River kwi- J1BUFF-EWR5

ii-Metrics	ii-EcoSpecs: PES	ii-TPCs: PES
Ii-ions zeetyuwa zendalo		
Sulphate as SO ₄	i-95 th percentile yeenkcukacha iphakathi kwe ≤ 60 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 48 - 60 mg/L.
Sodium as Na	i-95 th percentile yeenkcukacha iphakathi kwe ≤ 80 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 64 - 80 mg/L.
Magnesium as Mg	i-95 th percentile yeenkcukacha iphakathi kwe ≤ 25 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 20 - 25 mg/L.
Calcium as Ca	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 50 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 40 - 50 mg/L.
Chloride as Cl	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 125 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 100 - 125 mg/L.
Potassium as K	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 6.0 mg/L.	i- 95 th percentile yeenkcukacha iphakathi kwe 4.8 - 6.0 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	i- 95 th percentile yeenkcukacha iphakathi kwe ≤ 85 mS/m.	i- 95 th percentile yeenkcukacha iphakathi kwe 68 - 85 mS/m.
i-pH	i- 5 th percentile yeenkcukacha iphakathi kwe 5.9 - 6.5, ze i-95 th percentile ibe phakathi kwe- 8.0 - 8.8.	i- 5 th percentile yeenkcukacha ngu ≤ 5.7 ze i-95 th percentile ibe ngu ≥ 8.6.
Ubushushu	Inguqu ephakathi yobushushu ilindelekile ngenxa yedama iFloriskraal kumphezulu womjelo.	Qalisa uhlolo olusisiseko loku kwahluka.
i-oksijini enyibilikisiweyo	i-5 th percentile yeenkcukacha iphakathi kwe ≥ 7.0 mg/L. Nangona likhona nje ifuthe elilindelekileyo ngenxa yedama iFloriskraal kumphezulu womjelo, ubukhulu bomlambo buya kulithomalalisa eli futhe..	1- 5 th percentile yeenkcukacha iphakathi kwe 7.2 - 7.0 mg/L. Qalisa uhlolo olusisiseko loku kwahluka.
Ubukho bodaka	Inguqu ebonakalayo yodaka ibangelwa ziiinguqulelo ezingephi ezenziwa ngabantu. Kulindeleke ukuba kubekho iintlenge	Qalisa uhlolo olusisiseko loku kwahluka.
Izondlo		
TIN-N	i-50 th percentile yeenkcukacha iphakathi kwe ≤ 0.48 mg/L.	i-50 th percentile yeenkcukacha iphakathi kwe 0.38 - 0.48 mg/L.
PO ₄ -P	i- 50 th percentile yeenkcukacha iphakathi kwe ≤ 0.015 mg/L.	i-50 th percentile yeenkcukacha iphakathi kwe 0.012 - 0.015 mg/L.
Ukwahluka ngokwemvakalelo		
Chl-a phytoplankton	i- 50 th percentile yeenkcukacha iphakathi kwe < 10 µg/L.	i- 50 th percentile yeenkcukacha iphakathi kwe 8 - 10 µg/L.
Chl-a periphyton	i- 50 th percentile yeenkcukacha iphakathi kwe ≤ 12 mg/m ² .	i- 50 th percentile yeenkcukacha iphakathi kwe 10 - 12 mg/m ² .
Iityhefu		
Fluoride	i-50 th percentile yeenkcukacha iphakathi kwe ≤ 1.5 mg/L.	i-50 th percentile yeenkcukacha iphakathi kwe 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	i-50 th percentile yeenkcukacha iphakathi kwe ≤ 0.015 mg/L.	i- 50 th percentile yeenkcukacha iphakathi kwe 0.012 - 0.015 mg/L.
Ezinye iityhefu	i- 95 th percentile yeenkcukacha ikwi - TWQR ekuthethwa ngayo kuDWAF (1996) okanye kumda weBakala A	Likho ifuthe elilindelekileyo ukuba ngaba i-95 th percentile yeenkcukacha ibetha ngaphaya kwe-TWQR njengoko kwaxelwayo

ii-Metrics	ii-EcoSpecs: PES	ii-TPCs: PES
	Category ekuthethwa ngayo kuDWAF (2008).	nguDWAF (ngo1996) okanye ke kumda ophezulu weBakala A njengoko kwaxelwayo nguDWAF (ngo2008).

(a) N/A: Khange kubekho ziinkcukacha kolu hlolo.

UTafle 5.17: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iGouritz kwi-J4GOUR-EWR6

UMLAMBO		Umlambo iGouritz	IINDAWO ZOHLLO LWEKWALITI YAMANZI	
I-WQRU			I-RC	linkcukacha zifunyenwe kudonga elinqamlezileyo lokuthatha umlinganiselo lukaDWS olungu-J4H002Q01 (ikhowudi ye-WMS engu-102201), elime kumjelo ophezulu wesikhundla se-EWR. (iinkcukacha ezibhalisiweyo: 1965 – 1967; n = 29)
ISIKHUNDLA EWR	SE-	J4GOUR-EWR6	I-PES	linkcukacha zifunyenwe kudonga olunqamlezileyo lokuthatha umlinganiselo lukaDWS olungu-J4H002Q01 (iinkcukacha ezibhalisiweyo: 2010 – 2014; n = 85).
Uvavanyo lokuba nakho		Inqanaba lokuba nakho - 3		
IZithako zeKwaliti yaManzi		Ixabiso le-PES	IBakala/Amagqabantshintshi	
lityuwa ezingezozendalo (mg/L)	SO ₄	693.0	Asikho isikhokelo, kodwa ziya kwehla okuya lihamba ixesha.	
	Na	964.0	Zibetha ngaphaya kwe- ≤ 70 mg/L (TWQR) zineda kwezolimo: kunkcenkeshelo. Ziya ngokwenyuka okuya lihamba ixesha.	
	Mg	127.0	Asikho isikhokelo.	
	Ca	123.3	Asikho isikhokelo,	
	Cl	1 289.3	Zibetha ngaphaya kwe ≤ 100 mg/L (TWQR) zineda kwezolimo: kunkcenkeshelo.	
	K	9.81	Azikho isikhokelo.	
Izondlo (mg/L)	SRP	0.015	B/C	
	TIN	0.05	A	
Ukwahluka kwendalo	pH (5 th – 95 th %)	7.8 no- 8.65	B	
	Ubushushu (°C)	N/A	Lincinci ifuthe elilindelekileyo	
	i-oksijini enyibilikisiweyo	N/A		
	Ubukho bodaka (NTU)	N/A		
	Ukutsala umbane (mS/m)	542.5	Ayikho inguqu eyenzekileyo ukusukela kwii1970s.	
Ukwahluka kwemvakalelo	Chl a: phytoplankton	N/A	N/A	
	Ubukhulu bezinto eziphilayo ezingenamathambo	75.0%	C	
	iiDiatoms	10.2 (i-avareji)	C/D	
	Inqaku lemigqeku yeentlanzi	50.1%	D	
lityhefu	Ammonia (as N)	0.015	A	
	Fluoride (as F)	1.082	A. zonyuke ngokuthe xhaxhe ukusukela kwii1960s.	
LULONKE UHLELO LWESIKHUNDLA			IBakala B/C	

UTafle 5.18: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala B/C) kumlambo iGouritz kwi- J4GOUR-EWR6

ii-Metrics	ii-EcoSpecs	ii-TPCs
Ii-ions zeetyuwa ezingezizo zendalo		
Sulphate as SO ₄	i-95 th percentile yeenkukacha iphakathi kwe- ≤ 690 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 550 - 690 mg/L.
Sodium as Na	i- 95 th yeenkukacha iphakathi kwe ≤ 960 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 770 - 960 mg/L.
Magnesium as Mg	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 130 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 105 - 130 mg/L.
Calcium as Ca	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 120 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 95 - 120 mg/L.
Chloride as Cl	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 1300 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 1050 - 1300 mg/L.
Potassium as K	i-95 th percentile yeenkukacha iphakathi kwe ≤ 10 mg/L.	i-95 th percentile yeenkukacha iphakathi kwe 8 - 10 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 550 mS/m.	i-95 th percentile yeenkukacha iphakathi kwe 450 - 550 mS/m.
I-pH	i-5 th percentile yeenkukacha iphakathi kwe 5.9 - 6.5, ze i- 95 th ibe percentile 8.0 - 8.8.	i- 5 th percentile yeenkukacha nguis ≤ 5.7 ze i- 95 th percentile ibe ngu ≥ 8.6.
Ubushushu	Uluhlu lobushushu bendalo.	Qalisa uhlolo olusisiseko loku kwahluka
I-oksijini enyibilikisiweyo	i- 5 th percentile yeenkukacha iphakathi kwe ≥ 7.0 mg/L.	i- 5 th percentile yeenkukacha iphakathi kwe 7.2 - 7.0 mg/L. Qalisa uhlolo olusisiseko loku kwahluka.
Ubukho bodaka	Inguqu ebonakalayo yodaka ibangelwa ziinguqulelo ezingephi ezenziwa ngabantu. Kulindeleke ukuba kubekho iintlenge	Qalisa uhlolo olusisiseko loku kwahluka
Izondlo		
TIN-N	i-50 th percentile yeenkukacha iphakathi kwe ≤ 0.25 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.2 - 0.25 mg/L.
PO ₄ -P	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 0.015 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.012 - 0.015 mg/L.
Ukwahluka ngemvakalelo		
Chl-a phytoplankton	i-50 th percentile yeenkukacha iphakathi kwe < 10 µg/L.	i-50 th percentile yeenkukacha iphakathi kwe 8 - 10 µg/L.
Chl-a periphyton	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 12 mg/m ² .	i-50 th percentile yeenkukacha iphakathi kwe 10 - 12 mg/m ² .
Iityhefu		
Fluoride	i-50 th percentile yeenkukacha iphakathi kwe ≤ 1.5 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	i-50 th percentile yeenkukacha iphakathi kwe ≤ 0.015 mg/L.	i-50 th percentile yeenkukacha iphakathi kwe 0.012 - 0.015 mg/L.
Ezinye iityhefu	i- 95 th percentile yeenkukacha ikwi - TWQR ngokubhaliweyo kuDWAF (1996) okanye kumda weBakala A Category ekuthethwa ngayo kuDWAF (2008).	Likho ifuthe elilindelekileyo ukuba ngaba i-95 th percentile yeenkukacha ibetha ngaphaya kwe-TWQR ngokubhaliweyo kuDWAF (ngo1996) okanye ke kumda ophezulu weBakala A njengoko kwaxelwayo nguDWAF (ngo2008).

(a) N/A: Khange kubekho zinkukacha zikhoyo kolu hlolo.

UTafle 5.19: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iKeurbooms kwi-K6KEUR-EWR8

UMLAMBO	Umlambo iKeurbooms River	IINDAWO ZOHLOLO LWEKWALITI YAMANZI	
I-WQRU		RC	Kusetyenziswe iitafle zolingiso zeBakala A ngokubhaliweyo nguDWAF (2008).
ISIKHUNDLA SE-EWR	K6KEUR-EWR8	PES	Iinkcukacha zifunyenwe kudonga olunqamlezileyo lokuthatha umlinganiselo luka DWS olungu-K6H001Q01 (ikhowudi ye-WMS engu-102295), olumi kude lee (kangange- 20 km) kumphezul womjelo wesikhundla se- EWR . (Iinkcukacha ezibhalisiweyo: 2007 – 2014; n = 121; Fluorine (F) = 107)
Uvavanyo lokuba nakho		Inqanaba lokuba nakho 3	
IZithako zeKwaliti yaManzi		Ixabiso le-PES	IBakala/AmagComment
lityuwa ezingezozendalo (mg/L)	SO ₄	27.90	Asikho isikhokelo.
	Na	70.24	Ngaphandle nje ko ≤ 70 mg/L (TWQR) zineda kwezolimo: kunkcenkeshelo.
	Mg	11.25	Asikho isikhokelo.
	Ca	12.08	Asikho isikhokelo.
	Cl	129.02	zibetha ngaphaya kwe ≤ 100 mg/L (TWQR) zineda kwezolimo: kunkcenkeshelo.
	K	2.76	Asikho isikhokelo.
Izondlo (mg/L)	SRP	0.012	B
	TIN	0.06	A
Ukwahluka kwendalo	pH (5 th – 95 th %)	6.6 no- 7.8	B
	Ubushushu (°C)	N/A	B. Iincinci ifuthe elilindelekileyo, nangona iinkcukacha zesikhundla zisabonisa amanqanaba aphezulu . B. Inguqu ebonakalayo yodaka ibangelwa ziiinguqulelo ezingephi ezenziwa ngabantu.
	I-oksijini enyibilikisiweyo	N/A	
	Ubukho bodaka (NTU)	N/A	
	Ukutsala umbane (mS/m)	54.6	B
Ukwahluka kwemvakalelo	Chl a: phytoplankton	N/A	
	Ubukhulu bezinto eziphilayo ezingenamathambo	64.0%	C
	iiDiatoms	9.9 (i-avareji)	C/D
	Inqaku lomgqeku weentlanzi	76.4%	C
lityhefu	Ammonia (as N)	0.001	A
	Fluoride (as F)	0.26	A
LULONKE UHLELO LWESIKHUNDLA		IBakala B	

UTafale 5.20: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala B) kumlambo iKeurbooms kwi- K6KEUR-EWR8

ii-Metrics	ii-Eco Specs	ii-TPCs
Ii-ion zeetyuwa ezingezozendalo		
Sulphate as SO ₄	i-95 th percentile yeenkukacha iphakathi kwe ≤ 28 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 22 - 28 mg/L.
Sodium as Na	i-95 th percentile yeenkukacha iphakathi kwe ≤ 70 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 56 - 70 mg/L.
Magnesium as Mg	i-95 th percentile yeenkukacha iphakathi kwe ≤ 12 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 10 - 12 mg/L.
Calcium as Ca	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 12 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 10 - 12 mg/L.
Chloride as Cl	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 130 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 104 - 130 mg/L.
Potassium as K	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 3 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 2.4 - 3.0 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	i-95 th percentile yeenkukacha iphakathi kwe ≤ 55 mS/m.	i- 95 th percentile yeenkukacha iphakathi kwe 45 - 55 mS/m.
I-pH	i-5 th and 95 th yeenkukacha iphakathi kwe 6.5 - 8.0.	I- 5 th ne- 95 th yeenkukacha ngu ≥ 6.3 no- ≤ 8.2.
Ubushushu	Natural temperature range.	Qalisa uhlolo olusisiseko loku kwahluka.
i-oksijini enyibilikisiweyo	i- 5 th percentile yeenkukacha iphakathi kwe ≥ 7.0 mg/L.	i-5 th percentile yeenkukacha iphakathi kwe 7.2 - 7.0 mg/L. Qalisa uhlolo olusisiseko loku kwahluka.
Ubukho bodaka	Inguqu ebonakalayo yodaka ibangelwa ziinguqulelo ezingephi ezenziwa ngabantu. Kulindeleke ukuba kubekho iintlenge.	Qalisa uhlolo olusisiseko loku kwahluka.
Izondlo		
TIN-N	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 0.25 mg/L.	i-50 th percentile yeenkukacha iphakathi kwe 0.2 - 0.25 mg/L.
PO ₄ -P	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 0.015 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.012 - 0.0715 mg/L.
Ukwahluka kwemvakalelo		
Chl-a phytoplankton	i- 50 th percentile yeenkukacha iphakathi kwe < 10 µg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 8 - 10 µg/L.
Chl-a periphyton	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 12 mg/m ² .	i- 50 th percentile yeenkukacha iphakathi kwe 9.6 - 12.0 mg/m ² .
Iityhefu		
Fluoride	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 1.5 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 0.015 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.012 - 0.015 mg/L.
Ezinye iityhefu	i- 95 th percentile yeenkukacha ikwi - TWQR ngokubhaliweyo kuDWAF (1996) okanye kumda weBakala A Category ekuthethwa ngayo kuDWAF (2008).	Likho ifuthe elilindelekileyo ukuba ngaba i-95 th percentile yeenkukacha ibetha ngaphaya kwe-TWQR ngokubhaliweyo kuDWAF (ngo1996) okanye ke kumda ophezulu weBakala A njengoko kwaxelwayo nguDWAF (ngo2008).

(a) N/A: Khangela kubekho zinkukacha zikhoyo kolu vavanyo.

UTafale 5.21: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo i-Olifants kwi- J3OLIF-EWR9

UMLAMBO		Umlambo i-Olifants	IINDAWO ZOHOLO LOWEKVALITI YAMANZI	
I-WQRU			I-RC	linkcukacha zolwazi malunga neemeko zekwaliti yamanzi nosetyenziso lomhlaba ebezisezandleni zenkcutshe yezekwaliti yamanzi ziye zafumaneka, futhi ke iitafle zolingiso zeBakala A ngokubhaliweyo kuDWAF (2008) ziye zabanwa zingafanelekanga zona.
ISIKHUNDLA SE-EWR		J3OLIF-EWR9	I-PES	linkcukacha zifumaneka kudonga olunqamlezileyo lokuthatha umlinganiselo lukaDWS olungu-J3H021Q01 (ikhowudi yeWMS ingu-102192) zaza zasetyenziswa kuvavanyo lwemeko yangoku kumazantsi womjelo wesikhundla seEWR nakumphezulu womjelo wedama iStompdrift. (iinkcukacha ezibhalisiweyo: 1982 – 1993; n = 127).
Uvavanyo lokuba nakho		Iqondo lokuba nakho: 2.5		
IZithako zeKwaliti yaManzi		Ixabiso le-PES	IBakala/Amagqabantsshintshi	
lityuwa ezingezozendalo (mg/L)	SO ₄	1 353.4	Asikho isikhokelo, kodwa ukujiya kwazo kuthande ukuba kwinganaba eliphezulu	
	Na	1 774.5	Zibetha ngaphaya kwe ≤ 70 mg/L (TWQR) zanceda kwezolimo: kunkcenkceshelo.	
	Mg	336.0	Asikho isikhokelo	
	Ca	284.4	Asikho isikhokelo	
	Cl	3 113	Zibetha ngaphaya kwe ≤ 100 mg/L (TWQR) zanceda kwezolimo: kunkcenkceshelo.	
	K	30.16	Asikho isikhokelo.	
Izondlo (mg/L)	SRP	0.019	B/C	
	TIN	0.11	A	
Ukwahluka kwendalo	pH (5 th – 95 th %)	7.3 no-9.0	B/C, kodwa kuthelekelelwa ukuba kunxulumene nenkangeleko yamanzi angaphantsi komhlaba.	
	Ubushushu (°C)	N/A	C. Lincinci ifuthe elilindelekileyo.	
	i-oksijini enyibilikisiweyo	N/A		
	Ubukho bodaka (NTU)	N/A	B/C. Ifuthe lilindelekile ngenxa yemfuyo eninzi nokhukuliseko lomhlaba olwandileyo kulo mmandla..	
	Ukutsala umbane (mS/m)	1 078.7	Iqondo lobukho beetyuwa emanzini ngokwendalo kulindeleke ukuba libe phezulu ngenxa yokwakheka komhlaba kulo mmandla.	
Ukwahluka ngokwemvakalelo	Chl a: phytoplankton	N/A	N/A	
	Ubukhulu bezinto eziphilayo ezingenamathambo	69.0%	C	
	iiDiatoms	6.0 (i-avareji)	D/E	
	Inqaku lemigqeku yeentlanzi	N/A		
lityhefu	Ammonia (as N)	0.038	B	
	Fluoride (as F)	0.678	A	
LULONKE UHLELO LWESIKHUNDLA		IBakala C		

(a) N/A: Khange kubekho zinkcukacha zikhoyo kolu vavanyo.

UTafle 5.22: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala C) kumlambo i-Olifants kwi-J3OLIF-EWR9

ii-Metrics	ii-Eco Specs	ii-TPCs
Ii-ions zeetyuwa ezingezozendalo		
Sulphate as SO ₄	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 1350 mg/L.	i-95 th percentile yeenkukacha iphakathi kwe 1080 - 1350 mg/L.
Sodium as Na	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 1775 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 1420 - 1775 mg/L.
Magnesium as Mg	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 335 mg/L.	i-95 th percentile yeenkukacha iphakathi kwe 270 - 335 mg/L.
Calcium as Ca	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 285 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 230 - 285 mg/L.
Chloride as Cl	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 3000 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 2400 - 3000 mg/L.
Potassium as K	i- 95 th percentile yeenkukacha iphakathi kwe ≤ 30 mg/L.	i- 95 th percentile yeenkukacha iphakathi kwe 24 - 30 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	The 95 th percentile of the data is between ≤ 1100 mS/m.	i-95 th percentile yeenkukacha iphakathi kwe 880 - 1100 mS/m.
I-pH	i-5 th percentile yeenkukacha iphakathi kwe 5.9 - 6.5, ze i- 95 th percentile ibe ngu 8.8 - 9.2.	i- 5 th percentile yeenkukacha ngu ≤ 5.7 ze i- 95 th percentile ibe ngu ≥ 9.0.
Ubushushu	Umphezulu we-Olifants uhamba ngaphantsi ubukhulu becala, kubekho namanzi abonakala kwiindawo ezithile. La asingomanzi angaphantsi komhlaba asuka kwii-akhwifa, kodwa ingamanzi asuka kumda we-vadose. Amaqondo obushushu awonyukileyo namanqanaba e- DO alindelekile kwezi meko. Kungoko ii-EcoSpecs nee-TPCs kunzima ukuzilungiselela oku kwahluka nje, endaweni yoko kumele zinxulumyaniswe nokuhlangabezana neemfuno neemvakalelo zebiotic ezinxulumene nohlolo.	Qalisa uhlolo olusisiseko
i-oksijini enyibilikisiweyo		Qalisa uhlolo olusisiseko loku kwahluka, apho lufuneka khona ukuba kunokwenzeka.
Ubukho bodaka	Inguqu ebonakalayo yodaka ibangelwa ziiinguqulelo ezingephi ezenziwa ngabantu. Kulindeleke ukuba kubekho iintlenge.	Qalisa uhlolo olusisiseko loku kwahluka.
Izondlo		
TIN-N	i-50 th percentile yeenkukacha iphakathi kwe ≤ 0.25 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.2 - 0.25 mg/L.
PO ₄ -P	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 0.025 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 0.02 - 0.025 mg/L.
Ukwahluka ngokwemvakalelo^(a)		
Chl-a phytoplankton	i- 50 th percentile yeenkukacha iphakathi kwe < 15 µg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 12 - 15 µg/L.
Chl-a periphyton	i-50 th percentile yeenkukacha iphakathi kwe ≤ 21 mg/m ² .	i- 50 th percentile yeenkukacha iphakathi kwe 17 - 21 mg/m ² .
Iityhefu		
Fluoride	i- 50 th percentile yeenkukacha iphakathi kwe ≤ 1.5 mg/L.	i- 50 th percentile yeenkukacha iphakathi kwe 1.2 - 1.5 mg/L.

ii-Metrics	ii-Eco Specs	ii-TPCs
Ammonia (NH ₃ -N)	i-50 th percentile yeenkukacha iphakathi kwe ≤ 0.044 mg/L.	i-50 th percentile yeenkukacha iphakathi kwe 0.035 - 0.044 mg/L.
Ezinye iityhefu	i- 95 th percentile yeenkukacha ikwi - TWQR ngokubhaliweyo kuDWAF (1996) okanye kumda weBakala A Category ekuthethwa ngayo kuDWAF (2008).	Likho ifuthe elilindelekileyo ukuba ngaba i-95 th percentile yeenkukacha ibetha ngaphaya kwe-TWQR ngokubhaliweyo kuDWAF (ngo1996) okanye ke kumda ophezulu weBakala A njengoko kwaxelwayo nguDWAF (ngo2008).

UTafle 5.23: Amabakala e-PES nohlolo lwesikhundla lulonke kumlambo iKammanassie kwi-J3KAMM-EWR10

MLAMBO	Umlambo iKammanassie	IINDAWO ZOHLOLO LWEKWALITI YAMANZI	
I-WQRU		RC	Azikhange zibekho iinkcukacha zovavanyo lwekwali yamanzi. Usetyenziso lomhlaba nolwazi olufumanekayo, iinkcukacha ze-diatom, iinkcukacha zekwali yamanzi kwisikhundla ngqo neenkukacha zophando – onke la mabakala asetyenzisiwe khonukuze inkutshe yezekwali ikwazi ukunika iibono zayo, iqambe nomzekelo we-PAI igqibe nangebakala elibumbeneyo lwekwali yamanzi kweso sikhundla siphatsi kophando.
ISIKHUNDLA SE-EWR SITE	J3KAMM-EWR10	PES	N/A
Uvavanyo lokuba nakho		Inqanaba lokuba nakho: 2	
IZithako zeKwali yaManzi		Ixabiso le-PES	IBakala/Amagqabantshishi
lityuwa ezingezozendalo (mg/L)	SO ₄	N/A	N/A
	Na	N/A	N/A
	Mg	N/A	N/A
	Ca	N/A	N/A
	Cl	N/A	N/A
	K	N/A	N/A
Izondlo (mg/L)	SRP	N/A	N/A
	TIN	N/A	N/A
Ukwahluka kwendawo	pH (5 th – 95 th %)	N/A	N/A
	Ubushushu (°C)	N/A	N/A
	i-oksijini enyibilikisiweyo	N/A	N/A
	Ubukho bodaka (NTU)	N/A	N/A
	Ukutsala umbane (mS/m)	N/A	N/A
Ukwahluka ngokwemvakalelo	Chl a: phytoplankton	N/A	N/A
	Izinto eziphilayo ezingenamathambo	C/D	
	iiDiatoms	C/D	Ikwaliti yamanzi yendalo kwesi sikhundla ibiphakathi. Amanqanaba ezondlo, ongcoliswo lwendalo nawobukho betyuwa emanzini anyukele phezulwana – apho amanqanaba obukho betyuwa emanzini edibene nawongcoliso lwendalo ethande ukuxaka. Inkqubela ebonakalayo ngekwali yamanzi eyenziwa bubukho be-diatom inganxulunyaniswa namaqondo aphezulu angenyanga yeKhala ku2014 wona abangela ukuba kube lula ukucocwa kokungcola, kuba imigqeku ye-diatom enxulunyaniswa namaqanaba aphezulu ayo ibixhaphake kakhulu
	Inqaku lemigqeku yeentlanzi	D	
lityhefu	Ammonia (as N)		
	Fluoride (as F)		
LULONKE UHLELO LWESIKHUNDLA		IBakala C	

a) N/A: Khangela kubekho nkcukacha kolu hlolo.

Qaphela ukuba azanelanga kuyaphi iinkcukacha zekwaliti yamanzi ngobugcisa bomlambo iKammanassie. Kungoko uvavanyo lwekwaliti yamanzi alo mlambo belungqiyame nje kuphela ngolwazi obelufumaneka neembono zenkcutshe yezekwaliti yamanzi.

UTafle 5.24: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iGroot Brak kwi-WQSU 1 & 2. (iimbono zenkcutshe yezekwaliti yamanzi)

UMLAMBO		Umlambo iGroot Brak	IINDAWO ZOHLOLO LWEKVALITI YAMANZI	
I-WQSU		WQSU 1 + 2	I-RC	Iitafile zomda ezingagqibekanga ngomlambo weBakala "A"
ISIKHUNDLA SE-EWR		Asikho	I-PES	Umlambo iGroot Brak kwidama i-Ernest Robertson - K2H005-Q01 (1983 – 1996; n = 29)
Uvavanyo lokuba nakho		Inqanaba lokuba nakho lisezantsi kakhulu kuba linxulunyaniswa namanzi aphuphumela ngaphandle kwedama, hayi nenkcukacha zangoku.		
IZithako zeKwaliti yaManzi		IXabiso	Ibakala/Amagqabantshintshi	
lityuwa ezingezozendalo (mg/L)	MgSO ₄		Azikho iinkcukacha	
	Na ₂ SO ₄			
	MgCl ₂			
	CaCl ₂			
	NaCl			
	CaSO ₄			
Izondlo (mg/L)	SRP (mg/l)	0.042	Kusetyenziswe iinkcukacha ezingagqibekanga	
	TIN (mg/l)	<0.25	Ibakala A	
Ukwahluka kwendalo	pH (5 th – 95 th %)	4.62	Zine-asidi ngendalo	
	Ubushushu		Azikho iinkcukacha	
	I-oksijini enyibilikisiweyo		Kulindeleke kube kwinqanaba eliphazulu	
	Ubukho bodaka (NTU)		Kulindeleke kube kwinqanaba eliphantsi	
	Ukutsala umbane (mS/m)	<30	Ibakala A	
Ukwahluka ngokwemvakalelo	Chl a: periphyton		Azikho iinkcukacha	
	Chl a: phytoplankton		Azikho iinkcukacha	
	Ubukhulu bezinto eziphilayo ezingenamathambo (ASPT)		Azikho iinkcukacha	
	Inqaku lemigqeku yeentlanzi		Azikho iinkcukacha	
lityhefu			Azikho iinkcukacha – kulindeleke ukuba zibe mbalwa	
LULONKE UHLELO LWESIKHUNDLA			A/B (ngokweembono zenkcutshe)	

UTafite 5.25: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala A/B) kumlambo i-Groot Brak kwi-WQSU 1&2

UMLAMBO		Umlambo iGroot Brak		IINDAWO ZOHLULO LWEKWALITI YAMANZI		
I-WQSU		WQSU 1&2		Okwangoku akukabikho sifishi sohlolo		
ISIKHUNDLA SE-EWR		GB 1		Okwangoku akukabikho sikhundla sohlolo		
Ukukholelwa kuvavanyo lwe-PES		Kwiqondo eliphantsi kakhulu kuba lukhutshelwe ngaphandle kwe-WQSU 3				
IZithako zekwaliti yamanzi		IBakala le-PES	ii-Ecospecs ze-WQ	Ingaba inkqubela iyatuneka?	I-TPC	Ixesha elimiyo lohlolo
Iityuwa ezingezozendalo (mg/L)	MgSO ₄		≤23 mg/L		I-95 th percentile ibe ngu < 23 mg/L	
	Na ₂ SO ₄		≤33 mg/L		I-95 th percentile ibe ngu < 33 mg/L	
	MgCl ₂	Azikhho	≤30 mg/L	N/A	I-95 th percentile ibe ngu < 30 mg/L	Ngenyanga
	CaCl ₂		≤57 mg/L		I-95 th percentile ibe ngu < 57 mg/L	
	NaCl		≤191 mg/L		I-95 th percentile ibe ngu < 191 mg/L	
Izondlo (mg/L)	SRP	IBakala = C.	≤0.012 mg/L	Ewe ukuya ku- A	I-50 th percentile ibe ngu e < 0.012 mg/L	Ngenyanga
	TIN	IBakala = A	≤0.25 mg/L	Hayi	I- percentile ibe ngu < 0.25 mg/L	Ngenyanga
	pH	Umlambo one-asidi ngendalo	< 7.9	Hayi	I-95 th percentile ibe ngu < 7.9	Ngenyanga
Ukwahluka kwendalo	Ubushushu	Azikhho iinkcukacha, kodwa kungekho nto ifihlile	Gcina uluhlu	N/A	Gcina uluhlu lwendalo	Ngenyanga
	I-oksijini enyibilikisiweyo	ingxaki kulo mlambo..	7 – 8 mg/L	N/A	I-5 th percentile ibe ngu > 7 mg/L	Ngenyanga
	Ubukho bodaka (NTU)		Inguqu ephakathi	N/A	Moderate change allowed	Ngenyanga
	Ukutsala umbane (mS/m)	IBakala = A	≤30 mS/m	Hayi	I-95 th percentile ibe ngu < 30 mS/m	Ngenyanga
Ukwahluka ngokwemvaka lelo	Chi a: periphyton	Azikhho iinkcukacha. Uhlolo ngokubona Kwindla & Silimeia. Alukho uphawu olubonakalayo lwe- algae.	≤12 mg/m ² (B category)	N/A	I-50 th percentile ibe ngu < 12 mg/m ²	Ngekota
	Chi a: phytoplankton	A (esi sifundo)	≤ 15 µg/L (IBakala B)		I-50 th percentile ibe ngu < 15 µg/L	
	Ubukhulu bezinto eziphilayo ezingenamathambo lelo	C (esi sifundo)				
Iityhefu	Inqaku lemigqeku yeentlanzi	Ukutyhefeka komphakathi womjelo				
	Ukutyhefeka komphakathi womjelo	Azikhho iinkcukacha. Kungenzeka ukuba zikho izibulala zinambuzane				
Bona ii-Ecospecs zeentlanzi nezezinto eziphilayo ezingenamathambo, ngokulandelelanayo						
Vavanya kuphela xa neziphumo zohlolo lwendalo zibonisa ukuba ikho ingxaki futhi isizathu sayo asaziwa.						

UTafife 5.26: Amabakala e-PES nohlolo lwesikhundla lulonke kumlambo iGroot Brak kwi- WQSU 3

UMLAMBO	Umlambo iGroot Brak	1.1.1 IINDAWO ZOHLOLO LWEKWALITI YAMANZI	
I-WQSU	WQSU 3	Moordkuij R. at Bariff - K1H005 (1979 – 1982 n = 91)	
ISIKHUNDLA SE-EWR	GB 1	K1H005 (2002 – 2006 n = 51)	
Uvavanyo lokuba nakho		Phakathi. Qaphela: Kuba ukhuthelwe ngaphandle kwenye iindawo yoboniselelo ngamanzi. Iinkcukacha zendalo ziyayixhasa ikwaliti yamanzi ethengiswayo.	
IZithako zeKwaliti yaManzi	Ixabiso le-RC	Ixabiso le-PES	IBakala/Amagqabantshintshi
Iityuwa ezingezozendalo (mg/L)	MgSO ₄		
	Na ₂ SO ₄		
	MgCl ₂		
	CaCl ₂		
	NaCl		
Izondlo (mg/L)	CaSO ₄		Azikho iinkcukacha
	SRP	0.006*	IBakala = C. kukho ukwenyuka okuthile ngokwesimbo
	TIN	0.04	IBakala = A. Isimbo sisesihle
	pH (5 th – 95 th %)	Ayibalwanga	6.6 – 7.9
Ukwahluka kwendalo	Ubushushu		Azikho iinkcukacha. Kodwa ke oku akuthethi kuthi oku kukwahluka okunengxaki, kuba akubangelwa ngamazantsi onjelo wedama.
	i-oksijini enyibilikisiweyo		Bekubonakala ubukho bentwana yodaka emanzini kutyelelo
	Ubukho bodaka (NTU)		IwangeyoKwindla nolwangeyeSilimela ku2007 (kodwa ke emva kweemvula ezinkulu).
Ukwahluka ngokwemvakalelo	Ukutsala umbane (mS/m)	40	Isimbo sihle kancinci
	Chl a: periphyton		Bekubonakala ubukho bentwana yodaka emanzini kutyelelo
	Chl a: phytoplankton		IwangeyoKwindla nolwangeyeSilimela. Azikho iimpawu ze-algae ezibonakalayo
	Ubukhulu bezinto eziphilayo ezingenamathambo (ASPT)	-	ngokwendalo; ASPT = 8.0; SASS = 192 (olu phando)
	Inqaku lemiggeku yeentlanzi	-	Olu phando

lityhefu				Azikho iinkukacha. Kungenzeka ukuba zibekho izibulala zinambuzane eizuka kwezolimo.
LULONKE UHLOLO LWESIKHUNDLA				
B (ngokomzekeliso we- PAI)				

UTafle 5.27: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala B) kumlambo i-Groot Brak - WQSU 3

UMLAMBO		INDAWO ZOHLOLO LWEKWALITI	
Umlambo iGroot Brak		Okwangoku asikabikho isitishi sohlolo	
3		Okwangoku asikabikho isikhundla sohlolo	
I-WQSU		DWAF WQ WMS	RHP
ISIKHUNDLA SE-EWR SITE		Phantsi – phakathi, kuba ukhutshelwe ngaphandle komlambo iMoordkui, linkcukacha zendalo zixhasa i-PES ethengisiweyo yekwaliti yamanzi.	
Ukukholelwa uhlobo lwe-PES		I-Ecospecs ze-WQ	
IZithako zeKwaliti yaManzi		Ibakala -PES	Inqubo ikho inkqubo efunekayo?
Iityuwa ezingezozendalo (mg/L)	MgSO ₄	≤23 mg/L	i-95 th percentile ibe ngu < 23 mg/L
	Na ₂ SO ₄	≤33 mg/L	i-95 th percentile ibe ngu < 33 mg/L
	MgCl ₂	≤30 mg/L	i-95 th percentile ibe ngu < 30 mg/L
	CaCl ₂	≤57 mg/L	i-95 th percentile ibe ngu < 57 mg/L
	NaCl	≤191 mg/L	i-95 th percentile ibe ngu < 191 mg/L
Izondlo (mg/L)	SRP	≤0.02 mg/L	i-50 th percentile ibe ngu < 0.02 mg/L
	TIN	≤0.25 mg/L	i-50 th percentile ibe ngu < 0.25 mg/L
Ukwahluka kwendalo	pH	< 7.9	i-95 th percentile ibe ngu < 7.9
	Ubushushu	Gcina uluhlu	Gcina uluhlu lwendalo
	i-oksijini enyibilikisiweyo	7 – 8 mg/L	i-5 th percentile ibe ngu > 7 mg/L
	Ubukho bodaka (NTU)	Inguqu iphakathi	Inguqu ephakathi ivumelekile
	Ukutsala umbane (mS/m)	IBakala = A	i-95 th percentile ibe ngu < 30 mS/m
Ukwahluka ngokwemvaka lelo	Chi a: periphyton	≤12 mg/m ² (IBakala B)	i-50 th percentile ibe ngu < 12 mg/m ²
	Chi a: phytoplankton	≤ 15 µg/L (IBakala B)	i-50 th percentile to be < 15 µg/L
Ubukhulu bezinto eziphilayo ezingenamathambo (ASPT)	A (olu phando)		
	C (olu phando)		
Inqaku lemiggeku yeentlanzi		Bona ii-Ecospecs zeentlanzi nezinto eziphilayo ezingenamathambo ngokulandelelanayo	

ityhefu	Ukutyhefeka komphakathi womjelo	Azikho iinkcukacha Azikho iinkcukacha. Kungenzeka ukuba zikho ezo zibulala zihambuzane zithile	Vavanya kuphela xa iziphumo zohlolo lwendalo zibonisa ukuba ikho ingxaki eqatseleyo futhi isizathu singaziwa.
---------	---------------------------------	--	---

UTafale 5.28: amabakala e-PES nohlolo lwesikhundla lulonke kumlambo iGroot Brak kwi-WQSU 4 (*ixabiso lomda liphuculwe) (iimbono zencutshe)

UMLAMBO		Umlambo iGroot Brak		IINDAWO ZOHLOLO LWEKWALITI YAMANZI	
I-WQSU	4	I-RC	Umlambo iGroot Brak. kwiWolwedans - K2H002-Q01 (1976 – 1978; n = 68)		
ISIKHUNDLA SE-EWR	Asikho	I-PES	K2H002 (2002 – 2006; n = 57)		
Uvavanyo lokuba nakho		Kuhle kakhulu. Isitishi sohlolo simi e-WQSU, kumphezulu wentsukaphi yendawo. Iinkcukacha zifunyenwe > kwiminyaka eli-15 phambi kokwakhiwa kwedama.			
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	Inqanaba/Amagqabantshintshi	
Iityuwa ezingezozendalo (mg/L)	MgSO ₄			Azikho iinkcukacha	
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
Izondlo (mg/L)	SRP	0.016*	0.037	IBakala = C. isimbo siyanda	
	TIN	0.04	0.075	IBakala = A. isimbo siyanda	
ssUkwakheka kwendalo	pH (5 th – 95 th %)		6.8 – 8.1	Kuxhomekeke kwisitishi sohlolo esimi ku-WQSU, ngaphezulu kwentsukaphi yendawo.	
	Ubushushu		Azikho iinkcukacha	Inokubakho ingxaki xa ucinga ngamazantsi omjelo wedama.	
	i-oksijini enyibilikisiweyo		Azikho iinkcukacha	Ifuna uhlobo	
	Ubukho bodaka (NTU)		Azikho iinkcukacha		
	Ukutsala umbane (mS/m)	30	52	IBakala = B. isimbo siyanda	
Ukwakheka ngokwemvakalelo	Chl a: periphyton			Azikho iinkcukacha	
	Chl a: phytoplankton			Azikho iinkcukacha	
	Ubukhulu bezinto eziphilayo ezingenamathambo (ASPT)			Azikho iinkcukacha	
	Inqaku lemigqeku yeentlanzi			Azikho iinkcukacha	
Iityhefu				Azikho iinkcukacha	
LOLUNKE UHLELO LWESIKHUNDLA		B/C (Iimbono zencutshe)			

UTafie 5.29: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs (zeBakala B/C) kumlambo i-Groot Brak kwi- WQSU 4

UMLAMBO		IINDAWO ZOHLOLO LWEKWALITI YAMANZI	
I-WQSU		DWAF WQ WMS	RHP
ISIKHUNDLA SE-EWR		Okwangoku, asikabikho isitishi sohlolo	
Ukukholelwa kuvavanyo lwe-PES		Okwangoku, asikabikho isitishi sohlolo	
Ukukholelwa kuvavanyo lwe-PES		Iphantsi kakhulu, kuba ikhutshelwe ngaphandle kwe-WQSU 3.	
IZithako zekwaliti yaManzi		IBakala le-PES	Ingaba ikho inkqubela efunekayo?
Ityuwa ezingezozendalo (mg/L)		Ecospecs ze-WQ	I-TPC
Izondlo (mg/L)			Ixesha elimiyo lohlolo
MgSO ₄		≤23 mg/L	i-95 th percentile ibe ngu < 23 mg/L
Na ₂ SO ₄		≤33 mg/L	i-95 th percentile ibe ngu < 33 mg/L
MgCl ₂		≤30 mg/L	i-95 th percentile ibe ngu < 30 mg/L
CaCl ₂	Azikhho	≤57 mg/L	i-95 th percentile ibe ngu < 57 mg/L
NaCl		≤191 mg/L	i-95 th percentile ibe ngu < 191 mg/L
SRP	IBakala = C.	≤0.025 mg/L	i-50 th percentile ibe ngu < 0.025 mg/L
TIN	IBakala = A	≤0.25 mg/L	i-50 th percentile ibe ngu < 0.25 mg/L
pH	Umlambo one-asidi ngendalo	< 7.9	i-95 th percentile ibe ngu < 7.9
Ubushushu	Azikhho iinkcukacha, kodwa asingatsho	Gcina uluhlu	Gcina uluhlu lwendalo
I-oksijini enyibilikisiweyo	ukuba ikho ingxaki kulo mlambo.	7 – 8 mg/L	i-5 th percentile ibe ngu > 7 mg/L
Ubukho bodaka (NTU)		Inguqu ephakathi	Inguqu ephakathi ivumelekile
Ukutsala umbane (mS/m)	IBakala = A	≤30 mS/m	i-95 th percentile ibe ngu < 30 mS/m
Chl a: periphyton	Azikhho iinkcukacha. Uhlolo	≤12 mg/m ² (IBakala B)	i-50 th percentile ibe ngu < 12 mg/m ²
Chl a: phytoplankton	lolwangeyoKwindla nolwangeyeSilimela. Akukho zimpawu zibonakalayo ze-algae.	≤ 15 µg/L (IBakala B)	i-50 th percentile ibe ngu < 15 µg/L
Izinto eziphilayo ezingenamathambo (ASPT)	A (olu phando)		
Inqaku lemigqeku yeentlanzi	C (olu phando)		
Ukwahluka ngemvakalelo			Ngekota
			Bona ii-Ecospecs zeentlanzi nezezinto eziphilayo ezingenamathambo ngokulandelelanayo

lityhefu	Ukutyhefeka komphakathi womjelo	Azikho iinkcukacha Azikho iinkcukacha. Kungenzeka ukuba zikho izibulala zihambuzane ezithile	Vavanya kuphela xa iziphumo zohlolo lwendalo zibonisa ukuba ikho ingxaki eqatseleyo futhi isizathu singaziwa.
----------	---------------------------------	--	---

UTafle 5.30: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) nee-TPCs kumlambo i-Malgas kwi- WQSU2

UMLAMBO		IINDAWO ZOHLOLO LWEKWALITI YAMANZI			
I-WQSU		RC	Amxabiso angagqibekiyo		
ISIKHUNDLA SE-EWR SITE		PES	Malgas R. at Blanco - K3H004-Q01 ('01 – '06 n = 53)		
Uhlolo lokuba nakho		Uyilungele xa iyonke i-WQSU. Uphantsi xa sithetha ngesikhundla se-EWR.			
Izithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	IBakala/Amagqabantshintshi	G-power (ukuba nakho)
Iityuwa ezingezozendalo (mg/L)	MgSO ₄				Azikho iinkcukacha
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
Izondlo (mg/L)	SRP	≤0.005	0.038	IBakala = C. Isimbo siyanda.	0.116 (Phantsi)
	TIN	≤0.25	0.413	IBakala = A/B. Isimbo siyanda	
Ukwahluka kwendalo	pH (5 th + 95 th %ile)		4.3 + 7.2	Une-asidi ngendalo.	0.84 (Phezulu)
	Ubushushu	16.3 (50%ile)	-	Azikho iinkcukacha ze-PES. Akulindelekanga ngxaki kuba akukho dama phaya kumphezulu womjelo	
	I-oksijini enyibilikisiweyo				
	Ubukho bodaka (NTU)			Azikho iinkcukacha. Ingakho yona ingxaki ngenxa yekwari	
	Ukutsala umbane (mS/m)	<30	15	IBakala = A. Isimbo = senyuka kancinane	0.64 (Phakathi)
Ukwahluka ngokwemvakalelo	Chl a: periphyton			Azikho iinkcukacha	Azikho iinkcukacha (kodwa uhlolo olwenziweyo kwisikhundla se-EWR lubonisa ingxaki).
	Chl a: phytoplankton			Azikho iinkcukacha	
	Ubukhulu bezinto eziphilayo ezingenamathambo			IBakala A ; ASPT = 8.2, SASS = 164 (olu phando)	
	Inqaku lemigqeku yeentlanzi			C/D (olu phando)	
Iityhefu				Azikho iinkcukacha. Kungenzeka kubekho ukutyhefeka ngenxa yemisebenzi yekwari/yesamente/ ye-asphalt, nangenxa yamanzi asuka ezindlwini nalawo asuka kwimisebenzi yezolimo	
LULONKE UHLELO LWESIKHUNDLA		IBakala B (xa sisebenzisa umzekelo we-PAI)			

UTafle 5.31: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo i-Maalgate

UMLAMBO		Umlambo i-Moeras/i-Maalgate	IINDAWO ZOHLOLO LWEKWALITI YAMANZI	
I-WQSU		N/A	RC	litafle zomda ezingagqibekiyo ngomlambo webakala A
ISIKHUNDLA SE-EWR		Moe 1 & Maa 2	PES	Umlambo iMaalgate . @ Noetze Kamma K3H003 ('02 – '06; n = 52)
Uvavanyo lokuba nakho		Kuhle.		
IZithako zeKwaliti yaManzi		Ixabiso	IBakala/Amagqabantshintshi	
Izondlo (mg/L)	SRP	0.019	IBakala = B (TP = 0.015 mg/L)	
	TIN	0.1	IBakala = A	
Ukwahluka kwendalo	pH (5 th – 95 th %)	5.7 – 7.7	Une-asidi ngendalo	
	Ubushushu		Azikho iinkcukacha. Usenokuchaphazeleka (ngakumbi xa kutsalwa amanzi)	
	I-oksijini enyibilikisiweyo		Azikho. Usenokuchaphazeleka (ngakumbi xa amanzi etsalwa ngokugqithisileyo)	
	Ubukho bodaka (NTU)		Azikho iinkcukacha. Utyelelo kwesi sikhundla lubonisile ukuba umlambo awunywanga.	
	Ukutsala umbane (mS/m)	63	IBakala = C	
Ukwahluka ngemvakalelo	Chl a: periphyton		Azikho iinkcukacha. Utyelelo lwesikhundla lubonise i-periphyton ekumaqondo aphezulu	
	Chl a: phytoplankton		Azikho iinkcukacha	
	Ubukhulu bezinto eziphilayo ezingenamathambo (ASPT)		D (uphando lwangoku) Kuhle (RHP)	
	Inqaku lemigqeku yeentlanzi		D (uphando lwangoku)	
lityhefu			Azikho iinkcukacha – kulindeleke ukuba kubekho izibulala zinambuzane ezisuka kwezolimo	
LULONKE UHLELO LWESIKHUNDLA		Ibakala C (ngeokweembono zencutshe)		
Amaxabiso emida				
Isiqinisekisi		Ixabiso lomda osezantsi	Ixabiso lomda ophezulu	
TIN (mg/L) – 50 th %ile			2.0	
PHYTOPLANKTON Chl a (µg/L) – 50 th %ile		15	20	
SRP or PO4 (mg/L) – 50 th %ile			0.058	
PERIPHYTON Chl a (mg/m ²) – 50 th %ile		12	21	
pH – 5 th %ile and 95 th %ile		i-5 th percentile: 5.00 – 5.23		
UBUSHUSHU (°C) – 10 th %ile and 90 th %ile		Lwahluka hayi ngaphezulu kwe- 2°C kuluhlu lwendalo		
I-oksijini enyibilikisiweyo (mg/L) – 5 th %ile		6		
ISALATHISI SENDALO (ASPT)		5.67		

UTafle 5.32: Amabakala e-PES nohlolo lwesikhundla lulonke kumlambo i-Gwaing (i-Malgas/i-Keur) kwi-WQSU1

UMLAMBO	Umlambo i-Malgas (iKeur)	IINDAWO ZOHLOLO LWEKWALITI YAMANZI		
I-WQSU	WQSU 1	RC	Umlambo iRooi @ George K3H002-Q01 ('77 – '79 n = 84)	
ISIKHUNDLA SE-EWR	Asikho	PES	K3H002-Q01 ('01 – '06 n = 65)	
Uhlolo lokuba nakho		Phantsi. Qaphela: akhutshelwe ngaphandle kwendawo yoboniselo ngamanzi ekufuphi		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	Amabakala/Amagqabantshintshi
lityuwa ezingezozendalo (mg/L)	MgSO ₄			Azikho
	Na ₂ SO ₄			
	MgCl ₂			
	CaCl ₂			
	NaCl			
	CaSO ₄			
Izondlo (mg/L)	SRP	0.014	≤0.014*	IBakala = A
	TIN	0.04	≤0.25*	IBakala = A
Ukwahluka kwendalo	pH (5 th – 95 th %)		3.7 – 6.7	Une-asidi ngendalo
	Temperature			Azikho iinkcukacha kodwa kulindeleke ukuba uhlale unjengoko udaliwe kuba uphuhliso olukumphezulu womjelo alukho kangako.
	i-oksijini enyibilikisiweyo			
	Ubukho bodaka (NTU)			
	Ukutsala umbane (mS/m)	16	17	IBakala = A
Ukwahluka ngokwemvakalelo	Chl a: periphyton			Azikho iinkcukacha
	Chl a: phytoplankton			
	Ubukhulu bezinto eziphilayo ezingenamathambo			Kuhle (RHP)
	Inqaku lemigqeku yeentlanzi			kwendalo (RHP)
lityhefu				Azikho iinkcukacha
LULONKE UHLELO LWESIKHUNDLA		IBakala A (ngokweembono zencutshe)		
Amaxabiso emida				
Isiqinisekisi		Ixabiso lomda osezantsi	Ixabiso lomda ophezulu	
TIN (mg/L) – 50 th %ile			≤ 0.25	
PHYTOPLANKTON Chl a (µg/L) – 50 th %ile			<10	
SRP or PO ₄ (mg/L) – 50 th %ile			≤ 0.005	
PERIPHYTON Chl a (mg/m ²) – 50 th %ile			< 1.7	
pH – 5 th %ile and 95 th %ile		6.5 – 8.00		
UBUSHUSHU (°C) – 10 th %ile and 90 th %ile		Uluhlu lobushushu lwendalo		
I-oksijini enyibilikisiweyo (mg/L) - 5 th %ile		>8		
ISALATHISI SENDALO (ASPT)		7		

UTafle 5.33: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo i- Gwaing (i-Malgas) kwi-WQSU2

UMLAMBO	Umlambo iMalgas	IINDAWO ZOHOLO LWEKWALITI YAMANZI			
I-WQSU	WQSU 2	I-RC	Amaxabiso okungagqibeki		
ISIKHUNDLA SE-EWR	Mal 1	I-PES	Umlambo iMalgas . @ Blanco - K3H004-Q01 ('01 – '06 n = 53)		
Uvavanyo lokuba nakho		Kuhle ngayo iyonke i- WQSU. Phantsi ngesikhundla se-EWR.			
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES Value	IBakala/Amagqabantsintshi	G-power (Ukuba nakho)
lityuwa ezingezozendalo (mg/L)	MgSO ₄			Azikho iinkcukacha	
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
Izondlo (mg/L)	SRP	≤0.005	0.038	IBakala = C. isimbo siyanda	0.116 (Phantsi)
	TIN	≤0.25	0.413	IBakala = A/B. Isimbo siyanda	
Ukwahluka kwendalo	pH (5 th – 95 th %)		4.3 – 7.2	Une-asidi yendalo.	0.84 (Phezulu)
	Ubushushu	16.3 (50%ile)	-	Azikho iinkcukacha ze-PES. Akukho ngxaki ilindelekileyo kuba alikho idama kumphezulu womjelo	
	i-oksijini enyibilikisiweyo			Azikho iinkcukacha. Ingakho ingxaki ngenxa yekwari	
	Ubukho bodaka (NTU)			IBakala = A. Isimbo sanda kancinci nje	0.64 (Phakathi)
	Ukutsala umbane (mS/m)	<30	15		
Ukwahlula ngokwemvakalelo	Chl a: periphyton			Azikho iinkcukacha	Azikho iinkcukacha (kodwa uphando lubonisile ukuba ikho ingxaki kule ndawo).
	Chl a: phytoplankton			Azikho iinkcukacha	
	Ubukhulu bezinto eziphilayo ezingenamathambo			IBakala A: ASPT = 8.2, SASS = 164 (olu phando)	
	Inqaku lemigqeku yeentlanzi			C/D (olu phando)	
lityhefu				Azikho iinkcukacha. Kungakho ukutyhefeka ngenxa yemisebenzi yekwari/yesamente/ye-asphalt, amanzi asuka emakhayeni nabuyela emlanjeni.	
LULONKE UHLELO LWESIKHUNDLA		IBakala B (ngokomzekelo we-PAI). Iukhangeleka lunokubangcono kwisikhundla se-EWR uqobo.			
Amaxabiso emida					
Isiqinisekisi		Ixabiso lomda osezantsi		Ixabiso lomda ophezulu	
TIN (mg/L) – 50 th %ile				0.75	
PHYTOPLANKTON Chl a (µg/L) – 50 th %ile		5		10	
SRP or PO ₄ (mg/L) – 50 th %ile				0.02	

PERIPHYTON Chl a (mg/m ²) – 50 th %ile	1.7	12
pH – 5 th %ile and 95 th %ile	I-5 th percentile: 6.00 – 6.24 I-95 th percentile: 8.37 – 8.69	
UBUSHUSHU (°C) – 10 th %ile and 90 th %ile	Uluhu lobushushu bendalo	
I-oksijini enyibilikisiweyo (mg/L) - 5 th %ile	7	
ISALATHISI SENDALO (ASPT)	6.34	

UTafle 5.34: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo i- Gwaing kwi-WQSU3

UMLAMBO	Umlambo i-Gwaing	IINDAWO ZOHLOLO LWEKWALITI YAMANZI		
I-WQSU	WQSU 3	I-RC	Amaxabiso okungagqibeki	
ISIKHUNDLA SE-EWR	Asikho	I-PES	Asikho isitishi sohlolo sikaDWS	
Uvavanyo lokuba nakho		Une-RC ephantsi ngenxa yokusetyenziswa kwamaxabiso okungagqibeki. Une-PES ephakathi kuba kungekho sitishi sohlolo, kodwa ke uxhaswa ziinkcukacha zendalo neenkukacha ezingephi ezongezelelweyo.		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	IBakala /Amagqabantshintshi
Iityuwa ezinezozendalo (mg/L)	MgSO ₄			Azikho iinkcukacha
	Na ₂ SO ₄			
	MgCl ₂			
	CaCl ₂			
	NaCl			
	CaSO ₄			
Izondlo (mg/L)	SRP	≤0.005	1.4 mg/L	Ibakala = F
	TIN	≤0.25		IBakala = D
Ukwahluka kwendalo	pH (5 th – 95 th %)			Azikho iinkcukacha. Kungenzeka ibekho ingxaki ngenxa ye-DO esezantsi nobukho bodaka obukwinqanaba eliphezulu
	Ubushushu			
	i-oksijini enyibilikisiweyo			
	Ubukho bodaka (NTU)			
	Ukutsala umbane (mS/m)	<30	49	
Ukwahluka ngokwemvakalelo	Chl a: periphyton			Azikho iinkcukacha
	Chl a: phytoplankton			
	Ubukhulu bezinto eziphilayo ezingenamathambo (ASPT)			RHP = "amanzi anekwaliti embi"
	Inqaku lemigqeku yeentlanzi			
Iityhefu				Azikho iinkcukacha, kodwa ingakho ingxaki
LULONKE UHLELO LWESIKHUNDLA		IBakala D/E (iimbono zencutshe)		
Amaxabiso emida				
Isiqinisekisi		Ixabiso lomda osezantsi	Ixabiso lomda eliphezulu	
TIN (mg/L) – 50 th %ile			4.0	
PHYTOPLANKTON Chl a (µg/L) – 50 th %ile		20	30	
SRP or PO ₄ (mg/L) – 50 th %ile			0.125	
PERIPHYTON Chl a (mg/m ²) – 50 th %ile		21	84	

pH – 5 th %ile and 95 th %ile	i-5 th percentile: 5.46 – 5.7 i-95 th percentile: 8.56 – 10.00	
UBUSHUSHU (°C) – i-10 th %ile ne-90 th %ile	Awuguquki ngaphezu kwe- 4°C xa uwuthelekisa noluhlu lwendalo	
I-oksijini enyibilikisiweyo (mg/L) - 5 th %ile	4	
ISALATHISI SENDALO (ASPT)	5	

Apho kuye kwafunyanwa umahluko kumaxabiso ekwaliti yamanzi kwiMijelo yokuphilisana yeNdalo, kulapho kuye kwakhethwa ixabiso eliqatha okanye lokhuselo ngecandelo lekwaliti yamanzi laloo mjelo.

UTafale 5.35: amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iKaaimans kwi-WQSU 2

UMLAMBO		Umlambo i-Kaaimans River	IINDAWO ZOHLLOLO LWEKWALITI YAMANZI		
I-WQSU		WQSU 2	RC	Kaaimans R. @ Barbierskraal - K3H001-Q01 ('77 – '81; n = 175)	
ISIKHUNDLA SE-EWR		Ka1	PES	K3H001-Q01 ('01 – '06; n = 56)	
Uvavanyo lokuba nakho		Uphezulu ngokwe-RC ne-PES, kuba isitishi sohlolo sikwi- WQSU, unobude obuvisayo beenkcukacha, kuba iincinci inguqu yomhlaba kwindawo yobonisele ngamanzi.			
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	IBakala/Amagqabantshintshi	G-power (ukuba nakho)
lityuwa ezingezozendalo (mg/L)	MgSO ₄				Azikho iinkcukacha
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
Izondlo (mg/L)	SRP	0.011	0.028*	IBakala = B. isimbo = sande kancinane	0.122 (Phantsi)
	TIN	0.04	0.061	IBakala = A. isimbo sande nje kancinci	
Ukwahluka kwendalo	pH (5 th – 95 th %)		4.4 – 7.4	Amanzi ane-asidi yendalo	0.87 (Phezulu)
	Ubushushu			Azikho iinkcukacha.	
	i-oksijini enyibilikisiweyo			Ayinakude ibange ingxaki le nto	
	Ubukho bodaka (NTU)			Azikho iinkcukacha. Uphando ngokubona = ubukho bodaka kwinqanaba elisezantsi	
	Ukutsala umbane (mS/m)	16	17	IBakala A. isimbo = sizinzile	0.87 (Phezulu)
Ukwahluka ngokwemvakalelo	Chl a: periphyton			Azikho iinkcukacha. Uphando ngokubona = alukho baxekileyo ukhulo lwe- algal	
	Chl a: phytoplankton				
	Ubukhulu bezinto eziphilayo ezingenamathambo			IBakala A; ASPT = 8.0, SASS = 175 (olu phando). kwendalo (RHP)	
	Inqaku lemigqeku yeentlanzi			B	
lityhefu				Azikho iinkcukacha, kodwa ayinakude ibe yingxaki le nto	
LULONKE UHLELO LWESIKHUNDLA		IBakala A (lithathelwe kumzekelo we-PAI)			

UTafle 5.36: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) kumlambo iKaamans River kwi- K30C

IZithako zeKwaliti yamanzi	umda	IiMfuno zoMjelo weNdalo	UMqathango o weeMfuno ezisisiseko zoLuntu ⁵	UMqathango woMjelo: ikwaliti yamanzi
Ikhemistri ngokuthe gabalala-iityuwa ezingezizendalo eziphambili	MgSO ₄ (mg/l) ¹	< 16	N/A	< 16
	Na ₂ SO ₄ (mg/l) ¹	< 20	N/A	< 20
	MgCl ₂ (mg/l) ¹	< 15	N/A	< 15
	CaCl ₂ (mg/l) ¹	< 21	N/A	< 21
	NaCl (mg/l) ¹	< 45	N/A	< 45
Ikhemistri ngokuthe gabalala – ii-ioni eziphambili	Sodium (mg/l)	N/A	<200	<200
	Magnesium (mg/l)	N/A	<100	<100
	Chloride (mg/l)	N/A	<200	<200
	Calcium (mg/l)	N/A	<80	<80
	Sulphate (mg/l)	N/A	<400	<400
Izondlo	Phosphate (PO ₄) (mg/l) ²	<0.02mg/L	N/A	<0.02mg/L
	Iyonke i- Inorganic Nitrogen (mgN/l) ²	<0.25mg/L	N/A	<0.25mg/L
Ikwality ebonakalyo yamanzi	pH (range) i-5 th percentile i-95 th percentile	6.7 7.4	5 9.5	5 7.4
	i-oksijini enyibilikisiweyo (mg/l)¹	>1.7 mg/L	N/A	>1.7 mg/L
	Ubushushu¹	Incinci inguqu xa ubuthelekisa nobendalo		Incinci inguqu xa ubuthelekisa a nobendalo
	Ukutsala umbane (mS/m) – WUSEBENZISE KUPHELA XA IITYUWA EZIZIZIGAQA ZIVELISIWE	≤30mS/m	0-70	≤30mS/m
Ikwality yamanzi yendalo	Chl-a: periphyton ³	<1.7 mg/m ²	N/A	
	Chl-a: phytoplankton³	< 10 µg/L (A category)	N/A	
	Ukwakheka komgqeku we-Biotic – Ubukhuly bezinto eziphilayo ezingenamathambo s	ASPT: 8 (A category)		
	Ukutyhefeka ngaphakathi komjelo	Ukutyhefeka ngaphakathi		

IZithako zeKwaliti yamanzi	umda	IiMfuno zoMjelo weNdalo	UMqathango weMfuno ezisisiseko zoLuntu ⁵	UMqathango woMjelo: ikwaliti yamanzi
		komjelo makungenzeki		
lityhefu nemixube entsonkothileyo	lityhefu (njengoko kudwelisiwe ku DWAF, 1996 ⁶)	≤ TWQR	≤ TWQR	≤ TWQR

PHAWULA:

¹: Ukuthobela imiqathango yi-95th percentile. ²: ukuthobela imiqathango yi-50th percentile. ³ i-Chl-a ayibandakanyeki kwizifundo ngemijelo zekhompyutha. ⁴ ukuthobela imiqathango kukwinqanaba le-90th percentile

⁵ ref: *South African Water Quality Guidelines, Volume 1: Domestic Water Use, 2nd Ed.* 1996. Department of Water Affairs and Forestry. Pretoria, South Africa.

⁶ ref: *South African Water Quality Guidelines, Volume 7: Aquatic Ecosystems, 2nd Ed.* 1996. Department of Water Affairs and Forestry. Pretoria, South Africa.

UTafale 5.37: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iDiep River kwi- WQSU 3

UMLAMBO		IINDAWO ZOHLOLO LWEKWALITI YAMANZI		
I- WQSU	Umlambo iDiep River WQSU 3 (Kuqala inqanaba II le-Ecoregion 20.02 ukuya ekuqaleni komda womjelo wasezintabeni)	I-RC	Umlambo iDiep R. @ uMjelo weHlathi i-Woodville K4H003Q01 ('77 – '80; n = 58)	
ISIKHUNDLA SE-EWR	3	I-PES	K4H003Q01 ('03 – '07; n = 36)	
Uvavanyo lokuba nakho		Phakathi ukuya Phezulu		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	IBakala /Amagqabantsintshi
Iityuwa ezingezozendalo (mg/L)	MgSO ₄	8.86	15.87	U-TEACHA usetyenziselwe ukuqokelela iinkcukacha zohlolo. Iityuwa = iBakala A
	Na ₂ SO ₄	0.00	0.00	
	MgCl ₂	14.83	14.52	
	CaCl ₂	7.18	9.89	
	NaCl	102.10*	94.31	
Izondlo (mg/L)	SRP	0.003	0.018	IBakala B/C.
	TIN	0.04	0.07	IBakala A.
Ukwahluka kwendalo	pH (5 th + 95 th %ile)	4.8 + 6.2	4.8 + 7.5	IBakala B.
	Ubushushu	-	-	Azikho iinkcukacha kodwa akukho futhe lilindelekileyo.
	I-oksijini enyibilikisiweyo	-	-	Azikho iinkcukacha. Kubonakala kancinci imizila yeentlenge.
	Ubukho bo (NTU)	-	-	IBakala A
Ukwahluka ngokwemvakalelo	Ukutsala umbane (mS/m)	-	18.0	IBakala A
	Chl a: periphyton	-	21.25	Kukho ukwenyuka okuthile kwezondlo okuboniswa ziinkcukacha ze- periphyton data (IBakala C/D category; n=1)
	Chl a: phytoplankton	-	0.18	IBakala B lesimo sangoku.
	Ubukhulu bezinto eziphilayo ezingenamathambo	ASPT [§] : ubuncikane be- 6.58	ASPT: 7.3 MIRAI ^{**} : 86.1%	IBakala B lesimo sangoku.
	Inqaku lemigqeku yeentlanzi.	-	FRAI [*] : 86.1%	IBakala B lesimo sangoku.
Iityhefu	ii-Diatoms	-	SPI [#] =17.6	Ikwali ephezulu yamanzi
		-	-	Azikho iinkcukacha, kodwa ke likho ifuthe elilindelekileyo ngenxa yezibulala zinambuzane ezisetyenziswa ezifama nezichumisi
LULONKE UHLELO LWESIKHUNDLA		B (Ngokomzekelo we-PAI)		

*: ixabiso lomda lihlangahlengisiwe

sovavanyo

-: Azikho iinkcukacha

*: FRAI = Fish Response Assessment Index (Isalathisi lokuvakalelwa kweentlanzi)

**: MIRAI = Macro Invertebrate Response Assessment Index (Isalathisi sovavanyo lokuvakalelwa kobukhulu bezinto eziphilayo ezingenamathambo)

#: Specific Pollution Index (Isalathisi soqobo lokungcola)[§]: ASPT = inqaku le-avareji ngokwe-Taxon

UTafale 5.38: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) kumlambo iDiep River (kwi-WQSU 3, K40A)

Umlambo: iDiep		Isikhundla se-EWR: 3	Isikhundla sohlolo: K4H003Q01
Ii- metrics zekwaliti yamanzi		i-ECOSPEC	
Iityuwa ezingezozendalo*	MgSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 16 mg/L.	
	Na ₂ SO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 20 mg/L.	
	MgCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 15 mg/L.	
	CaCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 21 mg/L.	
	NaCl	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 191 mg/L.	
	CaSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 351 mg/L.	
Ukwahluka kwendalo	EC	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 30 mS/m.	
	pH	i-5 th ne-95 th percentiles yeenkcukacha mayibe phakathi ko-4.5 no- 7.5.	
	Ubushushu	Kukho ukuphambuka okuncinane kuluhlu lobushushu bendalo	
	i-oksijini enyibilikisiweyo	i-5 th percentile yeenkcukacha mayibe ngu ≥ 7.5 mg/L.	
	Ubukho bodaka	Vary by a small amount from the natural turbidity range; minor silting of instream habitats acceptable.	
Izondlo	TIN	i- 50 th percentile yeenkcukacha mayibe ngu ≤ 0.25 mg/L.	
	PO ₄ -P	i- 50 th percentile yeenkcukacha mayibe ngu ≤ 0.025 mg/L.	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i- 50 th percentile yeenkcukacha mayibe ngu <15 µg/L.	
	Chl-a periphyton	i- 50 th percentile yeenkcukacha mayibe ngu ≤ 52.5 mg/m ² .**	
Iityhefu		i-95 th percentile yeenkcukacha mayibe phakathi koLuhlu lweKwaliti yaManzi eNgqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe kuDWAf (1996).	

* Mayiveliswe ngokusebenzisa i-TEACHA xa i-TPC ye- EC ilindelekile okanye uncoliso lwetyuwa lulindelekile.

** i-Periphyton (21.25 mg/m²) kanye-kanye ikwiBakala C/D (C = 12 - 21 no- D = 21 - 84 mg/m²; DWAf, 2008), ngoko ke umda ophezulu kaC/D uchazwe njenge- EcoSpec.

UTafale 5.39: ii-TPC zekwaliti yamanzi kumlambo iDiep River (kwi-WQSU 3, K40A)

Umlambo: iDiep		ISIKHUNDLA SE-EWR: 3	Isikhundla sohlolo: K4H003Q01
Ii- metrics zekwaliti yamanzi		I-TPC	
Iityuwa ezingezozendalo*	MgSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu 13 – 16 mg/L.	
	Na ₂ SO ₄	i- 95 th percentile yeenkcukacha mayibe ngu 16 – 20 mg/L.	
	MgCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu 12 – 15 mg/L.	
	CaCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu 17 – 21 mg/L.	
	NaCl	i- 95 th percentile yeenkcukacha mayibe ngu 36 – 45 mg/L.	
	CaSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu 153 – 191 mg/L.	
Ukwahluka kwendalo	EC	i-95 th percentile yeenkcukacha mayibe ngu 24 – 30 mS/m.	
	pH	i- 5 th ne-95 th percentiles yeenkcukacha mayibe ngu <4.7 and >7.3.	
	Ubushushu	Luncinane ukuya phakathi uphambuko xa uthelakisa noluhlu lobushushu S bendalo. Ikho loo migqeku ingamininzanga nengaxhaphakanga kuyaphi kunoko kulindelekileyo ngokofuniselo.	
	i-oksijini enyibilikisiweyo	i- 5 th percentile yeenkcukacha mayibe ngu 7.8 – 7.5 mg/L.	
	Ubukho bodaka	Zikho iinguqu eziphakathi kusetyenziso lomhlaba wendawo yoboniselo ngamanzi, nezithi zibangele (okwexeshana nje) iingqumba zeentlenga ezikwinqanaba eliphezulu ngokuqatseleyo nobukho bodaka obukwinqanaba eliphezulu.	

Umlambo: i-Diep		ISIKHUNDLA SE-EWR: 3	Isikhundla sohlolo: K4H003Q01
Ii- metrics zekwaliti yamanzi		I-TPC	
Izondlo	TIN	i-50 th percentile yeenkcukacha mayibe ngu 0.2 – 0.25 mg/L.	
	PO ₄ -P	i-50 th percentile yeenkcukacha mayibe ngu 0.02 – 0.025 mg/L.	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i- 50 th percentile yeenkcukacha mayibe ngu 12 – 15 µg/L.	
	Chl-a periphyton	i- 50 th percentile yeenkcukacha mayibe ngu 42 – 52.5 mg/m ² .	
Iityhefu	i95 th percentile yeenkcukacha mayibe phakathi koLuhlu lweKwaliti yaManzi eNgqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe ku DWAF (1996).		

UTafle 5.40: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iKarataru kwi-WQSU 5

UMLAMBO	Umlambo iKarataru	IINDAWO ZOHLOLO LWEKVALITI YAMANZI			
I-WQSU	WQSU 5 (endaweni yemveli ukuya eSwartvlei)	I-RC	iKarataru R. @ kuMjelo weHlathi iKarataru K4H002Q01 ('76 – '79; n = 115)		
ISIKHUNDLA SE-EWR	4	I-PES	K4H002Q01 ('03 – '07; n = 36)		
Uvavanyo lokuba nakho	Phakathi, kuba zanele iinkcukacha zovavanyo lentsukaphi nezesimo sangoku				
Izithako zeKwaliti yaManzi	Ixabiso le-RC	Ixabiso le-PES	IBakala/Amagqabantshintshi		
Iityuwa ezingezizo zendalo (mg/L)	MgSO ₄	7.62	12.63	U-TEACHA uye wasetyenziswas kuvavanyo lweenkcukacha used for data assessment. Iityuwa = ibakala A	
	Na ₂ SO ₄	2.05	3.01		
	MgCl ₂	4.43	4.16		
	CaCl ₂	9.89	9.16		
	NaCl	35.59	36.15		
Izondlo (mg/L)	CaSO ₄	0.73	0.73		
	SRP	0.022 *	0.047	IBakala B.	
Ukwahluka kwendalo	TIN	0.07	0.128	IBakala A.	
	pH (5 th + 95 th %ile)	3.7 + 7.7	3.8 + 6.6	IBakala A.	
	Ubushushu	-	-	Azikho iinkcukacha kodwa ke akukho futhe lilindelekileyo	
	i-oksijini enyibilikisiweyo	-	-	Azikho iinkcukacha. Buncinane ubungqina bobukho beentlenge.	
	Ubukho bodaka (NTU)	-	-	iBakala A.	
Ukwahluka ngokwemvakalelo	Ukutsala umbane (mS/m)	-	7.0		
	Chl a: periphyton	-	9.91	IBakala B nge-periphyton.	
	Chl a: phytoplankton	-	0.09		
	Ubukhulu bezinto eziphilayo ezingenamathambo	ASPT: 7.3	ASPT: 8.1 MIRAI: 92.3%	IBakala A/B ngesimo sangoku.	
	Inqaku lemigqeku yeentlanzi	-	FRAI: 82.4%	IBakala B ngesimo sangoku.	
Iityhefu	ii-Diatoms	-	SPI=19.9	Amanzi anekwaliti epehezulu	
		-	-	Azikho iinkcukacha, kodwa likho elo futhe libangelwa kukwenziwa kwamaplangubut e-Geelhoutvlei.	
LULONKE UHLELO LWESIKHUNDLA		ngUB ngeWQSU 5 , nangona ilibakala A ekunabeni komlambo womjelo osemantla (umzekelo we-PAI)			

*: Ixabiso lomda lilihlengahlengisiwe

-: Azikho iinkcukacha

Table 5.41: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) kumlambo iKaratarata kwi-WQSU

5

River: Karatarata		EWR Site: 4	Monitoring site: K4H002Q01
Water quality metrics		ECOSPEC	
lityuwa ezingezozendalo*	MgSO ₄	i-95 th percentile yeenkcukacha mayibe ngu ≤ 16 mg/L.	
	Na ₂ SO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 20 mg/L.	
	MgCl ₂	i-95 th percentile yeenkcukacha mayibe ngu ≤ 15 mg/L.	
	CaCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 21 mg/L.	
	NaCl	i-95 th percentile yeenkcukacha mayibe ngu ≤ 191 mg/L.	
	CaSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 351 mg/L.	
Ukwahluka kwendalo	EC	i-95 th percentile yeenkcukacha mayibe ngu ≤ 30 mS/m.	
	pH	i-5 th ne- 95 th percentiles yeenkcukacha mayibe phakathi ko- 4.5 no- 7.5.	
	Ubushushu	Uluhlu lobushushu bendalo.	
	i-oksijini enyibilikisiweyo	i-5 th percentile yeenkcukacha mayibe ngu ≥ 8.0 mg/L.	
	Ubukho bodaka	Vary by a small amount from the natural turbidity range; minor silting of instream habitats acceptable.	
Izondlo	TIN	i-50 th percentile yeenkcukacha mayibe ngu ≤ 0.25 mg/L.	
	PO ₄ -P	i-50 th percentile yeenkcukacha mayibe ngu ≤ 0.075 mg/L.	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i-50 th percentile yeenkcukacha mayibe ngu < 10 µg/L.	
	Chl-a periphyton	i-50 th percentile yeenkcukacha mayibe ngu ≤ 12 mg/m ² .	
lityhefu		i-95 th percentile yeenkcukacha mayibe phakathi koluhlu lwekwaliti yamanzi engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe ku DWAF (1996).	

* ziza kufunyanwa ngokusebenzisa u-TEACHA xa ne- TPC ye-EC ibethe ngaphaya okanye ungcoliswe lweentyuwa lulindelekile.

UTafale 5.42: Ii-TPCs zekwaliti yamanzi kumlambo iKaratarata kwi- WQSU 5

Umlambo: iKaratarata		Isikhundla se-EWR: 4	Isikhundla sohlolo: K4H002Q01
Ii-metrics zekwaliti yamanzi		I-TPC	
lityuwa ezingezizo zendalo*	MgSO ₄	i-95 th percentile yeenkcukacha mayibe ngu 13 – 16 mg/L.	
	Na ₂ SO ₄	i- 95 th percentile yeenkcukacha mayibe ngu 16 – 20 mg/L.	
	MgCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu 12 – 15 mg/L.	
	CaCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu 17 – 21 mg/L.	
	NaCl	i- 95 th percentile yeenkcukacha mayibe ngu 36 – 45 mg/L.	
	CaSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu 153 – 191 mg/L.	
Ukwahluka kwendalo	EC	i-95 th percentile yeenkcukacha mayibe ngu 24 – 30 mS/m.	
	pH	i- 5 th and 95 th percentiles yeenkcukacha mayibe ngu <4.7 and >7.3.	
	Ubushushu	Kukho ukuphambuka kancinci kolwala luhlu lobushushu bendalo.	
	i-oksijini enyibilikisiwe	i- 5 th percentile yeenkcukacha mayibe ngu 8.2 – 8.0 mg/L.	
	Ubukho bodaka	Zincinci ukuya kweziphakathi iinguqu kusetyenziso lomhlaba kwindawo yoboniselwe ngamanzi, nto leyo ibangela ifuthe elincinci nelilelexeshana nje leentlenga zeeendawo zokuphilisana.	
Izondlo	TIN	i- 50 th percentile yeenkcukacha mayibe ngu 0.2 – 0.25 mg/L.	
	PO ₄ -P	i- 50 th percentile yeenkcukacha mayibe ngu 0.06 – 0.075 mg/L.**	

Umlambo: iKarata		Isikhundla se-EWR: 4	Isikhundla sohlolo: K4H002Q01
Ii-metrics zekwaliti yamanzi		I-TPC	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i- 50 th percentile yeenkcukacha mayibe ngu 8 – 10 µg/L.	
	Chl-a periphyton	i- 50 th percentile yeenkcukacha mayibe ngu 10 – 12 mg/m ² .	
lityhefu		i-95 th percentile yeenkcukacha mayibe phakathi koluhlu lwekwaliti yamanzi engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe ku DWAF (1996).	

* Ziza kufunyanwa ngokusebenzisa u-TEACHA xa ne-TPC ye-EC ibethe ngaphaya okanye ungcolisio lomoya lulindelekile .

** nangona umda ophezulu we le-phosphate elichanileyo ungu-0.125 mg/L, i- TPC yona yenzelwe ku-0.075 mg/L njengoko amanqanaba e-PES ye- phosphate ebengu- 0.046 mg/L.

UTafile 5.43: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iHoëkraal kwi- WQSU 4

UMLAMBO		IINDAWO ZOHLOLO LWEKWALITI YAMANZI		
I-WQSU	I-WQSU 4 (ukusuka nokuya kutsho ekuqaleni kweenduli zeenyawo ezisemazantsi)	I-RC	Hoëkraal R. @ Eastbrook K4H001Q01 ('77 – '80; n = 83)	
ISIKHUNDLA SE-EWR	-	I-PES	K4H001Q01 ('03 – '07; n = 28)	
Uvavanyo lokuba nakho		Phantsi - Phakathi njengoko ziincinci nenkcukacha kuhlobo lwesimo sangoku. Kwindawo esemazantsi kumda owongezelelweyo ukusuka kwichibi iSwartvlei. Qaphela ukuba udonga olunqamlezileyo lokuthatha umlinganiselo lumi kumazantsi omlambo i- Hoëkraal.		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	IBakala/Amagqabantshintshi
Iityuwa ezingezizo zendalo (mg/L)	MgSO ₄	35.87	153.20	U-TEACHA usetyenziselwe ukuhlalutya iinkcukacha ze-WMS. Iziphumo mazingahoywa kuba iinkcukacha zibonisa ukungenelela kweetyuwa kwichibi lokukhupha iityuwa emanzini iSwartvlei.
	Na ₂ SO ₄	9.95	159.5	
	MgCl ₂	35.92	100.33	
	CaCl ₂	24.80	45.24	
	NaCl	399.69	1560.41	
	CaSO ₄	0.59	0.73	
Izondlo (mg/L)	SRP	0.014 *	0.034	IBakala B.
	TIN	0.06	0.088	IBakala A.
Ukwahluka kwendalo	pH (5 th + 95 th %ile)	4.4 + 7.2	4.5 + 7.8	IBakala A.
	Ubushushu	-	-	Azikho iinkcukacha kodwa alikho ifuthe elilindelekileyo, nangona nje lomlambo unzulu futhi ubanzi ngaphezulu kwesikhundla esisemazantsi, nto leyo inokubangela ukuhla kwamanqanaba e-oksijini.
	i-oksijini enyibilikisiweyo	-	-	
	Ubukho bodaka (NTU)	-	-	Azikho iinkcukacha kodwa bukho obo bungqina bobukho beentlenge.
Ukutsala umbane (mS/m)		4.2: isikhundla esiphezulu	IBakala A	
Ukwahluka ngokwemvakalelo	Chl a: periphyton	-	4.81: isikhundla esiphezulu 152.93: isikhundla esisemazantsi	Isikhundla esisemazantsi sibonisa izindlo ezakheka ukwenyuka emanzini.
	Chl a: phytoplankton	-	0.14: isikhundla esiphezulu 0.47: isikhundla esisemazantsi	
	Ubukhulu bezilwanyana ezingenamathambo	-	-	Khangela kuvavanywe kuba esi ayisosikhundla se- EWR.
	Inqaku lemigqeku yeentlanzi	-	-	
	ii-Diatoms	-	SPI=19.8: Umlambo ophezulu. SPI=16.2: umlambo osezantsi	Isikhundla esiphezulu: Amanzi anekwaliti ephezulu. Isikhundla esisemazantsi: amanzi anekwaliti entle.
iityhefu		-	-	Azikho iinkcukacha, kodwa ke akukho futhe lilindelekileyo.
LULONKE UHLELO LWESIKHUNDLA		IBakala C Category (umzekelo we-PAI)		

* Ixabiso lomda lihlengahlengisiwe

Table 5.44: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) kumlambo i-Hoëkraal kwi-WQSU

Umlambo : iHoëkraal		Isikhundla se-EWR: -	Isikhundla sohlolo: K4H001Q01, nangona simi kwindawo esemazantsi yomlambo futhi nokungenelela kwamanzi anetyuwa kuyarhaneleka ngenxa yenkxaso yechibi i- Swartvlei.
Ii-metrics zekwaliti yamanzi		ECOSPEC	
Iityuwa ezingezozendalo*	MgSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 16 mg/L.	
	Na ₂ SO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 20 mg/L.	
	MgCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 15 mg/L.	
	CaCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 21 mg/L.	
	NaCl	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 191 mg/L.	
	CaSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 351 mg/L.	
Ukwahluka kwendalo	EC	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 30 mS/m.	
	pH	i- 5 th ne-95 th percentiles yeenkcukacha mayibe phakathi ko- 4.5 no- 7.5.	
	Ubushushu	Kukho ukuphambuka kancinci kolwaa luhlu lobushushu bendalo.	
	i-oksijini enyibilikisiweyo	i- 5 th percentile yeenkcukacha mayibe ngu ≥ 7.5 mg/L.	
Izondlo	Ubukho bodaka	Uguquka nje kancinci kolwa luhlu lobukho bodaka bendalo; okwa kubanentlengana kweendawo zokuphilisana zangaphakathi komjelo kwamkelekile.	
	TIN	i-50 th percentile yeenkcukacha mayibe ngu ≤ 0.25 mg/L.	
Ukwahluka ngokwemvakalelo	PO ₄ -P	i- 50 th percentile yeenkcukacha mayibe ngu ≤ 0.125 mg/L.	
	Chl-a phytoplankton	i-50 th percentile yeenkcukacha mayibe ngu yeenkcukacha mayibe ngu <10 µg/L.	
Iityhefu	Chl-a periphyton	i- 50 th percentile yeenkcukacha mayibe ngu ≤ 12 mg/m ² .	
	i-95 th percentile yeenkcukacha mayibe phakathi koluhlu lwekwaliti yamanzi engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubaliwe ku DWAf (1996).		

* ziza kufunyanwa ngokusebenzisa u-TEACHA xa ne-TPC ye- EC ibetha ngaphaya nongcoliso lweyutuwa lulindelekile.

UTafale 5.45: Ii- TPCs zekwaliti yamanzi kumlambo i-Hoëkraal kwi- WQSU 4

Umlambo: iHoëkraal		Isikhundla se-EWR: -	Isikhundla sohlolo: K4H001Q01
Ii- metrics zekwaliti yamanzi		i-TPC	
Iityuwa ezingezozendalo*	MgSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu- 13 – 16 mg/L.	
	Na ₂ SO ₄	i- 95 th percentile yeenkcukacha mayibe ngu- 16 – 20 mg/L.	
	MgCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu- 12 – 15 mg/L.	
	CaCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu- 17 – 21 mg/L.	
	NaCl	i- 95 th percentile yeenkcukacha mayibe ngu- 36 – 45 mg/L.	
	CaSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu- 153 – 191 mg/L.	
Ukwahluka kwendalo	EC	i- 95 th yeenkcukacha mayibe ngu- 24 – 30 mS/m.	
	pH	i-5 th ne- 95 th yeenkcukacha mayibe ngu <4.7 no>7.3.	
	Ubushushu	Ziphantsi ukuya phakathi iinguqu zosetyenziso lomhlaba kwindawo yobonisele ngamanzi, nto leyo ibangela ifuthe elincinci nelilexeshana lokuba neentlengo kweendawo zokuphilisana.	
	i-oksijini enyibilikisiweyo	i- 5 th percentile yeenkcukacha mayibe ngu- 7.8 – 7.5 mg/L.	
Ubukho bodaka	Ziphakathi iinguqu zosetyenziso lomhlaba kwindawo yobonisele ngamanzi, nto leyo ibangela iingqumba zexeshana zeentlengo ezikwiqondo eliphezulu ngokungaqhelekanga nobukho bodaka obukumanganaba aphezulu.		

Umlambo: iHoekraal		Isikhundla se-EWR: -	Isikhundla sohlolo: K4H001Q01
li- metrics zekwaliti yamanzi		i-TPC	
Izondlo	TIN	i- 50 th percentile yeenkcukacha mayibe ngu- 0.2 – 0.25 mg/L.	
	PO ₄ -P	i- 50 th percentile yeenkcukacha mayibe ngu- 0.1 – 0.125 mg/L.	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i- 50 th percentile yeenkcukacha mayibe ngu- 8 – 10 µg/L.	
	Chl-a periphyton	i- 50 th percentile yeenkcukacha mayibe ngu- 10 – 12 mg/m ² .	
Iityhefu		i- 95 th percentile yeenkcukacha mayibe phakathi koluhlu lwekwaliti yamanzi engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe ku DWAF (1996).	

UTafle 5.46: Amabakala ePES novavanyo lwesikhundla lulonke kumlambo iGoukamma kwi- WQSU 1

UMLAMBO	Umlambo iGoukamma	IINDAWO ZOHLOLO LWEKVALITI YAMANZI		
I-WQSU	WQSU 1 (kwintsukaphi ukuya esiphelweni sommandla wentaba)	RC	iRooi R. @ George K3H002-Q01 ('77 – '79 n = 84)	
ISIKHUNDLA SE-EWR	Asikho	PES	K3H002-Q01 ('01 – '06 n = 65)	
Uvavanyo lokuba nakho	Phantsi ukuya Phakathi kuba asikho isitishi sohlolo sakwaDWAF. Kodwa ke indawo yobonisele ngamanzi ayichaphazeleki futhi ke uza kuse ufane nje neminye imijelo yeentaba.			
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	IBakala/Amagqabantshintshi
Iityuwa ezingezizo zendalo (mg/L)	MgSO ₄			Azikho iinkcukacha
	Na ₂ SO ₄			
	MgCl ₂			
	CaCl ₂			
	NaCl			
Izondlo (mg/L)	SRP	0.014	≤0.014*	IBakala A.
	TIN	0.04	≤0.25*	IBakala A.
Ukwahluka kwendalo	pH (5 th – 95 th %)		3.7 – 6.7	Une-asidi ngendalo.
	Ubushushu			Azikho iinkcukacha kodwa ke bulindeleke ukuba bubebobendalo kuba luncinci uphuhliso olwenzekayo phaya kumphezulu womjelo wendawo yobonisele ngamanzi.
	i-oksijini enyibilikisiweyo			
	Ubukho bodaka (NTU)			
	Ukutsala umbane (mS/m)	16	17	IBakala A.
Ukwahluka ngokwemvakalelo	Chl a: periphyton			Azikho iinkcukacha
	Chl a: phytoplankton			Azikho iinkcukacha
	Ubukhulu bezinto eziphilayo ezingenamathambo			Azikho iinkcukacha
	Inqaku lemigqeku yeentlanzi			Azikho iinkcukacha
Iityhefu				Azikho iinkcukacha.
LULONKE UHLELO LWESIKHUNDLA		A		

uTafite 5.47: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) kumlambo iGoukamma kwi- WQSU 2

UMLAMBO		Umlambo iGoukamma	IINDAWO ZOHLOLO LWEKWALITI YAMANZI				
I-WQSU		2	DWAF WQ WMS	Ayikho			
ISIKHUNDLA SE-EWR SITE		Asikho	RHP	Okwangoku kukho isikhundla se-RHP phezulu phaya kwindawo yobonisele ngamanzi			
Ukukholelwa kuvavanyo lwe-PES							
Iphantsi kuba ukhutshelwe ngaphandle kwenye indawo yobonisele ngamanzi (iKarata R) ne-WQSU (Goukamma WQSU 2). Iinkcukacha zendalo (ukusuka kwiprojekthi yangoku ne-RHP yomphezulu wendawo yobonisele ngamanzi) zixhasa i-PES eqikelelwayo yekwaliti yamanzi.							
IZithako zekwaliti yamanzi		IBakala lePES	ii-Eco specs zeWQ	Ingaba ikho inkqubela efunekayo?	I-TPC	Ixesha elimiyo lohlolo	
Iityuwa ezingezozendalo (mg/L)	MgSO ₄		≤16 mg/L		i-95 th percentile ibe ngu < 16 mg/L	Rhoqo ngeenyanga ezimbini	
	Na ₂ SO ₄		≤20 mg/L		i-95 th percentile ibe ngu < 20 mg/L		
	MgCl ₂	Azikho	≤15 mg/L	N/A	i-95 th percentile ibe ngu < 15 mg/L		
	CaCl ₂		≤21 mg/L		i-95 th percentile ibe ngu < 21 mg/L		
	NaCl		≤45 mg/L		i-95 th percentile ibe ngu < 45 mg/L		
Izondlo (mg/L)	SRP	A	≤0.012 mg/L	Yes - B/C	i-50 th percentile ibe ngu < 0.025 mg/L	Rhoqo ngeenyanga ezimbini	
	TIN	A	≤0.25 mg/L	No	i-50 th percentile ibe ngu < 0.25 mg/L	Rhoqo ngeenyanga ezimbini	
Ukwahluka kwendalo	pH	Umlambo one-asidi ngendalo	< 6.4	No	i-95 th percentile ibe ngu < 6.4	Rhoqo ngeenyanga ezimbini	
	Ubushushu	Azikho iinkcukacha. Uhlolo olwenziweyo alubonisanga ngaki yabukho bodaka.	Uluhlu lwendalo	N/A	Uluhlu lobushushu bendalo	Rhoqo ngeenyanga ezimbini	
	i-oksijini enyibilikisiweyo		8 mg/L	N/A	i-5 th percentile ibe ngu > 8 mg/L	Rhoqo ngeenyanga ezimbini	
	Ubukho bodaka (NTU)		Alukho utshintsho	N/A	Ayikho inguqu evumelekileyo	Rhoqo ngeenyanga ezimbini	
	Ukutsala umbane (mS/m)		A – Iizinzile	≤30 mS/m	No	i-95 th percentile ibe ngu < 30 mS/m	Rhoqo ngeenyanga ezimbini
	Chl a: periphyton		Azikho iinkcukacha. Uhlolo olwenziweyo alubonisanga ngaki yabukho bodaka.	≤ 1.7 mg/m ² (Ibakala A) ≤ 10 µg/L (Ibakala B)	N/A	i-50 th percentile ibe ngu < 1.7 mg/m ²	Ngekota
Ukwahluka ngokwemvaka lelo	Ubukhulu bezinto eziphilayo ezingenamathambo (ASPT)	B (olu phando)					
Bona ii-Ecospecs zeentlanzi nezezinto eziphilayo ezingenamathambo ngokulandelelanayo.							

lityhefu	Inqaku lemigqeku yeentlanzi Ukutyhefeka komphakathi womjelo	C (olu phando) Azikho iinkcukacha Azikho iinkcukacha.	Ayinakubakho ingxaki. Vavanya kuphela xa neziphumo zohlolo lwendalo zibonisa ukuba ikho ingxaki eqatseleyo futhi isizathu sayo asaziwa.

UTafle 5.48: Amabakala e- PES novavanyo lwesikhundla lulonke kumlambo iGoukamma kwi-WQSU 2

UMLAMBO		IINDAWO ZOHLOLO LWEKWALITI YAMANZI			
I-WQSU	I-WQSU 2 (Encamini yommandla weentaba namahlathi ukuya esiphelweni esiphezulu sefutha lamaza – kanye phaya ngakwibhulorho u- N2)	I-RC	Karataro R. @ Ihlathi likaRhulumente iKarataro K4H002-Q01 ('71 – '76; n = 107)		
ISIKHUNDLA SE-EWR	Gou 1	I-PES	K4H002-Q01 ('01 – '06; n = 51)		
Uvavanyo lokuba nakho	Kwinqanaba eliphantsi. Isitishi sohlolo yindawo yoboniselo ngamanzi ekufuphi.				
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	Ibakala/Amagqabantshintshi	G-power (ukuba nakho)
lityuwa ezingezozendalo (mg/L)	MgSO ₄				Azikho iinkcukacha
	Na ₂ SO ₄				
	MgCl ₂				
	CaCl ₂				
	NaCl				
	CaSO ₄				
izondlo (mg/L)	SRP	0.018	0.048*	Ibakala = C. isimbo = siyanda	0.232 (phantsi)
	TIN	0.11	0.11	ibakala = A. isimbo = sizinzile	
Ukwahluka kwendalo	pH (5 th – 95 th %)		3.8 – 6.4	Amanzi ane-asidi ngendalo	0.781 (phakathi)
	Ubushushu			Azikho iinkcukacha. Ayinakubakho ingxaki	
	i-oksijini enyibilikisiweyo			Azikhjo iinkcukacha. Uhlolo ngokubona = ubukho bodaka busezantsi	
	Ubukho bodaka (NTU)				
	Ukutsala umbane (mS/m)	10	12	Ibakala = A. isimbo = sizinzile	0.997 (phezulu)
Ukwahluka kwemvakalelo	Chl a: periphyton			Azikho iinkcukacha. Uhlolo ngokubona = alukho ukhulo olubaxekileyo lwe- algae	
	Chl a: phytoplankton				
	Ubukhulu bezinto ezingenamathambo			Ibakala B; ASPT = 7.2, SASS = 143 (uphando lwangoko). Lwendalo (RHP)	
	Inqaku lemigqeku yeentlanzi			C (olu phando)	
lityhefu				Azikho iinkcukacha, kodwa ayinakubakho ingxaki	
LULONKE UHLELO LWESIKHUNDLA		Ibakala A (umzekelo wePAI)			

UTafle 5.49: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) kumiambo iGoukamma kwi-WQSU 2

UMLAMBO		Umlambo iGoukamma	IINDAWO ZOHLOLO LWEKWALITI YAMANZI			
I-WQSU		2	DWAF WQ WMS	Ayikho		
ISIKHUNDLA SE-EWR		Gou 1	RHP	Okwangoku isikhundla se- RHP siphaya phezulu kwindawo yobonisele ngamanzi		
Ukukholelwa kuvavanyo lwe- PES						
Kwibakala eliphantsi – kweliphakathi kuba ikhutshelwe ngaphandle ukusuka kwenye indawo yohlolo (iKarataro R). iinkcukacha (ukusuka kwiprojekthi yangoku nendawo yobonisele ngamanzi ephezulu iRHP) xhasa i-PES ethengisiweyo yekwaliti yamanzi.						
IZithako zeKwaliti yaManzi		IBakala le-PES	ii-Eco specs ze-WQ	Ingaba ikho inkqubo efunekayo?	I-TPC	Ixesha elimiyo lohlolo
Iityuwa ezingezozendalo (mg/L)	MgSO ₄		≤16 mg/L		i-95 th percentile ibe ngu < 16 mg/L	
	Na ₂ SO ₄		≤20 mg/L		i-95 th percentile ibe ngu < 20 mg/L	
	MgCl ₂	Azikho	≤15 mg/L	N/A	i-95 th percentile ibe ngu < 15 mg/L	Rhoqo ngeenyanga ezimbini
	CaCl ₂		≤21 mg/L		i-95 th percentile ibe ngu < 21 mg/L	
	NaCl		≤45 mg/L		i-95 th percentile ibe ngu < 45 mg/L	
Izondlo (mg/L)	SRP	C	≤0.025 mg/L	Ewe- B/C	i-50 th percentile ibe ngu < 0.025 mg/L	Rhoqo ngeenyanga ezimbini
	TIN	A	≤0.25 mg/L	Hayi	i-50 th percentile ibe ngu < 0.25 mg/L	Rhoqo ngeenyanga ezimbini
Ukwahluka kwendalo	pH	Umlambo one-asidi ngendalo	< 6.4	Hayi	i-95 th percentile ibe ngu < 6.4	Rhoqo ngeenyanga ezimbini
	Ubushushu	Azikho iinkcukacha. Uhlolo ngokubona alubonisanga ngxaki yabukho bodaka.	Uluhlu lwendalo	N/A	Uluhlu lobushushu bendalo	Rhoqo ngeenyanga ezimbini
	i-oksijini enyibilikisiweyo		8 mg/L	N/A	5 th percentile ibe ngu > 8 mg/L	Rhoqo ngeenyanga ezimbini
	Ubukho bodaka (NTU)		Ayikho inguqu	N/A	Ayikho inguqu evumelekileyo	Rhoqo ngeenyanga ezimbini
	Ukutsala umbane (mS/m)		A – kuzinzile	≤30 mS/m	No	i-95 th percentile ibe ngu < 30 mS/m
Ukwahluka ngokwemvaka lelo	Chl a: periphyton	Azikho iinkcukacha. Uhlolo ngokubona alubonisanga ngxaki yabukho bodaka	≤ 1.7 mg/m ² (ibakala A)	N/A	i-50 th percentile ibe ngu < 1.7 mg/m ²	Ngekota
	Chl a: phytoplankton		≤ 10 µg/L (ibakala B)		i-50 th percentile ibe ngu < 10 µg/L	
	Ubukhulu bezinto eziphilayo ezingenamathambo (ASPT)	B (olu phando)				
	Inqaku lemigqeku yeentlanzi	C (olu phando)				
Bona ii-Ecospecs zeentlanzi nezezinto eziphilayo ezingenamathambo ngokulandelelanayo.						

iityhefu	Ukutyhefeka kwangaphakathi komjelo	Azikho iinkcukacha Azikho iinkcukacha.	Ayinakubakho ingxaki. Vavanya kuphela xa neziphumo zohlolo lwendalo zibonisa ukuba ikho ingxaki eqatseleyo futhi isizathu sayo asaziwa.
----------	------------------------------------	---	---

UTafle 5.50: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iKnysna kwi- WQSU 1

UMLAMBO	Umlambo iKnysna	IINDAWO ZOHLOLO LWEKWALITI YAMANZI		
I-WQSU	I-WQSU 1	I-RC	Knysna R. @ Millwood K5H002Q01 ('77 – '80; n = 75)	
ISIKHUNDLA SE-EWR	1	I-PES	K5H002Q01 ('04 – '07; n = 26)	
Uvavanyo lokuba nakho		Kwinqanaba eliphakathi njengoko iinkcukacha zovavanyo lwentsukaphi nesimo sangoku zikho ngokwaneleyo		
IZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	Ibakala/Amagqabantshintshi
Iityuwa ezingezozendalo (mg/L)	MgSO ₄	7.18	7.19	U-TEACHA usetyenziselwe uhlolo lweenkcukacha. Iityuwa = ibakala A.
	Na ₂ SO ₄	1.49	4.20	
	MgCl ₂	2.60	2.73	
	CaCl ₂	11.50	3.92	
	NaCl	50.83*	39.54	
Izondlo (mg/L)	SRP	0.011*	0.021	Ibakala A
	TIN	0.06	0.112	Ibakala A
Ukwahluka kwendalo	pH (5 th + 95 th %ile)	4.0 + 6.9	4.5 + 7.2	Ibakala A
	Ubushushu	-	-	Azikho iinkcukacha kodwa alikho ifuthe elilindelekileyo, nangona nje imibala engenziwa kangako ingabangela ukwenyuka kancinci kobushushu bangaphakathi emjelweni.
	i-oksijini enyibilikisiweyo	-	-	Azikho iinkcukacha. Bukho ubungqina bobukho beentlengana ezithile obuboniweyo .
	Ubukho bodaka (NTU)	-	-	Ibakala A.
Ukwahluka ngokwemvakalelo	Ukutsala umbane (mS/m)		9.0	Ibakala A.
	Chl a: periphyton	-	4.08	Ukwanda okuthile kwezondlo kuboniswe ngamaxabiso e-periphyton (ibakala B; n=1).
	Chl a: phytoplankton	-	0.12	
	Ubukhulu bezinto eziphilayo ezingenamathambo	ASPT: mean of 8.1	ASPT: 6.9 MIRAI: 86.92%	Ibakala B lwesimo sangoku.
	Inqaku lemigqeku yeentlanzi	-	FRAI: 86.4%	Ibakala B lwesimo sangoku.
iiDiatoms	-	SPI=18.9	Amanzi anekwaliti ephezulu	
Iityhefu		-	-	Azikho iinkcukacha kodwa alikho ifuthe elilindelekileyo
LULONKE UHLELO LWESIKHUNDLA		A/B kwi-WQSU 1 , nangona iibakala A kwi-EWR 1 nakumphezulu womjelo wesikhundla (umzekelo we-PAI)		

UTafle 5.51: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) kumlambo iKnysna kwi- WQSU

1

Umlambo : iKnysna		I-EWR: 1	Isikhundla sohlolo: K5H002Q01
ii-metrics zekwaliti yamanzi		ii-ECOSPEC	
lityuwa ezingezozendalo*	MgSO ₄	i-95 th percentile yeenkcukacha mayibe ngu ≤ 16 mg/L.	
	Na ₂ SO ₄	i-95 th percentile yeenkcukacha mayibe ngu ≤ 20 mg/L.	
	MgCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 15 mg/L.	
	CaCl ₂	i-95 th percentile yeenkcukacha mayibe ngu ≤ 21 mg/L.	
	NaCl	i-95 th percentile yeenkcukacha mayibe ngu ≤ 45 mg/L.	
	CaSO ₄	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 351 mg/L.	
Ukwahluka kwendalo	EC	i- 95 th percentile yeenkcukacha mayibe ngu ≤ 30 mS/m.	
	pH	-5 th ne- 95 th percentiles yeenkcukacha mayibe phakathi ko-4.5 no- 7.5.	
	Ubushushu	Uluhlu lobushushu bendalo.	
	i-oksijini enyibilikisiweyo	i-5 th percentile yeenkcukacha mayibe ngu ≥ 8.0 mg/L.	
	Ubukho bodaka	Yahluka kancinci kolwaa luhlu lobukho bodaka lwendalo; ukubakho kweentlenge kwiindawo zokuphilisana zangaphakathi komjelo kuvumelekile.	
Izondlo	TIN	i-50 th percentile yeenkcukacha mayibe ngu ≤ 0.25 mg/L.	
	PO ₄ -P	i- 50 th percentile yeenkcukacha mayibe ngu ≤ 0.025 mg/L.	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i-50 th percentile yeenkcukacha mayibe ngu <15 µg/L.	
	Chl-a periphyton	i- 50 th percentile yeenkcukacha mayibe ngu ≤ 12 mg/m ² .	
lityhefu		i-95 th percentile yeenkcukacha mayibe phakathi koluhlu lwekwaliti yamanzi engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubaliwe ku DWAF (1996).	

* Ziza kufunyanwa ngokusebenzisa u-TEACHA xa ne-TPC ye-EC ibethe ngaphaya okanye ungcoliso lwetyuwa lulindelekile.

UTafle 5.52: ii-TPCs zekwaliti yamanzi kumlambo iKnysna kwi- WQSU 1

Umlambo: iKnysna		I-EWR: 1	Isikhundla sohlolo: K5H002Q01
ii-metrics zekwaliti yamanzi		I-TPC	
lityuwa ezingezozendalo*	MgSO ₄	i-95 th percentile yeenkcukacha mayibe ngu- 13 – 16 mg/L.	
	Na ₂ SO ₄	i- 95 th percentile yeenkcukacha mayibe ngu- 16 – 20 mg/L.	
	MgCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu- 12 – 15 mg/L.	
	CaCl ₂	i- 95 th percentile yeenkcukacha mayibe ngu- 17 – 21 mg/L.	
	NaCl	i- 95 th percentile yeenkcukacha mayibe ngu- 36 – 45 mg/L.	
	CaSO ₄	i-95 th percentile yeenkcukacha mayibe ngu- 280 – 351 mg/L.	
Ukwahluka kwendalo	EC	i-95 th percentile yeenkcukacha mayibe ngu- 24 – 30 mS/m.	
	pH	i- 5 th ne- 95 th percentiles yeenkcukacha mayibe phakathi kuka- <4.7 no>7.3.	
	Ubushushu	Uphambuka kancinci kolwaa luhlu lobushushu bendalo.	
	i-oksijini enyibilikisiweyo	i-5 th percentile yeenkcukacha mayibe ngu- 8.2 – 8.0 mg/L.	
	Ubukho bodaka	Iinguqu eziphakathi kusetyenziso lomhlaba wendawo yoboniseko ngamanzi, oko kubangela iingqumba zeentlenge eziphezulu ngokugqithisileyo nobukho bodaka obukumanqanaba aphezulu Moderate.	
Izondlo	TIN	i-50 th percentile yeenkcukacha mayibe ngu 0.2 – 0.25 mg/L.	
	PO ₄ -P	i- 50 th percentile yeenkcukacha mayibe ngu 0.02 – 0.025 mg/L.	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i- 50 th percentile yeenkcukacha mayibe ngu 12 – 15 µg/L.	

Umlambo: iKnysna	I-EWR: 1	Isikhundla sohlolo: K5H002Q01
li-metrics zekwaliti yamanzi	I-TPC	
Chl-a periphyton	i-50 th percentile yeenkcukacha mayibe ngu 10 – 12 mg/m ² .	
lityefu	i-95 th percentile yeenkcukacha mayibe phakathi koluhlu lwekwaliti yamanzi engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe ku DWAF (1996).	

* Ziza kufunyanwa ngokusebenzisa u- TEACHA xa ne-TPC ye-EC ibetha naphaya nongcoliso lweetyuwa lulindelekile.

UTafle 5.53: Amabakala e-PES novavanyo lwesikhundla lulonke kumlambo iGouna kwi-WQSU 2

UMLAMBO	Gouna River	IINDAWO ZOHLOLO LWEKVALITI YAMANZI		
I-WQSU	I-WQSU 2	I-RC	Gouna R. @ Gouna Commonage K5H001Q01 ('77 – '80; n = 76)	
ISIKHUNDLA SE-EWR	2	I-PES	K5H001Q01 ('81 – '84; n = 30)	
Uvavanyo lokuba nakho	Liphantsi inwanaba ngenxa yeenkcukacha ezingephi, ngakumbi kwisimo sangoku			
iZithako zeKwaliti yaManzi		Ixabiso le-RC	Ixabiso le-PES	Ibakala/Amagqabantshintshi
lityuwa ezingezozendalo (mg/L)	MgSO ₄	14.45	23.19	UTEACHA usetyenzisiwe ukuze kwenziwe uhlobo lweenkcukacha. Ukunyukela phezulwana kweetyuwa kuboniwe, i.e. kwinqanaba A/B
	Na ₂ SO ₄	3.78	0.30	
	MgCl ₂	10.75	12.11	
	CaCl ₂	10.59	9.35	
	NaCl	95.29*	102.20	
Izondlo (mg/L)	SRP	0.009*	0.011	Inqanaba A
	TIN	0.070	0.120	Inqanaba A
Ukwahluka kwendalo	pH (5 th + 95 th %ile)	4.0 + 6.8	4.0 + 5.6	Inqanaba A njengoko incinane inguqu xa uyithelekisa naleyo yendalo.
	Ubushushu	-	-	Aziko iinkcukacha kodwa alikho ifuthe elilindelekileyo.
	i-oksijini enyibilikisiweyo	-	-	Aziko iinkcukacha. Zikho iinlenge ezibonakala kumazantsi e- WQSU, i.e. ngaphantsi kwesitishi sokumpompa.
	Ubukho bodaka (NTU)	-	-	
	Ukutsala umbane (mS/m)		15.0	Ibakala A.
Ukwahluka ngokwemvakalelo	Chl a: periphyton	-	43.70	Kuboniswa kukwenyuka komgangatho wexabiso le-periphyton value (Ibakala D; n=1).
	Chl a: phytoplankton	-	0.09	
	Ubukhulu bezinto eziphilayo ezingenamathambo	ASPT: 6.9 – 7.6	ASPT: 7.6 MIRAI: 92.8%	Ibakala A lesimo sangoku.
	Inqaku lemigqeku yeentlanzi	-	FRAI: 93.8%	Ibakala A lesimo sangoku.
	iiDiatoms	-	SPI=19.8	Amanzi anekwaliti ephezulu.
lityhefu		-	-	Aziko iinkcukacha kodwa azikho iityhefu ezilindelekileyo.
LULONKE UHLELO LWESIKHUNDLA		B nge-WQSU 2 , nangona ilibakala A ekunabeni komlambo ukuya kumphezulu womjelo wesitishi sokumpompa (umzekelo we-PAI)		

UTafale 5.54: Imiqathango yendalo yekwaliti yamanzi (ii-EcoSpecs) kumlambo iGouna kwi-WQSU 2

Umlambo : i-Gouna		I-EWR: 2	Isikhundla sohlolo: K5H001Q01
ii-metrics zekwaliti yamanzi		I-ECOSPEC	
lityuwa ezingezozendalo*	MgSO ₄	i-95 th percentile yeenkukacha mayibe ngu ≤ 23 mg/L.	
	Na ₂ SO ₄	i-95 th percentile yeenkukacha mayibe ngu ≤ 20 mg/L.	
	MgCl ₂	i-95 th percentile yeenkukacha mayibe ngu ≤ 15 mg/L.	
	CaCl ₂	i-95 th percentile yeenkukacha mayibe ngu ≤ 21 mg/L.	
	NaCl	i-95 th percentile yeenkukacha mayibe ngu ≤ 191 mg/L.	
	CaSO ₄	i- i-95 th percentile yeenkukacha mayibe ngu ≤ 351 mg/L.	
Ukwahluka kwendalo	EC	i-95 th percentile yeenkukacha mayibe ngu ≤ 43 mS/m.	
	pH	i- 5 th ne-95 th percentiles yeenkukacha mayisuke ku- 4.5 ukuya ku7.5.	
	Ubushushu	Kwendalo ubukhulu becala ukuya kuphambuko oluncinane xa ukuthelekisa nolwaa luhlu lobushushu bendalo.	
	i-oksijini enyibilikisiweyo	i-5 th percentile yeenkukacha mayibe ngu ≥ 7.5 mg/L.	
	Ubukho bodaka	Yahluka kancinci kolwaa luhlu lobukho bodaka lwendalo; ukubakho kweentenge kwiindawo zokuphilisana zangaphakathi komjelo kuvumelekile.	
Izondlo	TIN	i- 50 th percentile yeenkukacha mayibe ngu ≤ 0.15 mg/L.	
	PO ₄ -P	i- 50 th percentile yeenkukacha mayibe ngu ≤ 0.025 mg/L.	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i-50 th percentile yeenkukacha mayibe ngu <15 µg/L.	
	Chl-a periphyton	i- 50 th percentile yeenkukacha mayibe ngu ≤ 84 mg/m ² .	
lityhefu		i-95 th percentile yeenkukacha mayibe phakathi koluhlu lwekwaliti yamanzi engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe ku DWAF (1996).	

* Ziza kufunyanwa ngokusebenzisa u-TEACHA xa ne-TPC ye-EC ibetha ngaphaya okanye ungcoliso lweetyuwa lulindelekile.

UTafale 5.55: Water quality ii-TPCs zekwaliti yamanzi kumlambo iGouna kwi-WQSU 2

Umlambo: iGouna		I-EWR: 2	Isikhundla sohlolo: K5H001Q01
ii-metrics zekwaliti yamanzi		i-TPC	
lityuwa ezingezozendalo*	MgSO ₄	i-95 th percentile yeenkukacha mayibe ngu- 18 – 23 mg/L.	
	Na ₂ SO ₄	i- 95 th percentile yeenkukacha mayibe ngu- 16 – 20 mg/L.	
	MgCl ₂	i-95 th percentile yeenkukacha mayibe ngu- 12 – 15 mg/L.	
	CaCl ₂	i- 95 th percentile yeenkukacha mayibe ngu- 17 – 21 mg/L.	
	NaCl	i-95 th percentile yeenkukacha mayibe ngu- 36 – 45 mg/L.	
	CaSO ₄	i-95 th percentile yeenkukacha mayibe ngu- 153 – 191 mg/L.	
Ukwahluka kwendalo	EC	i- 95 th percentile yeenkukacha mayibe ngu- 35 – 43 mS/m.	
	pH	i-5 th ne-95 th percentiles yeenkukacha mayibe ngu- <4.7 and >7.3.	
	Ubushushu	Luncinci ukuya phakathi uphambuko ukusuka kolwaa luhlu lobushushu bendalo. Iintlobo eziboyika kunene ubushushu azikho ninzi kangako futhi zingaxhaphakanga kunokuba bekulindelekile ngokwemvelaphi. Kuncinci ukuya phakathi ukuphambuka kolwaa luhlu lobushushu bendalo.	
	i-oksijini enyibilikisiweyo	i-5 th percentile of the mayibe ngu- 7.8 – 7.5 mg/L.	
	Ubukho bodaka	Iinguqu eziphakathi kusetyenziso lomhlaba wendawo yoboniselo ngamanzi, oko kubangela iingqumba zeentenge eziphezulu ngokugqithisileyo <u>zexeshana</u> nobukho bodaka obukumanqanaba aphezulu.	

Umlambo: iGouna		i-EWR: 2	Isikhundla sohlolo: K5H001Q01
ii-metrics zekwaliti yamanzi		i-TPC	
Izondlo	TIN	i- 50 th percentile yeenkcukacha mayibe ngu- 0.2 – 0.25 mg/L.	
	PO ₄ -P	i-50 th percentile yeenkcukacha mayibe ngu -0.012 – 0.015 mg/L.	
Ukwahluka ngokwemvakalelo	Chl-a phytoplankton	i-50 th percentile yeenkcukacha mayibe ngu- 12 – 15 µg/L.	
	Chl-a periphyton	i-50 th percentile yeenkcukacha mayibe ngu- 67 – 84 mg/m ² .	
lityhefu		i-95 th percentile yeenkcukacha mayibe phakathi koluhlu lwekwaliti yamanzi engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe ku DWAF (1996).	

6. ICANDELO LOBUNINZI BAMANZI ANGAPHANTSI KOMHLABA

Icandelo lobuninzi bamanzi angaphantsi komhlaba liqingqwe ngokuthi kusetyenziswe amaxabiso (okutsitsa ngokovuselelo nangokwamanzi angaphantsi komhlaba) afunyenwe ngelixa bekuqingqwa amahlelo emijelo yamanzi neenjongo ezibandakanyekayo ngokwekwaliti engqalileyo yomjelo kwindawo yoboniselo ngamanzi i- Breede-Gouritz WMA, (DWS 2017), ebonisiweyo kuTafle 6.1.

Amaxabiso abemi afunyenwe kuvimba weenkukacha weeNkonzo zaManzi ka2011 (i-Water Services dataset ka2011) kwindawo yoboniselo ngamanzi i-Breede, nkukacha ezo zithatyathelwe kuphando lwamahlelo emijelo kwindawoyoboniselo ngamanzi i-Gouritz. UMjelo ohoyene neeMfuno zoLuntu (i-Basic Human Needs Reserve ngelasemzini) uhambisa iinkonzo ezisisiseko zabantu abafumana iinkonzo zamanzi kuloo mjelo uthile – s oko kuquka amanzi okusela, okupheka nawokuzicoca. Kwasetyenziswa umlinganiselo oyi- 25 litres ngomntu ngamnye ngemini.

UTafle 6.1: Umjelo wamanzi angaphantsi komhlaba iBreede-Gouritz

Ummandla woboniselo ngamanzi weminyaka emine	Vuselelo (Mm ³ /a)	Abemi	iiMfuno ezisisiseko zoluntu (Mm ³ /a)	Amanzi angaphantsi komhlaba asisisiseko (Mm ³ /a)	Umjelo (Mm ³ /a)	Umjelo njenge % yovuselelo
G40A	13.06	15963	0.15	3.17	3.32	25.39
G40B	19.19	4612	0.04	5.33	5.37	27.99
G40C	45.16	38379	0.35	6.25	6.60	14.62
G40D	59.72	15963	0.15	14.45	14.60	24.44
G40E	13.19	7670	0.07	4.41	4.48	33.97
G40F	11.28	28422	0.26	2.12	2.38	21.09
G40G	16.02	17112	0.16	3.72	3.88	24.20
G40H	6.53	32767	0.30	1.58	1.88	28.77
G40J	6.92	1574	0.01	2.53	2.54	36.77
G40K	9.13	15963	0.15	4.67	4.82	52.75
G40L	13.96	15963	0.15	1.63	1.78	12.72
G40M	10.57	355	0.003	5.17	5.17	48.94
G50A	7.37	1370	0.01	2.61	2.62	35.58
G50B	6.59	1507	0.01	3.47	3.48	52.86
G50C	8.56	1748	0.02	2.05	2.07	24.13
G50D	5.39	9430	0.09	2.55	2.64	48.91
G50E	4.92	14355	0.13	1.37	1.50	30.51
G50F	6.64	1989	0.02	1.27	1.29	19.40
G50G	2.40	844	0.01	1.43	1.44	59.90
G50H	5.75	736	0.01	3.28	3.29	57.16
G50J	6.07	3325	0.03	1.90	1.93	31.80
G50K	2.72	1748	0.02	0.76	0.78	28.53
H10A	13.15	12494	0.11	0.76	0.87	6.65
H10B	12.20	12494	0.11	0.48	0.59	4.87
H10C	21.28	57300	0.52	2.00	2.52	11.86
H10D	14.89	12494	0.11	2.05	2.16	14.53
H10E	20.35	12494	0.11	3.20	3.31	16.29
H10F	25.24	20720	0.19	1.39	1.58	6.26
H10G	31.82	12494	0.11	0.44	0.55	1.74
H10H	28.48	12494	0.11	2.80	2.91	10.23

Ummandla woboniselolo ngamanzi weminyaka emine	Vuselelo (Mm ³ /a)	Abemi	iiMfuno ezisiSiseko zoluntu (Mm ³ /a)	Amanzi angaphantsi komhlaba asisiseko (Mm ³ /a)	Umjelo (Mm ³ /a)	Umjelo njenge % yovuselelo
H10J	61.45	1035	0.01	7.94	7.95	12.94
H10K	43.17	12494	0.11	7.40	7.51	17.41
H10L	2.76	4268	0.04	0.00	0.04	1.41
H20A	2.42	427	0.00	0.47	0.47	19.58
H20B	5.37	17136	0.16	0.17	0.33	6.08
H20C	2.84	1266	0.01	0.05	0.06	2.17
H20D	8.74	1266	0.01	2.11	2.12	24.27
H20E	14.68	1266	0.01	2.01	2.02	13.77
H20F	8.65	875	0.01	0.32	0.33	3.79
H20G	4.83	1266	0.01	0.47	0.48	9.97
H20H	1.56	140420	1.28	0.07	1.35	86.62
H30A	5.17	1102	0.01	0.33	0.34	6.58
H30B	6.04	39573	0.36	0.16	0.52	8.63
H30C	10.59	1317	0.01	0.07	0.08	0.77
H30D	3.18	926	0.01	0.06	0.07	2.15
H30E	2.95	9784	0.09	0.31	0.40	13.53
H40A	3.74	2233	0.02	0.87	0.89	23.81
H40B	12.26	2152	0.02	0.87	0.89	7.26
H40C	4.90	2233	0.02	0.86	0.88	17.97
H40D	4.18	2233	0.02	1.85	1.87	44.75
H40E	10.91	2233	0.02	0.20	0.22	2.02
H40F	1.07	1798	0.02	0.58	0.60	55.74
H40G	3.22	2233	0.02	0.23	0.25	7.78
H40H	4.71	1217	0.01	0.13	0.14	3.00
H40J	4.44	26455	0.24	0.18	0.42	9.49
H40K	2.99	3916	0.04	0.24	0.28	9.22
H40L	2.47	2290	0.02	0.42	0.44	17.85
H50A	1.42	3842	0.04	0.26	0.30	20.78
H50B	5.04	5825	0.05	0.78	0.83	16.53
H60A	30.87	10083	0.09	2.49	2.58	8.36
H60B	42.43	7900	0.0720875	7.28	7.35	17.33
H60C	30.89	15284	0.14	1.64	1.78	5.76
H60D	14.76	511	0.00	0.95	0.95	6.47
H60E	9.73	10305	0.09	0.71	0.80	8.26
H60F	7.65	3321	0.03	0.66	0.69	9.02
H60G	4.11	10083	0.09	0.64	0.73	17.81
H60H	7.49	10083	0.09	1.14	1.23	16.45
H60J	8.17	10083	0.09	1.31	1.40	17.16
H60K	3.59	10083	0.09	1.04	1.13	31.53
H60L	2.88	10083	0.09	0.87	0.96	33.40
H70A	5.55	4786	0.04	1.47	1.51	27.27
H70B	22.83	19350	0.18	4.17	4.35	19.04

Ummandla woboniselolo ngamanzi weminyaka emine	Vuselelo (Mm ³ /a)	Abemi	iiMfuno ezisiSiseko zoluntu (Mm ³ /a)	Amanzi angaphantsi komhlaba asisiseko (Mm ³ /a)	Umjelo (Mm ³ /a)	Umjelo njenge % yovuselelo
H70C	3.99	4915	0.04	0.23	0.27	6.89
H70D	20.70	4786	0.04	5.53	5.57	26.93
H70E	26.55	6729	0.06	5.16	5.22	19.67
H70F	15.50	2721	0.02	2.31	2.33	15.06
H70G	3.92	4786	0.04	1.26	1.30	33.26
H70H	2.80	746	0.01	1.89	1.90	67.74
H70J	3.95	4786	0.04	1.43	1.47	37.31
H70K	3.03	772	0.01	1.21	1.22	40.17
H80A	16.34	0	0.00	7.21	7.21	44.12
H80B	24.01	0	0.00	6.45	6.45	26.86
H80C	5.75	10 110	0.09	0.61	0.70	12.21
H80D	2.57	0	0.00	1.23	1.23	47.86
H80E	7.66	392	0.00	2.11	2.11	27.59
H80F	5.96	585	0.01	2.72	2.73	45.73
H90A	19.62	0	0.00	9.04	9.04	46.08
H90B	12.96	0	0.00	6.02	6.02	46.45
H90C	5.51	18 526	0.17	1.93	2.10	38.10
H90D	10.38	1 697	0.02	3.29	3.31	31.84
H90E	9.70	6 253	0.06	4.88	4.94	50.90
J11A	2.98	0	0.00	0.00	0.00	0.00
J11B	3.11	0	0.00	0.00	0.00	0.00
J11C	0.22	192	0.00	0.00	0.00	0.80
J11D	3.74	48	0.00	0.00	0.00	0.01
J11E	1.40	4 773	0.04	0.00	0.04	3.11
J11F	0.43	734	0.01	0.00	0.01	1.56
J11G	0.12	84	0.00	0.00	0.00	0.64
J11H	4.01	885	0.01	0.00	0.01	0.20
J11J	6.02	1 333	0.01	0.00	0.01	0.20
J11K	2.52	11 732	0.11	0.00	0.11	4.25
J12A	3.15	0	0.00	0.02	0.02	0.63
J12B	1.55	7 857	0.07	0.00	0.07	4.63
J12C	1.59	152	0.00	0.01	0.01	0.72
J12D	6.32	535	0.00	0.02	0.02	0.39
J12E	1.93	507	0.00	0.02	0.02	1.28
J12F	6.15	336	0.00	0.03	0.03	0.54
J12G	5.66	294	0.00	0.01	0.01	0.22
J12H	4.53	0	0.00	0.02	0.02	0.44
J12J	4.59	0	0.00	0.01	0.01	0.22
J12K	2.44	0	0.00	0.01	0.01	0.41
J12L	6.59	681	0.01	0.05	0.06	0.85
J12M	3.04	0	0.00	0.06	0.06	1.97
J13A	4.10	0	0.00	0.02	0.02	0.49

Ummandla woboniselongamanzi weminyakam emine	Vuselelo (Mm ³ /a)	Abemi	iiMfuno ezisiSiseko zoluntu (Mm ³ /a)	Amanzi angaphantsi komhlaba asisiseko (Mm ³ /a)	Umjelo (Mm ³ /a)	Umjelo njenge % yovuselelo
J13B	2.86	660	0.01	0.03	0.04	1.26
J13C	2.91	0	0.00	0.03	0.03	1.03
J21A	4.28	34 661	0.32	0.00	0.32	7.39
J21B	0.56	389	0.00	0.00	0.00	0.63
J21C	0.12	406	0.00	0.00	0.00	3.09
J21D	0.24	0	0.00	0.00	0.00	0.00
J21E	0.26	0	0.00	0.00	0.00	0.00
J22A	3.04	0	0.00	0.00	0.00	0.00
J22B	1.12	0	0.00	0.00	0.00	0.00
J22C	1.27	241	0.00	0.00	0.00	0.17
J22D	1.22	133	0.00	0.00	0.00	0.10
J22E	1.31	0	0.00	0.00	0.00	0.00
J22F	0.12	1 237	0.01	0.00	0.01	9.41
J22G	2.92	0	0.00	0.00	0.00	0.00
J22H	4.19	919	0.01	0.00	0.01	0.20
J22J	0.90	0	0.00	0.00	0.00	0.00
J22K	0.35	0	0.00	0.00	0.00	0.00
J23A	0.28	2 080	0.02	0.00	0.02	6.78
J23B	0.50	47	0.00	0.00	0.00	0.09
J23C	0.25	12	0.00	0.00	0.00	0.04
J23D	0.70	12	0.00	0.00	0.00	0.02
J23E	2.03	674	0.01	0.18	0.19	9.17
J23F	1.33	6 901	0.06	0.00	0.06	4.73
J23G	0.00	0	0.00	0.00	0.00	0.00
J23H	1.11	0	0.00	0.00	0.00	0.00
J23J	1.82	0	0.00	0.97	0.97	53.30
J24A	2.58	193	0.00	0.00	0.00	0.07
J24B	0.51	1 521	0.01	0.00	0.01	2.72
J24C	0.21	0	0.00	0.00	0.00	0.00
J24D	0.08	314	0.00	0.00	0.00	3.58
J24E	0.39	0	0.00	0.00	0.00	0.00
J24F	1.37	0	0.00	0.00	0.00	0.00
J25A	2.42	200	0.00	1.02	1.02	42.22
J25B	4.45	4 135	0.04	1.23	1.27	28.49
J25C	1.04	1 342	0.01	0.02	0.03	3.10
J25D	2.94	5 170	0.05	0.61	0.66	22.35
J25E	1.12	402	0.00	0.04	0.04	3.90
J31A	7.88	0	0.00	1.13	1.13	14.34
J31B	1.57	0	0.00	0.48	0.48	30.57
J31C	1.87	0	0.00	0.35	0.35	18.72
J31D	2.07	266	0.00	0.38	0.38	18.47
J32A	0.08	50	0.00	0.00	0.00	0.57

Ummandla woboniselongamanzi weminyaka emine	Vuselelo (Mm ³ /a)	Abemi	iiMfuno ezisiSiseko zoluntu (Mm ³ /a)	Amanzi angaphantsi komhlaba asisiseko (Mm ³ /a)	Umjelo (Mm ³ /a)	Umjelo njenge % yovuselelo
J32B	0.01	881	0.01	0.00	0.01	80.39
J32C	0.01	77	0.00	0.00	0.00	7.03
J32D	0.00	0	0.00	0.00	0.00	0.00
J32E	1.76	719	0.01	0.00	0.01	0.37
J33A	4.81	130	0.00	1.44	1.44	29.96
J33B	8.98	680	0.01	1.47	1.48	16.44
J33C	2.83	76	0.00	0.01	0.01	0.38
J33D	3.82	255	0.00	1.24	1.24	32.52
J33E	8.22	13 522	0.12	1.98	2.10	25.59
J33F	4.50	34 154	0.31	2.19	2.50	55.59
J34A	3.08	7 465	0.07	1.48	1.55	50.26
J34B	6.44	1 850	0.02	2.85	2.87	44.52
J34C	9.60	669	0.01	3.51	3.52	36.63
J34D	4.06	1 915	0.02	1.80	1.82	44.77
J34E	2.29	1 321	0.01	1.13	1.14	49.87
J34F	3.44	2 928	0.03	0.47	0.50	14.44
J35A	8.47	31 018	0.28	1.20	1.48	17.51
J35B	8.12	5 831	0.05	1.24	1.29	15.93
J35C	1.98	3 633	0.03	0.88	0.91	46.12
J35D	9.82	5 284	0.05	3.65	3.70	37.66
J35E	1.33	1 224	0.01	0.21	0.22	16.63
J35F	6.67	2 796	0.03	2.02	2.05	30.67
J40A	9.73	970	0.01	5.03	5.04	51.79
J40B	5.45	0	0.00	2.71	2.71	49.72
J40C	15.81	1 076	0.01	6.58	6.59	41.68
J40D	10.21	8 056	0.07	4.20	4.27	41.86
J40E	7.48	1 908	0.02	3.45	3.47	46.36
K10A	2.34	53 970	0.49	1.16	1.65	70.62
K10B	1.96	4 727	0.04	1.20	1.24	63.43
K10C	4.43	0	0.00	2.33	2.33	52.60
K10D	2.53	1 579	0.01	1.10	1.11	44.05
K10E	13.70	4 122	0.04	4.30	4.34	31.66
K10F	2.82	7 002	0.06	0.99	1.05	37.37
K20A	19.85	9 650	0.09	6.15	6.24	31.43
K30A	28.06	6 994	0.06	7.15	7.21	25.71
K30B	21.52	6 334	0.06	5.03	5.09	23.64
K30C	27.80	146 970	1.34	7.83	9.17	32.99
K30D	18.44	9 839	0.09	7.43	7.52	40.78
K40A	8.99	512	0.00	3.79	3.79	42.21
K40B	13.52	0	0.00	4.85	4.85	35.87
K40C	17.00	6 256	0.06	4.32	4.38	25.75
K40D	17.74	20 130	0.18	3.71	3.89	21.95

Ummandla woboniselol ngamanzi weminyaka emine	Vuselelo (Mm ³ /a)	Abemi	iiMfuno ezisiSiseko zoluntu (Mm ³ /a)	Amanzi angaphantsi komhlaba asisiseko (Mm ³ /a)	Umjelo (Mm ³ /a)	Umjelo njenge % yovuselelo
K40E	26.56	13 515	0.12	10.61	10.73	40.41
K50A	27.43	49	0.00	10.09	10.09	36.79
K50B	24.71	14 745	0.13	8.58	8.71	35.27
K60A	6.43	154	0.00	4.20	4.20	65.34
K60B	8.43	43	0.00	5.70	5.70	67.62
K60C	10.95	1 668	0.02	6.60	6.62	60.41
K60D	23.54	681	0.01	12.43	12.44	52.83
K60E	6.39	2 392	0.02	3.95	3.97	62.16
K60F	14.35	10 113	0.09	9.35	9.44	65.80
K60G	11.31	23 053	0.21	5.02	5.23	46.25
K70A	14.30	5 364	0.05	6.84	6.89	48.17
K70B	20.46	852	0.01	4.46	4.47	21.84

UMJELO OPHAKANYISWAYO WAMANZI ANGAPHANTSI KOMHLABA – IBAKALA LEKWALITI YAMANZI

Ikwaliti yamanzi angaphantsi komhlaba kwiindawo zoboniselol ngamanzi zeminyaka emine – xa sijonge ezi nkukukacha zikhoyo zekhemisti yamanzi – yavavanywa kusetyenziswa olwaa luhlu lomlinganiselo ongqaliweyo wamanzi nekwaliti yamanzi, njengoko kubonisiwe kuTafle 6.2 apha ngezantsi. Isishwankathelo seziphumo zohlelo lwekwaliti yamanzi angaphantsi komhlaba kwinqanaba leminyaka eminye ngokweemfuno ezisisiseko zoluntu sifakiwe kwiitafle ezilandelayo.

UTafle 6.2: ikwaliti yamanzi yendalo naleyo yenziwe ngeekhemikhali

Umda	Uluhlu olungqaliweyo lwekwaliti yamanzi ¹⁾				
	ii-Unithi ze-pH	Ihlelo 0	Ihlelo I	Ihlelo II	Ihlelo III
i-pH		6 – 9	5 – 6 & 9 – 9.5	4 – 5 & > 9.5 – 10	<4 & > 10
Ukutsala umbane	mS/m	< 70	70 - 150	150 – 370	> 370
I-Calcium njenge-Ca	mg/l	< 80	80 - 150	150 – 300	> 300
I-Magnesium as Mg	mg/l	< 70	70 - 100	100 – 200	> 200
I-Sodium as Na	mg/l	< 100	100 - 200	200 – 400	> 400
I-Chloride as Cl	mg/l	< 100	100 - 200	200 – 600	> 600
I-Sulphate as SO ₄	mg/l	< 200	200 - 400	400 – 600	> 600
I-Nitrate as NO _x -N	mg/l	< 6	6 - 10	10 – 20	> 20
I-Fluoride as F	mg/l	<0.7	0.7 – 1.0	1.0 – 1.5	> 1.5

1) Uluhlu lweencwadi ezisetyenzisiweyo: Ubugcisa bokuhlela ngokweNgxelo yeKomishoni yoPhando ngezaManzi– i-Water Research Commission: Quality of Domestic Water Supplies – Volume 1. Report No. TT 101/98, Second Edition, 1998.

Iqondo lekwaliti yamanzi zange livavanywe kwezi ndawo zoboniselol ngamanzi zilandelayo ngenxa yeenkcukacha ezinganelanga zolwazi olunekayo (oko kukuthi azikho iinkukukacha malunga nekwaliti yobumeli yamanzi angaphantsi komhlaba):

- G40B; G40D; G40E; G40G; G40K
- G50A; G50B; G50C; G50D; G50G; G50J
- H10D; H10E; H10J; H10K
- H20C; H20G; H20H
- H30B; H30E
- H40C; H40D; H40E; H40G; H40J; H40K; H40L
- H50A; H50B
- H60A; H60B; H60C; H60D; H60E; H60F; H60G; H60H; H60J; H60K; H60L
- H70A; H70C; H70D; H70E; H70F; H70G; H70H; H70J

- J11A; J11C; J11D; J11F; J11H; J11J; J11K
- J12A; J12C; J12E; J12J; J12M
- J13A; J13C, J22A; J22G; J22K, J23B; J23E; J23J, J24, J25B; J25D; J25E
- J31A; J31B; J31C, J32D, J33A; J33B; J33C, J34C; J34E, J35C; J35E; J35F
- J40A; J40B; J40C
- K10C; K10E; K10F
- K20A
- K30A; K30C; K30D
- K40A; K40B; K40C; K40D; K40E
- K50A
- K60A; K60B; K60C; K60D; K60E; K60F; K60G
- K70A; K70B

UTaflele 6.3. UMjelo weKwaliti yaManzi angaPhantsi komhlaba: kuMmandla woLawulo lwaManzi iBreede Gouritz
 iINdawo zoBoniselo ngaManzi: G40C, G40F, G40J, G40L, G40M, G40H, G40I, G40K, G40N, G40O, G40P, G40Q, G40R, G40S, G40T, G40U, G40V, G40W, G40X, G40Y, G40Z

Umda wekhemikhali	I-Unithi	Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾										Umjelo i-BHN ²⁾					Umjelo weKwaliti yaManzi ³⁾				
		INani leeSampulu					Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾					Umjelo we-BHN ²⁾					Umjelo weKwaliti yaManzi ³⁾				
		G40C	G40F	G40J	G40L	G40M	G40C	G40F	G40J	G40L	G40M	G40C	G40F	G40J	G40L	G40M	G40C	G40F	G40J	G40L	G40M
i-pH		39	61	45	43	7.09	7.26	7.01	8.16	5.0 - 9.5	6.38-7.80	6.53-7.99	6.31-7.71	7.34-8.97							
ukutsala umbane	mS/m	39	61	45	43	20.70	20.90	21.20	208.00	<150	22.77	22.99	23.32	228.80							
i-Calcium njengo- Ca	mg/l	36	57	42	40	7.94	7.67	7.68	105.32	<150	8.73	8.43	8.45	115.85							
i-Magnesium njengo- Mg	mg/l	36	57	42	40	3.30	3.30	3.40	33.40	<100	3.63	3.63	3.74	36.74							
i-Sodium njengo- Na	mg/l	36	55	42	38	18.08	18.40	18.32	253.05	<200	19.89	20.24	20.16	253.05							
i-Chloride njengo-Cl	mg/l	36	58	42	40	31.94	32.69	32.57	391.39	<200	35.13	35.96	35.83	391.39							
i-Sulphate njengo- SO ₄	mg/l	36	58	42	40	5.71	5.35	5.81	78.14	<400	6.28	5.89	6.39	85.95							
i-Nitrate njengo- NO ₃ -N	mg/l	36	57	42	38	0.04	0.05	0.06	5.17	<10	0.04	0.06	0.07	5.68							
i-Fluoride njengo- F	mg/l	36	54	42	35	0.30	0.31	0.29	0.28	<1.0	0.33	0.34	0.32	0.31							
Umda wekhemikhali		iINdawo zoBoniselo ngaManzi G40M, G50E, G50F, G50H										Umjelo weKwaliti yaManzi angaphantsi komhlaba³⁾									
		INani leeSampulu					Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾					Umjelo we-BHN ²⁾					Umjelo weKwaliti yaManzi ³⁾				
		G40M	G50E	G50F	G50H	G40M	G50E	G50F	G50H	G40M	G50E	G50F	G50H	G40M	G50E	G50F	G50H	G40M	G50E	G50F	G50H
i-pH		37	49	45	23	8.17	6.21	8.17	8.02	5.0 - 9.5	7.35-8.99	5.59-6.83	7.35-8.99	7.22-8.82							
ukutsala umbane	mS/m	37	49	45	23	109.10	54.90	91.90	1630.00	<150	120.01	60.39	101.09	1630.00							
i-Calcium njengo- Ca	mg/l	37	44	42	23	88.34	4.37	84.20	182.00	<150	97.18	4.81	92.62	182.00							
i-Magnesium njengo- Mg	mg/l	37	44	42	23	15.03	8.91	12.37	358.00	<100	16.54	9.81	13.61	358.00							
i-Sodium njengo- Na	mg/l	37	42	40	23	113.15	80.39	85.23	3107.18	<200	124.47	88.43	93.76	3107.18							
i-Chloride njengo-Cl	mg/l	37	44	42	23	203.98	143.85	153.44	5302.10	<200	203.98	158.24	188.79	5302.10							
i-Sulphate njengo- SO ₄	mg/l	37	44	41	23	31.52	20.58	25.59	742.40	<400	34.67	22.64	28.15	742.40							
i-Nitrate njengo- NO ₃ -N	mg/l	37	43	40	23	1.70	0.06	0.26	0.13	<10	1.87	0.07	0.29	0.14							
i-Fluoride njengo- F	mg/l	37	40	39	23	0.18	0.11	0.22	1.35	<1.0	0.20	0.12	0.25	1.49							

Umda wekhemikhali	I-Uniti	iIndawo zoBoniselo ngaManzi G50J, H10A, H10B, H10C													
		Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo 1)						Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾						
		INani leeSampulu		H10C		H10B			H10A		H10B		H10C		
G50J	H10A	H10B	H10C	G50J	H10A	H10B	H10C	G50J	H10A	H10B	H10C	G50J	H10A	H10B	H10C
i-pH		9	34	73	60	7.58	7.69	7.88	7.33	5.0 - 9.5	6.82-8.34	6.92-8.46	7.09-8.67	6.60-8.06	
ukutsala umbane	mS/m	9	34	73	60	142.90	180.00	49.00	45.60	<150	157.19	180.00	53.90	50.16	
i-Calcium njengo- Ca	mg/l	9	34	70	60	92.00	65.00	46.98	20.35	<150	101.20	71.50	51.67	22.39	
i-Magnesium njengo- Mg	mg/l	9	34	70	60	23.40	54.25	6.72	12.85	<100	25.74	59.68	7.39	14.14	
i-Sodium njengo- Na	mg/l	9	34	68	60	194.60	219.35	36.21	49.80	<200	200.00	241.29	39.83	54.78	
i-Chloride njengo-Cl	mg/l	9	34	70	60	338.90	380.55	80.09	106.25	<200	338.90	418.61	88.10	116.88	
i-Sulphate njengo- SO ₄	mg/l	9	34	70	60	38.70	153.60	21.75	22.30	<400	42.57	168.96	23.92	24.53	
i-Nitrate njengo- NO ₃ -N	mg/l	9	34	70	60	1.04	0.42	0.17	0.44	<10	1.14	0.46	0.19	0.48	
i-Fluoride njengo- F	mg/l	9	34	68	60	0.16	0.34	0.18	0.17	<1.0	0.18	0.37	0.20	0.19	
Umda wekhemikhali	I-Uniti	iIndawo zoBoniselo ngaManzi H10F, H10G, H10H, H10L													
		Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo 1)						Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾						
		INani leeSampulu		H10L		H10H			H10G		H10H		H10L		
H10F	H10G	H10H	H10L	H10F	H10G	H10H	H10L	H10F	H10G	H10H	H10L	H10F	H10G	H10H	H10L
i-pH		15	117	54	82	7.29	7.05	7.59	6.98	5.0 - 9.5	6.56-8.02	6.35-7.76	6.83-8.35	6.28-7.68	
ukutsala umbane	mS/m	15	117	54	82	24.70	13.40	117.70	9.87	<150	27.17	14.74	129.47	10.85	
i-Calcium njengo- Ca	mg/l	15	111	53	78	6.10	7.49	28.50	3.70	<150	6.71	8.24	31.35	4.07	
i-Magnesium njengo- Mg	mg/l	15	112	53	78	12.00	2.80	28.30	2.12	<100	13.20	3.08	31.13	2.34	
i-Sodium njengo- Na	mg/l	12	107	52	76	18.85	12.10	176.10	8.60	<200	20.74	13.31	193.71	9.46	
i-Chloride njengo-Cl	mg/l	15	114	54	79	30.50	18.45	308.75	14.80	<200	33.55	20.30	308.75	16.28	
i-Sulphate njengo- SO ₄	mg/l	12	109	52	79	4.15	5.50	53.65	3.00	<400	4.57	6.05	59.02	3.30	
i-Nitrate njengo- NO ₃ -N	mg/l	15	112	54	78	0.06	0.25	0.03	0.18	<10	0.07	0.28	0.03	0.20	
i-Fluoride njengo- F	mg/l	15	110	54	75	0.31	0.16	0.57	0.12	<1.0	0.34	0.17	0.62	0.13	

Umda wekhemikhali	i-Uniti	iIndawo zoBoniselo ngaManzi H20A, H20B, H20D, H20E													
		iNani leeSampulu				iInkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾				
		H20A	H20B	H20D	H20E	H20A	H20B	H20D	H20E		H20A	H20B	H20D	H20E	
i-pH		85	344	12	15	7.20	7.09	6.77	6.46	5.0 – 9.5	6.48-7.92	6.38-7.79	6.09-7.44	5.81-7.11	
ukutsala umbane	mS/m	85	344	12	15	50.60	40.75	12.75	2.30	<150	55.66	44.83	14.03	2.53	
i-Calcium njengo- Ca	mg/l	85	344	12	15	34.80	23.45	9.70	0.50	<150	38.28	25.80	10.67	0.55	
i-Magnesium njengo- Mg	mg/l	85	344	12	15	5.60	9.15	2.15	0.50	<100	6.16	10.07	2.37	0.55	
i-Sodium njengo- Na	mg/l	85	344	11	15	25.70	31.80	7.70	2.60	<200	28.27	34.98	8.47	2.86	
i-Chloride njengo-Cl	mg/l	85	344	12	15	55.00	50.55	13.90	1.50	<200	60.50	55.61	15.29	1.65	
i-Sulphate njengo- SO ₄	mg/l	85	344	11	15	81.20	39.65	4.00	2.00	<400	89.32	43.62	4.40	2.20	
i-Nitrate njengo- NO ₃ -N	mg/l	85	344	12	15	0.05	0.90	0.33	0.05	<10	0.06	0.99	0.36	0.06	
i-Fluoride njengo- F	mg/l	85	344	12	15	0.11	0.13	0.05	0.05	<1.0	0.12	0.14	0.06	0.06	

Umda wekhemikhali	i-Uniti	iIndawo zoBoniselo ngaManzi H20F, H30A, H30C, H30D													
		iNani leeSampulu				iInkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾				
		H20F	H30A	H30C	H30D	H20F	H30A	H30C	H30D		H20F	H30A	H30C	H30D	
i-pH		302	9	21	45	7.10	7.95	7.70	7.81	5.0 – 9.5	6.39-7.80	7.16-8.75	6.93-8.47	7.03-8.59	
ukutsala umbane	mS/m	302	9	21	45	29.00	142.00	41.70	19.50	<150	31.90	156.20	45.87	21.45	
i-Calcium njengo- Ca	mg/l	302	9	21	42	22.80	63.80	36.40	17.10	<150	25.08	70.18	40.04	18.81	
i-Magnesium njengo- Mg	mg/l	302	9	21	42	4.70	32.50	6.00	3.31	<100	5.17	35.75	6.60	3.64	
i-Sodium njengo- Na	mg/l	302	9	21	40	24.90	191.90	37.00	10.94	<200	27.39	200.00	40.70	12.04	
i-Chloride njengo-Cl	mg/l	302	9	21	43	28.30	273.50	82.00	18.08	<200	31.13	300.85	90.20	19.89	
i-Sulphate njengo- SO ₄	mg/l	302	9	21	43	28.00	121.90	48.00	4.30	<400	30.80	134.09	52.80	4.73	
i-Nitrate njengo- NO ₃ -N	mg/l	302	9	21	42	0.11	0.02	0.02	0.13	<10	0.12	0.02	0.02	0.14	
i-Fluoride njengo- F	mg/l	302	9	16	40	0.10	0.55	0.23	0.16	<1.0	0.11	0.61	0.25	0.18	

Umda wekhemikhali	I-Uniti	IIndawo zoBoniselo ngaManzi H40A, H40B, H40F, H70B																			
		Inani leeSampulu						Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾						Umjelo we-BHN ²⁾		Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾					
		H40A	H40B	H40F	H70B	H40A	H40B	H40F	H70B	H40A	H40B	H40F	H70B	H40A	H40B	H40F	H70B				
i-pH		54	14	17	9	7.86	7.35	6.54	8.21	112.50	10.88	28.90	925.00	123.75	11.97	31.79	7.07-8.65	6.62-8.09	5.89-7.19	7.39-9.03	
ukutsala umbane	mS/m	55	14	17	9	92.38	2.24	14.80	98.9	44.90	0.93	7.00	259.10	101.62	2.47	16.28	108.79	49.39	1.02	7.70	259.10
i-Calcium njengo- Ca	mg/l	48	14	17	9	88.94	21.43	26.30	1722.90	127.45	21.15	35.90	2741.20	97.83	23.57	28.93	140.20	23.27	39.49	2741.20	
i-Magnesium njengo- Mg	mg/l	47	12	17	9	229.91	3.50	20.70	338.60	0.06	0.18	4.70	0.02	252.90	3.85	22.77	0.06	0.20	5.27	0.02	
i-Sodium njengo- Na	mg/l	50	14	17	9	0.43	0.36	0.05	1.75					0.47	0.39	0.06					1.75
i-Chloride njengo- Cl	mg/l	50	14	17	9																
i-Sulphate njengo- SO ₄	mg/l	50	14	17	9																
i-Nitrate njengo- NO ₃ -N	mg/l	50	14	17	9																
i-Fluoride njengo- F	mg/l	46	13	17	9																
Umda wekhemikhali	I-Uniti	IIndawo zoBoniselo ngaManzi H70K, H90D, H90E, J11B																			
		Inani leeSampulu						Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾						Umjelo we-BHN ²⁾		Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾					
		H70K	H90D	H90E	J11B	H70K	H90D	H90E	J11B	H70K	H90D	H90E	J11B	H70K	H90D	H90E	J11B				
i-pH		29	21	85	37	7.50	8.12	8.09	7.80	121.00	102.70	179.00	81.30	6.75-8.25	7.31-8.94	7.28-8.90	7.02-8.58	133.10	112.97	179.00	89.43
ukutsala umbane	mS/m	26	21	81	37	46.20	53.66	87.18	60.50	19.20	17.40	28.50	18.40	50.82	59.02	95.89	66.55	21.12	19.14	31.35	20.24
i-Calcium njengo- Ca	mg/l	26	21	80	37	166.95	144.96	208.48	79.20	273.10	247.79	358.72	94.70	183.65	159.45	208.48	87.12	273.10	247.79	358.72	104.17
i-Magnesium njengo- Mg	mg/l	26	21	82	37	29.88	33.16	36.65	68.60	0.02	0.83	3.40	1.41	32.87	36.47	40.31	75.46	0.02	0.91	3.74	1.55
i-Sodium njengo- Na	mg/l	26	20	79	37	0.52	0.14	0.18	0.82					0.57	0.15	0.19	0.90				
i-Chloride njengo- Cl	mg/l	26	20	77	37																
i-Sulphate njengo- SO ₄	mg/l	26	20	77	37																
i-Nitrate njengo- NO ₃ -N	mg/l	26	20	77	37																
i-Fluoride njengo- F	mg/l	26	20	77	37																

Umda wekhemikhali	I-Unithi	iINDAWO ZO BONISELO NGA MANZI J11E, J11G, J12B, J12D												
		iNani leeSampulu				iNkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J11E	J11G	J12B	J12D	J11E	J11G	J12B	J12D		J11E	J11G	J12B	J12D
i-pH		81	20	17	11	8.19	8.18	7.73	8.23	5.0 - 9.5	7.37-9.01	7.36-9.00	6.96-8.50	7.41-9.05
ukutsala umbane	mS/m	81	20	17	11	174.30	164.50	368.00	195.00	<150	174.30	164.50	368.00	195.00
i-Calcium njengo- Ca	mg/l	77	20	17	11	98.70	115.05	117.10	68.30	<150	108.57	126.56	128.81	75.13
i-Magnesium njengo- Mg	mg/l	77	20	17	11	45.00	44.65	129.30	50.60	<100	49.50	49.12	129.30	55.66
i-Sodium njengo- Na	mg/l	73	20	17	11	197.89	190.65	335.70	252.80	<200	217.68	200.00	335.70	252.80
i-Chloride njengo-Cl	mg/l	75	20	17	11	250.10	294.70	726.10	415.00	<200	250.10	294.70	726.10	415.00
i-Sulphate njengo- SO ₄	mg/l	77	20	17	11	179.50	120.45	144.20	104.80	<400	197.45	132.50	158.62	115.28
i-Nitrate njengo- NO ₃ -N	mg/l	71	20	17	11	0.30	1.18	0.06	0.02	<10	0.33	1.29	0.07	0.02
i-Fluoride njengo- F	mg/l	67	20	17	11	0.88	0.92	0.80	0.53	<1.0	0.96	0.92	0.88	0.58
Umda wekhemikhali	I-Unithi	iINDAWO ZO BONISELO NGA MANZI J12F, J12G, J12H, J12K												
		iNani leeSampulu				iNkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J12F	J12G	J12H	J12K	J12F	J12G	J12H	J12K		J12F	J12G	J12H	J12K
i-pH		10	11	13	10	8.23	8.20	7.75	8.13	5.0 - 9.5	7.41-9.05	7.38-9.02	6.98-8.53	7.32-8.94
ukutsala umbane	mS/m	10	11	13	10	306.50	323.00	157.00	462.00	<150	306.50	323.00	157.00	462.00
i-Calcium njengo- Ca	mg/l	10	11	13	10	79.65	103.10	121.90	212.95	<150	87.62	113.41	134.09	212.95
i-Magnesium njengo- Mg	mg/l	10	11	13	10	81.35	91.80	26.00	112.55	<100	89.49	100.98	28.60	123.81
i-Sodium njengo- Na	mg/l	10	11	13	10	406.70	262.30	151.50	703.25	<200	406.70	262.30	166.65	703.25
i-Chloride njengo-Cl	mg/l	10	11	13	10	691.30	795.60	258.70	1168.85	<200	691.30	795.60	258.70	1168.85
i-Sulphate njengo- SO ₄	mg/l	10	11	13	10	241.45	107.20	156.00	459.80	<400	265.60	117.92	171.60	459.80
i-Nitrate njengo- NO ₃ -N	mg/l	10	11	13	10	0.02	0.02	0.02	0.02	<10	0.02	0.02	0.02	0.02
i-Fluoride njengo- F	mg/l	10	11	13	10	0.56	0.50	0.39	0.55	<1.0	0.61	0.55	0.43	0.61

Umda wekhemikhali	i-Unithi	iIndawo zoBoniselo ngaManzi J12L, J13B, J21A, J21B												
		iNani leeSampulu				Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Groundwater Quality Reserve ³⁾			
		J12L	J13B	J21A	J21B	J12L	J13B	J21A	J21B		J12L	J13B	J21A	J21B
i-pH		57	15	64	56	7.54	7.98	7.97	7.60	6.79-8.29	7.18-8.78	7.17-8.76	6.84-8.35	
ukutsala umbane	mS/m	57	15	64	56	25.50	418.00	61.40	100.50	28.05	418.00	67.54	110.55	
i-Calcium njengo- Ca	mg/l	54	14	60	56	16.11	94.90	58.18	60.10	17.72	104.39	64.00	66.11	
i-Magnesium njengo- Mg	mg/l	54	15	59	56	2.89	93.80	18.81	19.45	3.18	103.18	20.69	21.40	
i-Sodium njengo- Na	mg/l	52	15	60	54	21.85	623.60	43.23	116.95	24.03	623.60	47.55	128.65	
i-Chloride njengo- Cl	mg/l	53	15	58	56	30.71	906.39	18.77	104.70	33.78	906.39	20.84	115.17	
i-Sulphate njengo- SO ₄	mg/l	54	15	57	54	7.55	253.60	75.14	113.80	8.31	278.96	82.65	125.18	
i-Nitrate njengo- NO ₃ -N	mg/l	52	15	54	56	0.03	0.35	0.09	2.56	0.03	0.39	0.09	2.81	
i-Fluoride njengo- F	mg/l	51	15	54	56	0.21	0.72	0.46	0.87	0.23	0.79	0.51	0.96	
Umda wekhemikhali	i-Unithi	iIndawo zoBoniselo ngaManzi J21C, J21D, J21E, J22B												
		iNani leeSampulu				Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J21C	J21D	J21E	J22B	J21C	J21D	J21E	J22B		J21C	J21D	J21E	J22B
i-pH		107	40	26	21	7.75	7.66	7.96	8.00	6.98-8.53	6.89-8.42	7.16-8.76	7.20-8.80	
ukutsala umbane	mS/m	107	40	26	21	76.40	85.20	78.80	74.20	84.04	93.72	86.68	81.62	
i-Calcium njengo- Ca	mg/l	107	40	26	21	56.90	43.30	61.00	60.30	62.59	47.63	67.10	66.33	
i-Magnesium njengo- Mg	mg/l	107	40	26	21	16.40	17.70	16.00	24.80	18.04	19.47	17.60	27.28	
i-Sodium njengo- Na	mg/l	107	38	26	21	83.80	107.90	83.40	67.90	92.18	118.69	91.74	74.69	
i-Chloride njengo- Cl	mg/l	107	40	26	21	76.90	102.60	93.00	57.20	84.59	112.86	102.30	62.92	
i-Sulphate njengo- SO ₄	mg/l	107	38	26	21	83.50	95.70	53.05	63.40	91.85	105.27	58.36	69.74	
i-Nitrate njengo- NO ₃ -N	mg/l	107	39	26	21	3.13	3.55	2.23	1.95	3.44	3.91	2.45	2.15	
i-Fluoride njengo- F	mg/l	107	40	26	21	0.68	1.00	0.95	0.91	0.75	1.00	0.95	1.00	

Umda wekhemikhali	I-Unithi	IiNdawo zoBoniselo ngamaManzi J22C, J22D, J22E, J22F												
		Inani leeSampulu				Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J22C	J22D	J22E	J22F	J22C	J22D	J22E	J22F		J22C	J22D	J22E	J22F
i-pH		26	46	80	24	8.00	7.94	7.79	7.80	5.0 - 9.5	7.20-8.80	7.15-8.73	7.01-8.57	7.02-8.58
ukutsala umbane	mS/m	26	46	80	24	97.10	89.85	88.70	103.45	<150	106.81	98.84	97.57	113.80
i-Calcium njengo- Ca	mg/l	26	46	80	24	50.15	44.95	61.80	82.10	<100	55.17	49.45	67.98	90.31
i-Magnesium njengo- Mg	mg/l	26	46	80	24	27.00	24.80	21.00	18.95	<100	29.70	27.28	23.10	20.85
i-Sodium njengo- Na	mg/l	26	46	80	24	102.75	118.25	95.95	86.35	<200	113.03	130.08	105.55	94.99
i-Chloride njengo-Cl	mg/l	26	46	80	24	94.00	102.75	92.40	109.40	<200	103.40	113.03	101.64	120.34
i-Sulphate njengo- SO ₄	mg/l	26	46	80	24	123.80	121.30	90.80	110.70	<400	136.18	133.43	99.88	121.77
i-Nitrate njengo- NO ₃ -N	mg/l	26	46	80	24	1.78	1.47	4.12	3.82	<10	1.95	1.61	4.53	4.20
i-Fluoride njengo- F	mg/l	26	46	80	24	1.03	1.10	1.05	0.89	<1.0	1.03	1.10	1.05	0.97
Umda wekhemikhali	I-Unithi	IiNdawo zoBoniselo ngamaManzi J22H, J22J, J23A, J23C												
		Inani leeSampulu				Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J22H	J22J	J23A	J23C	J22H	J22J	J23A	J23C		J22H	J22J	J23A	J23C
i-pH		32	26	39	10	7.79	7.83	7.93	7.78	5.0 - 9.5	7.01-8.56	7.05-8.61	7.14-8.72	7.00-8.56
ukutsala umbane	mS/m	32	26	39	10	145.05	83.20	238.00	136.00	<150	145.05	91.52	238.00	149.60
i-Calcium njengo- Ca	mg/l	32	26	36	10	105.60	50.75	125.25	126.75	<150	116.16	55.83	137.78	139.43
i-Magnesium njengo- Mg	mg/l	32	26	36	10	38.25	20.45	16.26	32.70	<100	42.08	22.50	17.89	35.97
i-Sodium njengo- Na	mg/l	32	26	35	10	158.05	86.90	342.80	143.70	<200	173.86	95.59	342.80	158.07
i-Chloride njengo-Cl	mg/l	32	26	35	10	208.55	82.15	265.46	159.05	<200	229.41	90.37	265.46	174.96
i-Sulphate njengo- SO ₄	mg/l	32	26	36	10	225.10	117.50	497.01	112.50	<400	247.61	129.25	497.01	123.75
i-Nitrate njengo- NO ₃ -N	mg/l	32	26	35	10	3.31	3.02	0.04	1.10	<10	3.64	3.32	0.04	1.21
i-Fluoride njengo- F	mg/l	32	26	33	10	1.09	0.90	2.14	0.69	<1.0	1.19	0.99	2.14	0.76

Umda wekhemikhali	i-Unithi	iIndawo zoBoniselo ngalManzi J23D, J23F, J24A, J24B															
		iNani leeSampulu						Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾		Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J23D	J23F	J24A	J24B	J24C	J24D	J23F	J24A	J24B	J24C	J24D	J24E	J24F	J23D	J23F	J24A
i-pH		26	85	168	220	8.10	7.97	7.77	7.79	7.29-8.91	7.17-8.76	5.0 - 9.5	7.29-8.91	7.17-8.76	6.99-8.54	7.01-8.56	
ukusala umbane	mS/m	26	85	168	220	155.00	126.70	86.30	86.54	155.00	139.37	<150	155.00	139.37	94.93	95.19	
i-Calcium njengo- Ca	mg/l	26	83	168	213	98.95	51.07	70.60	76.59	108.85	56.17	<150	108.85	56.17	77.66	84.24	
i-Magnesium njengo- Mg	mg/l	26	83	168	214	40.65	28.60	18.55	17.15	44.72	31.46	<100	44.72	31.46	20.41	18.87	
i-Sodium njengo- Na	mg/l	26	80	168	210	166.55	120.85	91.25	83.50	183.21	132.94	<200	183.21	132.94	100.38	91.85	
i-Chloride njengo- Cl	mg/l	26	83	168	214	208.95	140.10	78.80	78.20	208.95	154.11	<200	208.95	154.11	86.68	86.02	
i-Sulphate njengo- SO ₄	mg/l	26	83	168	214	151.40	157.15	103.40	69.00	166.54	172.86	<400	166.54	172.86	113.74	75.90	
i-Nitrate njengo- NO ₃ -N	mg/l	26	83	168	212	1.89	0.12	1.85	3.45	2.08	0.13	<10	2.08	0.13	2.04	3.80	
i-Fluoride njengo- F	mg/l	26	81	168	209	0.79	0.48	0.92	0.81	0.86	0.52	<1.0	0.86	0.52	1.01	0.89	

Umda wekhemikhali	i-Unithi	iIndawo zoBoniselo ngalManzi J24C, J24D, J24E, J25A															
		iNani leeSampulu						Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾		Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J24C	J24D	J24E	J25A	J24C	J24D	J24E	J25A	J24C	J24D	J24E	J25A	J24C	J24D	J24E	J25A
i-pH		112	97	18	33	7.70	7.90	7.94	7.11	6.93-8.47	7.11-8.69	5.0 - 9.5	6.93-8.47	7.11-8.69	7.15-8.73	6.40-7.83	
ukusala umbane	mS/m	112	97	18	33	96.70	146.70	161.50	50.90	106.37	146.70	<150	106.37	146.70	161.50	55.99	
i-Calcium njengo- Ca	mg/l	112	93	18	33	68.45	94.20	106.00	14.80	75.30	103.62	<150	75.30	103.62	116.60	16.28	
i-Magnesium njengo- Mg	mg/l	112	93	18	33	18.00	28.80	30.95	10.90	19.80	31.68	<100	19.80	31.68	34.05	11.99	
i-Sodium njengo- Na	mg/l	112	86	18	33	103.40	162.40	198.65	50.40	113.74	178.64	<200	113.74	178.64	198.65	55.44	
i-Chloride njengo- Cl	mg/l	112	95	18	33	120.20	194.50	213.80	96.50	132.22	194.50	<200	132.22	194.50	213.80	106.15	
i-Sulphate njengo- SO ₄	mg/l	112	89	18	33	78.40	114.00	105.70	26.40	86.24	125.40	<400	86.24	125.40	116.27	29.04	
i-Nitrate njengo- NO ₃ -N	mg/l	112	95	18	32	3.90	1.76	0.82	0.02	4.28	1.94	<10	4.28	1.94	0.90	0.02	
i-Fluoride njengo- F	mg/l	112	93	18	32	0.79	0.81	1.05	0.31	0.87	0.89	<1.0	0.87	0.89	1.05	0.34	

Umda wekhemikhali	I-Unithi	iIndawo zoBoniselo ngalManzi J25C, J31D, J32A, J32B																	
		Inani leeSampulu						Inkangeleko yekWaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾		Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾					
		J25C	J31D	J32A	J32B	J25C	J31D	J32A	J32B	J25C	J31D	J32A	J32B	J25C	J31D	J32A	J32B		
i-pH		31	48	19	15	7.11	6.54	7.68	7.86	5.0 - 9.5	6.40-7.83	5.89-7.19	6.91-8.45	7.07-8.65					
ukutsala umbane	mS/m	31	48	19	15	50.90	14.06	165.80	187.00	<150	55.99	15.46	165.80	187.00					
i-Calcium njengo- Ca	mg/l	31	43	19	15	14.80	4.92	85.60	99.50	<150	16.28	5.41	94.16	109.45					
i-Magnesium njengo- Mg	mg/l	31	43	19	15	10.90	2.86	42.50	44.50	<100	11.99	3.14	46.75	48.95					
i-Sodium njengo- Na	mg/l	31	41	19	15	50.40	9.30	172.20	175.10	<200	55.44	10.23	189.42	192.61					
i-Chloride njengo-Cl	mg/l	31	43	19	15	96.50	23.80	203.20	242.60	<200	106.15	26.18	203.20	242.60					
i-Sulphate njengo- SO ₄	mg/l	31	43	19	15	26.40	9.90	180.30	170.80	<400	29.04	10.89	198.33	187.88					
i-Nitrate njengo- NO ₃ -N	mg/l	31	41	19	15	0.02	0.03	3.80	0.88	<10	0.02	0.03	4.18	0.96					
i-Fluoride njengo- F	mg/l	31	39	19	15	0.31	0.29	1.06	0.99	<1.0	0.34	0.32	1.06	0.99					
Umda wekhemikhali	I-Unithi	iIndawo zoBoniselo ngalManzi J32C, J32E, J33D, J33E																	
		Inani leeSampulu						Inkangeleko yekWaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾		Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾					
		J32C	J32E	J33D	J33E	J32C	J32E	J33D	J33E	J32C	J32E	J33D	J33E	J32C	J32E	J33D	J33E		
i-pH		29	41	40	149	7.76	7.55	7.53	6.64	5.0 - 9.5	6.98-8.54	6.80-8.30	6.78-8.29	6.96-7.30					
ukutsala umbane	mS/m	29	41	40	149	127.40	19.80	19.80	12.10	<150	140.14	21.78	21.78	13.31					
i-Calcium njengo- Ca	mg/l	29	40	39	148	80.80	11.07	11.00	2.76	<150	88.88	12.17	12.10	3.04					
i-Magnesium njengo- Mg	mg/l	29	40	39	148	30.90	3.50	3.50	2.07	<100	33.99	3.85	3.85	2.27					
i-Sodium njengo- Na	mg/l	29	39	38	146	143.90	12.40	12.39	13.40	<200	158.29	13.64	13.63	14.74					
i-Chloride njengo-Cl	mg/l	29	40	38	147	170.70	23.87	23.64	22.99	<200	187.77	26.26	26.01	25.29					
i-Sulphate njengo- SO ₄	mg/l	29	39	38	146	121.00	5.72	5.58	6.34	<400	133.10	6.29	6.14	6.97					
i-Nitrate njengo- NO ₃ -N	mg/l	29	39	38	147	2.15	0.04	0.04	0.19	<10	2.37	0.04	0.04	0.20					
i-Fluoride njengo- F	mg/l	29	37	36	146	0.90	0.27	0.27	0.11	<1.0	0.99	0.30	0.30	0.12					

Umda wekhemikhali	I-Unithi	iIndawo zoBoniselo ngamanzi J33F, J34A, J34B, J34D												
		Inani leeSampulu				Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo 1)				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J33F	J34A	J34B	J34D	J33F	J34A	J34B	J34D		J33F	J34A	J34B	J34D
i-pH		70	36	11	22	6.82	6.33	6.77	6.56	5.0 – 9.5	6.14–7.50	5.70–6.96	6.09–7.45	5.90–7.22
ukutsala umbane	mS/m	70	36	11	22	44.60	19.55	114.50	12.54	<150	49.06	21.51	125.95	13.79
i-Calcium njengo- Ca	mg/l	70	36	11	22	11.95	3.26	46.30	2.39	<150	13.15	3.68	50.93	2.63
i-Magnesium njengo- Mg	mg/l	70	36	11	22	7.05	3.44	21.80	1.73	<100	7.76	3.78	23.98	1.91
i-Sodium njengo- Na	mg/l	70	36	11	22	34.25	25.95	136.70	15.40	<200	37.68	28.55	150.37	16.94
i-Chloride njengo-Cl	mg/l	70	36	11	22	61.74	40.85	252.50	27.36	<200	67.92	44.94	252.50	30.10
i-Sulphate njengo- SO ₄	mg/l	70	36	11	22	27.50	5.22	42.20	2.50	<400	30.25	5.74	46.42	2.75
i-Nitrate njengo- NO ₃ -N	mg/l	68	36	11	22	0.08	0.17	0.37	0.22	<10	0.09	0.19	0.41	0.24
i-Fluoride njengo- F	mg/l	69	36	11	22	0.22	0.11	0.29	0.08	<1.0	0.24	0.12	0.32	0.08
Umda wekhemikhali	I-Unithi	iIndawo zoBoniselo ngamanzi J34F, J35A, J35B, J35D												
		Inani leeSampulu				Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo 1)				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾			
		J34F	J35A	J35B	J35D	J34F	J35A	J35B	J35D		J34F	J35A	J35B	J35D
i-pH		9	17	49	49	7.66	8.07	7.01	7.64	5.0 – 9.5	6.89–8.43	7.26–8.88	6.31–7.71	6.88–8.40
ukutsala umbane	mS/m	9	17	49	49	46.60	42.00	46.20	173.10	<150	51.26	46.20	50.82	173.10
i-Calcium njengo- Ca	mg/l	9	17	47	48	6.87	62.70	6.23	86.30	<150	7.55	68.97	6.85	94.93
i-Magnesium njengo- Mg	mg/l	9	17	47	48	7.70	6.30	8.70	35.65	<100	8.47	6.93	9.57	39.22
i-Sodium njengo- Na	mg/l	9	17	45	48	61.80	11.10	57.30	145.70	<200	67.98	12.21	63.03	160.27
i-Chloride njengo-Cl	mg/l	9	17	45	48	101.64	19.10	108.70	239.50	<200	111.81	21.01	119.57	239.50
i-Sulphate njengo- SO ₄	mg/l	9	17	47	48	27.90	11.50	4.85	95.95	<400	30.69	12.65	5.34	105.55
i-Nitrate njengo- NO ₃ -N	mg/l	9	17	45	48	0.05	0.16	4.51	0.64	<10	0.06	0.18	4.96	0.70
i-Fluoride njengo- F	mg/l	9	17	43	47	0.48	0.16	0.10	0.52	<1.0	0.53	0.18	0.11	0.57

Umda wekhemikhali	i-Unithi	iIndawo zoBoniselo ngaManzi J40D, J40E, K10A, K10B													
		iNani leeSampulu				Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾				
		J40D	J40E	K10A	K10B	J40D	J40E	K10A	K10B		J40D	J40E	K10A	K10B	
i-pH		48	63	20	14	7.30	7.52	7.88	7.51	5.0 – 9.5	6.57–8.03	6.77–8.28	7.09–8.67	6.76–8.26	
ukutsala umbane	mS/m	48	63	20	14	83.15	219.00	276.00	214.50	<150	91.47	219.00	276.00	214.50	
i-Calcium njengo- Ca	mg/l	48	63	20	14	12.41	59.40	52.50	32.87	<150	13.65	65.34	57.75	36.15	
i-Magnesium njengo- Mg	mg/l	48	63	20	14	14.72	31.97	44.17	35.74	<100	16.19	35.17	48.58	39.32	
i-Sodium njengo- Na	mg/l	45	63	20	14	131.41	288.45	399.60	288.65	<200	144.55	288.45	399.60	288.65	
i-Chloride njengo- Cl	mg/l	48	63	20	13	213.26	492.38	698.45	545.73	<200	213.26	492.38	698.45	545.73	
i-Sulphate njengo- SO ₄	mg/l	45	63	20	14	31.69	65.21	97.08	59.60	<400	34.86	71.73	106.79	65.56	
i-Nitrate njengo- NO ₃ -N	mg/l	47	62	20	12	0.09	0.04	0.05	0.27	<10	0.10	0.04	0.06	0.29	
i-Fluoride njengo- F	mg/l	48	63	20	13	0.14	0.22	0.22	0.25	<1.0	0.16	0.24	0.24	0.28	
Umda wekhemikhali	i-Unithi	iIndawo zoBoniselo ngaManzi K10D, K30B, K50B													
		iNani leeSampulu				Inkangeleko yeKwaliti yeGW okanye ixabiso eliphakathi layo ¹⁾				Umjelo we-BHN ²⁾	Umjelo wekwaliti yamanzi angaphantsi komhlaba ³⁾				
		K10D	K30B	K50B	K10D	K30B	K50B	K10D	K30B		K50B	K10D	K30B	K50B	
i-pH		11	47	9		7.85	6.83	7.48		5.0 – 9.5	7.07–8.64	6.15–7.51	6.73–8.22		
ukutsala umbane	mS/m	11	47	9		257.00	27.72	61.90		<150	257.00	30.49	68.09		
i-Calcium njengo- Ca	mg/l	11	43	9		30.82	4.27	15.08		<150	33.90	4.70	16.58		
i-Magnesium njengo- Mg	mg/l	11	43	9		28.10	7.49	9.80		<100	30.91	8.24	10.78		
i-Sodium njengo- Na	mg/l	11	41	9		426.18	28.35	85.57		<200	426.18	31.19	94.13		
i-Chloride njengo- Cl	mg/l	11	42	9		533.12	41.92	139.99		<200	533.12	46.11	153.96		
i-Sulphate njengo- SO ₄	mg/l	11	44	9		66.60	7.95	17.54		<400	73.26	8.74	19.29		
i-Nitrate njengo- NO ₃ -N	mg/l	10	43	9		0.09	7.99	0.48		<10	0.10	8.79	0.52		
i-Fluoride njengo- F	mg/l	10	23	9		0.84	0.16	0.34		<1.0	0.93	1.49	0.18		

¹⁾ Oko kukhokeleke koovimba beenkukacha zekwaliti yamanzi angaphantsi komhlaba bexesha elide (isuka kwi-DWS Water Management System). Inani (ubuncikane) lohlalutyo elisetyenzisiweyo kuphononongo lwamanani ngu-nine (9).

²⁾ umda ophuzulu wekwaliti yamanzi (okusela) yeHlelo II (isuka kwi-WRC et al. 2nd Edition, 1998, Volume 1: Assessment Guide), futhi

³⁾ ixabiso eliphakathi lidibene ne-10%. Apho umahluko kumaxabiso ekwaliti yamanzi ngokwenkangeleko yekwaliti yamanzi angaphantsi komhlaba neyeemfundo ezisisiseko zoluntu uye wafunyanwa, kulapho nexabiso lokhuselo elincinane okanye eliphezulu liye takhethelewa loo Mjelo uhlile wekwaliti yamanzi angaphantsi komhlaba. Apho ummandla wohlelo wathi abonisa inkangeleko yekwaliti yamanzi angaphantsi komhlaba, loo mmandla waye wakhethewa njengoMjelo wekwaliti yamanzi nexabiso lakhe lahiwa ukwenyuswa nge- 10 per cent kuphela xa nexabiso lingadlulanga kwele loMjelo we- BHN.

UTafale 6.4: Ushwankathelo lwehlelo lekwaliti yamanzi nemida ekhathazayo

Indawo yoboniselo ngamanzi yeminyaka emine	Ihlelo lekwaliti yamanzi (WRC, 1998)	imida ekhathazayo yekwaliti yamanzi
G40C	0	Ayikho
G40F	0	Ayikho
G40J	0	Ayikho
G40L	II	Sodium, Chloride
G40M	II	Chloride
G50E	I	Chloride
G50F	I	Chloride
G50H	III	Ukutsala umbane, Magnesium, Sodium, Chloride, Sulphate
G50J	II	Chloride, Sodium, ukutsala umbane
H10A	II	Sodium, Chloride, ukutsala umbane
H10B	0	Ayikho
H10C	I	Chloride
H10F	0	Ayikho
H10G	0	Ayikho
H10H	II	i-Chloride
H10L	0	Ayikho
H20A	0	Ayikho
H20B	0	Ayikho
H20D	0	Ayikho
H20E	0	Ayikho
H20F	0	Ayikho
H30A	II	iChloride
H30C	0	Ayikho
H30D	0	Ayikho
H40A	I	Ukutsala umbane, Calcium, iChloride
H40B	0	Ayikho
H40F	0	Ayikho
H70B	III	Ukutsala umbane, Magnesium, Sodium, Chloride
H70K	II	Chloride
H90D	II	Chloride
H90E	II	Ukutsala umbane, Sodium, Chloride
J11B	I	Ukutsala umbane
J11E	II	Ukutsala umbane, Chloride
J11G	II	Ukutsala umbane, Chloride
J12B	III	Ukutsala umbane, Magnesium, Sodium, Chloride
J12D	II	Ukutsala umbane, Sodium, Chloride
J12F	III	Ukutsala umbane, Sodium, Chloride
J12G	III	Ukutsala umbane, Sodium, Chloride
J12H	II	Ukutsala umbane, Chloride
J12K	III	Ukutsala umbane, Calcium, Sodium, Chloride, Sulphate
J12L	0	Ayikho
J13B	III	Ukutsala umbane, Sodium, Chloride
J21A	0	None
J21B	I	Ukutsala umbane, Sodium, Chloride
J21C	I	Ukutsala umbane
J21D	I	Ukutsala umbane, Sodium, Chloride
J21E	I	Ukutsala umbane
J22B	I	Ukutsala umbane
J22C	I	Ukutsala umbane, Sodium
J22D	I	Ukutsala umbane, Sodium, Chloride
J22E	I	Ukutsala umbane
J22F	I	Ukutsala umbane, Calcium, Chloride
J22H	II	Ukutsala umbane, Calcium, Sodium, Chloride
J22J	I	Ukutsala umbane
J23A	II	Ukutsala umbane, Sodium, Chloride, Sulphate
J23C	I	Ukutsala umbane, Calcium, Sodium, Chloride
J23D	II	Ukutsala umbane, Chloride
J23F	I	Ukutsala umbane, Sodium, Chloride

Indawo yoboniselwe ngamanzi yeminyaka emine	Ihlelo lekwalithi yamanzi (WRC, 1998)	imida ekhathazayo yekwalithi yamanzi
J24A	I	Ukutsala umbane
J24B	I	Ukutsala umbane
J24C	I	Ukutsala umbane, Sodium, Chloride
J24D	II	Ukutsala umbane, Chloride
J24E	II	Ukutsala umbane, Chloride
J25A	0	None
J25C	0	None
J31D	0	None
J32A	II	Ukutsala umbane, Chloride
J32B	II	Ukutsala umbane, Chloride
J32C	I	Ukutsala umbane, Calcium, Sodium, Chloride
J32E	0	Ayikho
J33D	0	Ayikho
J33E	0	Ayikho
J33F	0	Ayikho
J34A	0	Ayikho
J34B	II	Chloride
J34D	0	Ayikho
J34F	I	Chloride
J35A	0	Ayikho
J35B	I	Chloride
J35D	II	Ukutsala umbane, Chloride
J40D	II	Chloride, Sodium
J40E	II	Ukutsala umbane, Sodium, Chloride
K10A	III	Ukutsala umbane, Sodium, Chloride
K10B	II	Ukutsala umbane, Sodium, Chloride
K10D	III	Ukutsala umbane, Sodium, Chloride
K30B	0	Ayikho
K50B	I	Chloride

7. AMACHWEBA (ICANDELO LEKWALITI YAMANZI)

UTafale 7.1 Umthamo wamanzi

Indawo yoboniselo ngamanzi yeminyaka emine	Igama lechweba	I-PES	I-REC	nMAR (MCM)	EWR nMAR (MCM)
G40B	Rooiels	B	B	*9.44	n/a
G40D	Palmiet	C	B	*177.94	n/a
G40G	Bot/Kleinmond	C	B	*77.67	n/a
G40H	Onrus	E	D	*4.74	n/a
G40L	Klein	C	B	*51.21	n/a
G40M	Uilkraals	D	C	*6.82	n/a
G40F	Heuningnes	C	A	29.53	n/a
H70K	Breede	B	B	1785.00	954.00
H80E	Duiwenhoks	B	A	89.29	73.01
H90C	Goukou	C	B	115.95	91.73
J40B	Gouritz	C	B	623.52	377.23
K10A	Blinde	C	C	n/a	n/a
K10B	Hartenbos	D	C	n/a	n/a
K10F	Klein Brak	C	C	50.67	37.66
K20A	Groot Brak	D	C	36.79	11.11
K30A	Maalgate	B	C	41.51	24.41
K30B	Gwaing	B	C	35.07	21.7

Indawo yoboniselolngamanzi yeminyaka emine	Igama lechweba	I-PES	I-REC	nMAR (MCM)	EWR nMAR (MCM)
K30C	Kaaimans	B	B	53.6	41.3
K30D	Indawo yentlango: Touws	B	A	29.66	25.15
K40D	Swartvlei	B	B	83.4	56.6
K40E	Goukamma	B	A	57.5	48.8
K50B	Kynsna	B	B	83.2	63.4
K60 E and F	Keurbooms	A/B	A/B	232	214.10
K60G	Noetsie	B	B	4.8	n/a
K60G	Piesang	D	B/C	n/a	n/a
K70A	Groot (Wes)	B	A	n/a	n/a
K70A	Matjie	B	B	5.10	n/a
K70A	Sout	A	A	11.22	n/a
K70B	Bloukrans	A	A	n/a	n/a

* I-MAR yanamhla

IKWALITI YAMANZI ECHWEBE

UTafle 7.2: ii-EcoSpecs nomda wengxaki enokwehla kwiChweba iBlinde

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezinokwehla
I-Hydrology (Imeko yasemanzini)	Gcina imeko yokuhamba kwamanzi (iindawana zokuphila ezincinane zezona ziwadinga kakhulu amanzi ahambayo)	I-MAR ayiguquki ngaphezu kwe- 10% kuleyo yexesha langoku Iimpuphuma (ziboniswa yi- 1:10 yexesha lomnyaka) sukunciphisa ngaphezulu kwe- 5% kuleyo yexesha langoku Amanzi ahamba ezantsi akohluki ngaphezu kwe - 5% kulawo exesha langoku
I-Hydrodynamics (uguguququko lwasemanzini)	Gcina isimo somlomo wechweba sisihle ukuze lendawo yokuphilisana izilungele iintaka, iintlanzi, ii- macrophytes, i-microalgae (ubulembu obuncinanana) nekwaliti yamanzi	Isimo somlomo ovulekileyo wechweba Sanda/sincipha nge- 10% kweso sexesha sangoku Isimo somlomo wechweba sibuvaleka, siqhuba njalo sigalela amanzi elwandle. I-avareji yobunzulu bamanzi ngu <0.5 m (eli nani liza kuqinisekiswa luhlolo) Iqondo lokuguquka kwamanqana amanzi ngu > 30% kunelo lexesha langoku
Ikwali yamanzi	Ukusasazwa kwamanzi anetyuwa makungade kubangele ii-TPCs ezigqithileyo ezintlanzini, kwizinto eziphilayo ezingenamathambo, kwii- macrophytes nee-microalgae Ubukho bodaka ne-oksijini enyibilikisiweyo mabungade bubangele ii-TPCs ezigqithileyo kwii-biota Ubukho be-DIN/be-DIP mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes nee-microalgae Izinto ezinyehfu mazingade zibangele ii-TPCs ezigqithileyo kwii-biota	Iqondo lobukho betyuwa emanzini > 20 (uluhlu olulindelekileyo ngu-5-15) I-oksijini enyibilikisiweyo (DO) ngu< 5 mg/l ecwhebeni Ubukho bodaka > 10 NTU kumanzi ahamba kancinci Ubunzulu be-Secchi: ukuya ezantsi DIN >100 µg/l (i-avareji) DIP > 20 µg/l (i-avareji) Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAf, 1995) Iqondo lobukho beentlenge libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Uguquququko lweentlenge	Imeko yeempuphuma mayilandelee iipethini zosasazo lweentlenge nezeendawo zokuphilisana zasemanzini (zangaphakathi ngqo emjelweni) khonukuze ingade ibangele ii- TPCs ezigqithileyo ze-biota linguqu kubukhulu beepethini zosasazo lweenkoko zeentlenge mayingade ibangele ii- TPCs ezigqithileyo kwizinto eziphilayo ezingenamathambo zolwandle Inguqu kulwakihiwo lweentlenge (oluphakathi) neempawu Inguqu kwibathymetry e-avareji	I-avareji yokwakheka kweentlenge (i-% yamaqhezu) iphaya kwinguqu elindelekileyo echwebeni ukusuka kuleyo yesiqhelo (uza kuthatyathwa umlinganiselo wayo) ngo-30% (kuphando ngalunye) I-avareji yobunzulu phaya ngakwinguqu elindelekileyo kwelona-lona jelo ukusuka ku-30% waleyo yesiqhelo (iza kuqingqwa) (kwezi meko kulindeleke ukuba kubekho uguquququko olubalaseleyo kwi-bathymetry ephakathi kweempuphuma namaxesha ovaleleko awandisiweyo)
I-Microalgae (ubulembu obuncinanana)	Gcina ubukhulu bendalo bobulembu be-phytoplankton/bolwandle bebezezantsi /buphakathi Thintela ukuvela kwee-phytoplankton ezidubulayo	i-Phytoplankton ngu > 3.5 µg/l (phakathi) i-Benthic microalgae ngu >23 mg/m ² (phakathi) i-Phytoplankton ngu >20 µg/l /okanye ukushinyana kweeseli ngu >10 000 iiseli/ml (kanye nje)
Ii-Macrophytes	Gcina iindawo zokuphila ze-macrophyte zikwimeko entle. Thintela iingcongolo ukuze zinganabeli emanzini avulekileyo	Kukho inguqu kangange-20% kummandla we-macrophyte. (iingcongolo ziqgume ummandla kangange- 0.04 ha.)

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezinokwehla
	Thintela ukwanda kwezondlo nobulembu obudubulayo Thintela ukunaba kwemithi engeyoyomthonyama (njengomnga) kummandla wonxweme	Iingcongolo zithabatha ummandla kangange > 0.5 ha ii-Macro-algal ezidubulayo zigqume kangange > 50% yommandla ovulekileyo wamanzi ubukho bee-macrophytes ezitshabalalisayo zasemanzini, umzekelo i-Azolla, i-hyacinth yamanzi, njalo-njalo. Imithi etshabalalisayo yogqume i- > 50% yomda wonxweme
Izinto eziphilayo ezingenamathambo	Phanda ngobukho beproni zentlabathi ze-Callichirus kraussi kunxweme lwentlabathi olusechwebeni elisemazantsi Phanda ngobukho bee- Pseudodiaptomus hessei okanye ii- congeneric zechweba kwi- zooplankton yechweba	Ukuba imigqeku yangoku iphambuka kwimo yesiqhelo (njengoko kufunyanisiwe kula matyelelo mathathu okuqala) ngaphaya kwe-30%
Iintlanzi	Iimigqeku yeentlanzi mayi- amabakala anxulunyaniswa nechweba amahlano nabonisa iimpawu ezifanayo (ngokwahluka nangobuninzi) kwezo ziphantsi kohlo. Ngokwamanani, ubugqeku mabuquke oku kulandelayo: Abahlali basechwebeni (i-50-80% lobuninzi babo bedibene) i-Ib yemigqeku yasemanzini nasechwebeni (i-5-20%) i-IIa yaleyo impilo yayo ixhomekeke echwebeni (i-10-20%) i-IIb yemigqeku enxulunyaniswa nechweba (i-5-15%), i-IIc yezo zixhamla emanzini (i-20-80%) i-III yezibhadubhadu zasemanzini (ezingekho ngaphezulu kwe- 5%) i-IV yeentlanzi zomthonyama (i-1-5%) i-V yeentlobo ze-catadromous (i-1-5%) Iintlobo zamabakala mazibe nemigqeku ezinzileyo eziintlobo ezingekho ngaphantsi kwesibini (umzekelo i-G.aestuaria & ne-Hyporampus capensis). Amabakala aloo migqeku ampilo yayo ixhomekeke echwebeni makabenabameli abaziintlobo ezisetyenziswa kakhulu ezingekho ngaphantsi kwesibini (umzekelo ii- L. lithognathus, nee-Lichia amia). Iintlobo ze-REI (zasekudibaneni komlambo nechweba (i-River Estuary Interface) ezonganyelwe zizo zombini ii- Myxus capensis nee-G. aestuaria.	i-Ia yemigqeku yasechwebeni ngu<50% i-Ib yemigqeku yasemanzini nasechwebeni ngu- <10% i-IIa yemigqeku empilo yayo ixhomekeke echwebeni ngu <10% i-IIb yemigqeku ebandakanyeka echwebeni ngu <5% i-IIc yaloo migqeku ixhamla emanzini ngu < 20% i-III yezibhadu-bhadu zasemanzini ngu > 5% i-IV yeentlanzi zomthonyama ngu <1% i-V yezo ntlobo zifudukayo ukusuka emanzini ziye elwandle ukuya kuzala ngu <1%
Iintaka	Gcina imigqeku yamaqela okuqala eentaka ezikhoyo echwebeni.	Amanani eentaka kulo naliphi na iqela (ngaphandle kwezo ntlobo zanda ngokwengingqi, njengedada laseYiphutha) ehla aye kutsho ngaphantsi kwemeko ephakathi yesiqhelo (oku kuqinisekiswa ziinkcukacha zexesha elidlulileyo okanye uphando lwakuqala/lwangaphambili) Amanani eentlobo ezithile / eentaka ezibalwa izihlandlo ezithathu ezilandelelanayo zehlobo okanye zobusika

UTafule 7.3: ii-EcoSpecs nemida yeengxaki ezisenokwehla kwichweba iHartenbos

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
I-Hydrology (Imeko yasemanzini)	Gcina ubuncikane isimo sexesha langoku sokuhamba kwamanzi (iza kuqinisekiswa)	I-MAR ayiguquki ngaphezu kwe- 10% kuleyo yexesha langoku Iimpuphuma (ziboniswa yi- 1:10 yexesha lomnyaka) sukunciphisa ngaphezulu kwe- 5% kuleyo yexesha langoku Amanzi ahamba ezantsi akohluki ngaphezu kwe - 50% kulawo exesha langoku
I-Hydrodynamics (uguquguquko lwasemanzini)	Gcina isimo somlomo wechweba sisihle ukuze lendawo yokuphilisana izilungele iintaka, iintlanzi, ii- macrophytes, i-micro-algae (ubulembu obuncinanana) nekwali yamanzi	Isimo somlomo ovalekileyo wechweba asinciphi nge- 10% kweso sexesha sangoku I-avareji yobunzulu bamanzi ngu < 10% kunaleyo yexesha langoku Iqondo lokudlanyaza kwamaza (xa umlomo wechweba uvulekile) ngu > 20%
Ikwali yamanzi	Ukusasazwa kwamanzi anetyuwa makungade kubangele ii-TPCs ezigqithileyo ezintlanzini, kwizinto eziphilayo ezingenamathambo, kwii-macrophytes nee-microalgae Ubukho bodaka ne-oksijini enyibilikisiweyo mabungade bubangele ii-TPCs ezigqithileyo kwii-biota Ubukho be-DIN/be-DIP mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes nee-microalgae Izinto ezinyehfu mazingade zibangele ii-TPCs ezigqithileyo kwii-biota	I-avareji yobukho betyuwa emanzini ngasechwebeni incipha ngo> 5 ngaphantsi kwe-avareji yesiqhelo (iza kuqingqwa) I-oksijini enyibilikisiweyo (DO) ngu < 5 mg/l ecwhebeni Ubukho bodaka ngu > 20 NTU kumanzi ahamba kancinci i-Secchi kwiindawo ezibuhlaziyeka ngu: 0.5 m DIN >200 µg/l i-avareji (iza kuqinisekiswa) DIP > 50 µg/l i-avareji (iza kuqinisekiswa) Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAf, 1995) Iqondo lobukho beentlenge libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi-Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Uguquguquko lweentlenge	Imeko yeempuphuma mayilandelee iipethini zosasazo lweentlenge nezeendawo zokuphilisana zasemanzini (zangaphakathi ngqo emjelweni) khonukuze ingade ibangele ii- TPCs ezigqithileyo ze-biota linguqu kubukhulu beepethini zosasazo lweenkoko zeentlenge mayingade ibangele ii- TPCs ezigqithileyo kwizinto eziphilayo ezingenamathambo zolwandle Inguqu kulwakhawo lweentlenge (oluphakathi) neempawu Inguqu kwibathymetry e-avareji	I-avareji yokwakheka kweentlenge (i-% yamaqhezu) iphaya kwinguqu elindelekileyo echwebeni ukusuka kuleyo yesiqhelo (uza kuthatyathwa umlinganiselo wayo) ngo-30% (kuphando ngalunye) I-avareji yobunzulu phaya ngakwinguqu elindelekileyo kwelona-lona jelo ukusuka ku-30% waleyo yesiqhelo (iza kuqingqwa) (kwezi meko kulindeleke ukuba kubekho uguquguquko olubalaseleyo kwi-bathymetry ephakathi kweempuphuma namaxesha ovaleleko awandisiweyo)
I-Microalgae (ubulembu obuncinanana)	Gcina ubukhulu bendalo bobulembu be-phytoplankton/bolwandle /buphakathi Thintela ukuvela kwee-phytoplankton ezidubulayo	i-Phytoplankton ngu > 8 µg/l (phakathi) i-Benthic microalgae >42 mg/m ² (phakathi) i-Phytoplankton ngu >20 µg/l /okanye ukushinyana kweeseli ngun>10 000 iiseli/ml (kanye nje) i-Dinoflagellates, i-chlorophytes ne/okanye i-cyanobacteria ngu >10% yobuninzi obubalulekayo

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
Ii-Macrophytes	<p>Gcina iindawo zokuphila ze-macrophyte zikwimeko entle.</p> <p>Thintela iingcongolo ukuze zinganabeli emanzini avulekileyo</p> <p>Thintela ukwanda kwezondlo nobulembu obudubulayo</p> <p>Thintela ukunaba kwemithi engeyoyomthonyama (njengomnga) kummandla wonxweme</p> <p>Gcina imfezeko yomgxobhozo weetyuwa</p>	<p>Kukho inguqu kangange-20% kummandla we-macrophyte. (iingcongolo ziqgume ummandla kangange- 9 ha, umgxobhozo weetyuwa kangange-47 ha)</p> <p>ii-Macroalgal ezidubulayo zithabatha i- > 50% yommandla wamanzi avulekileyo</p> <p>ii-Macro-algal ezidubulayo ziqgume kangange > 50% yommandla ovulekileyo wamanzi</p> <p>Ubukho bee-macrophytes ezitshabalalisayo zasemanzini (umzekelo, i- <i>Azolla</i>, i-water hyacinth)</p> <p>Bugquma i- > 10% yethafa leempuphuma</p> <p>Iqondo lokwanda kwemimandla engenayo imigxobhozo yeetyuwa ngenxa yokuncipha kobumanzi nokunyuka kwamaqondo obukho beetyuwa emanzini yi- >30% yomgxobhozo weetyuwa</p>
Izinto eziphilayo ezingenamathambo	<p>Phanda ngobukho beeproni zentlabathi ze-<i>Callichirus kraussi</i> kunxweme lwentlabathi olusechwebeni elisemazantsi</p> <p>Phanda ngobukho bee- <i>Pseudodiptomus hessei</i> okanye ii- congeneric zechweba kwi- zooplankton yechweba</p>	<p>Ukuba ngaba iintlobo zexesha langoku ziphambuka kwi-avareji yesiqhelo (njengoko utyelelo lwezihlandlo ezithathu lubonisile) malunga nangaphaya kwe-30%</p>

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
iintlanzi	<p>Iimigqeku yeentlanzi mayiquke amabakala anxulunyaniswa nechweba amahlani nabonisa iimpawu ezifanayo (ngokwahluka nangobuninzi) kwezo ziphantsi kohlolo. Ngokwamanani, ubugqeku mabuquke oku kulandelayo:</p> <p>Abahlali basechwebeni (i-20-60% lobuninzi babo bedibene)</p> <p>i-Ib yemigqeku yasemanzini nasechwebeni (i-10-30%)</p> <p>i-IIa yaleyo impilo yayo ixhomekeke echwebeni (i-20-40%)</p> <p>i-IIb yemigqeku enxulunyaniswa nechweba (i-5- 20%),</p> <p>i-IIc yezo zixhamla emanzini (i-20-80%)</p> <p>i-III yezibhadubhadu zasemanzini (ezingekho ngaphezulu kwe- 5%)</p> <p>i-IV yeentlanzi zomthonyama (i-1-5%)</p> <p>i-V yeentlobo ze-catadromous (i-1-5%)</p> <p>Iintlobo zamabakala mazibe nemigqeku ezinzileyo eziintlobo ezingekho ngaphantsi kwesibini (umzekelo i- <i>G.aestuaria</i>, i-<i>Hyporhamphus capensis</i>, <i>Omobranchus woodii</i>).</p> <p>Amabakala aloo migqeku ampilo yayo ixhomekeke echwebeni makabenabameli abaziintlobo ezisetyenziswa kakhulu ezingekho ngaphantsi kwesibini (umzekelo, i- <i>A. japonicus</i>, i-<i>L. lithognathus</i>, i-<i>P. commersonii</i>, i-<i>Lichia amia</i>).</p> <p>Iintlobo ze-REI (zasekudibaneni komlambo nechweba (i-River Estuary Interface) ezonganyelwe zizo zombini ii- <i>Myxus capensis</i> nee-<i>G. aestuaria</i>.</p>	<p>i-Ia yemigqeku yasechwebeni ngu <20%</p> <p>i-Ib yemigqeku yasemanzini nasechwebeni ngu <10%</p> <p>i-IIa yemigqeku empilo yawo ixhomekeke echwebeni ngu <20%</p> <p>i-IIb yemigqeku ebandakanyeka echwebeni ngu <5%</p> <p>i-IIc yaloo migqeku ixhamla emanzini ngu < 20%</p> <p>i-IV yeentlanzi zomthonyama ngu <1%</p> <p>i-V yezo ntlobo zifudukayo ukusuka emanzini ziye elwandle ukuya kuzala ngu <1%</p> <p>i-Ia imelwe ngu- <i>G. aestuaria</i> kuphela</p> <p>i-IIa yeentlanzi ezixhatshazwayo ikumanani asezantsi kakhulu okanye ayikho</p> <p>iintlobo ze-REI zimelwe ngu- <i>G. aestuaria</i> kuphela. <i>Ii-aestuaria</i>, nee-<i>Myxus capensis</i> azikho</p>
Iintaka	Gcina imigqeku yamaqela okuqala eentaka ezikhoyo echwebeni.	<p>Amanani eentaka kulo naliphi na iqela (ngaphandle kwezo ntlobo zanda ngokwengingqi, njengedada laseYiphutha) ehla aye kutsho ngaphantsi kwemeko ephakathi yesiqhelo (oku kuqinisekiswa ziinkcukacha zexesha elidlulileyo okanye uphando lwakuqala/lwangaphambili)</p> <p>Amanani eentlobo ezithile / eentaka ezibalwa izihlandlo ezithathu ezilandelelanayo zehlobo okanye zobusika</p>

UTafle 7.4: ii-Eco-Specs nemida yeengxaki ezisenokwehla kwichweba iPiesang

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
I-Hydrology (Imeko yasemanzini)	Gcina ubuncikane isimo sexesha langoku sokuhamba kwamanzi (iza kuqinisekiswa)	I-MAR ayiguquki ngaphezu kwe- 10% kuleyo yexesha langoku Iimpuphuma (ziboniswa yi- 1:10 yexesha lomnyaka) azinciphi ngaphezulu kwe- 5% kwezo zexesha langoku Amanzi ahamba ezantsi akohluki ngaphezu kwe - 50% kulawo exesha langoku
I-Hydrodynamics (uguquguquko lwasemanzini)	Gcina isimo somlomo wechweba sisihle ukuze lendawo yokuphilisana izilungele iintaka, iintlanzi, ii- macrophytes, i-microalgae (ubulembu obuncinanana) nekwaliti yamanzi	Isimo somlomo ovalekileyo wechweba asinciphi nge-10% kweso sexesha sangoku I-avareji yobunzulu bamanzi ngu < 10% kunaleyo yexesha langoku Iqondo lokudlanyaza kwamaza (xa umlomo wechweba uvulekile) ngu > 20%
Ikwali yamanzi	Ukusasazwa kwamanzi anetyuwa makungade kubangele ii-TPCs ezigqithileyo ezintlanzini, kwizinto eziphilayo ezingenamathambo, kwii-macrophytes nee-microalgae Ubukho bodaka ne-oksijini enyibilikisiweyo mabungade bubangele ii-TPCs ezigqithileyo kwii-biota Ubukho be-DIN/be-DIP mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes nee-microalgae Izinto ezinetyhefu mazingade zibangele ii-TPCs ezigqithileyo kwii-biota	Iqondo lobukho betyuwa emanzini ngu > 20 (uluhlu olulindelekileyo ngu10-20) Iqondo lobukho betyuwa emanzini ngu < 5 (uluhlu olulindelekileyo ngu10-20) I-oksijini enyibilikisiweyo (DO) ngu< 5 mg/l ecwhebeni Ubukho bodaka ngu > 10 NTU kumanzi ahamba kancinci i-Secchi: iya ezantsi DIN >100 µg/l kanye nje DIP > 20 µg/l kanye nje Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) Iqondo lobukho beentlenga libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Uguquguquko kweentlenga	Imeko yeempuphuma mayilandelee iipethini zosasazo lweentlenga nezeendawo zokuphilisana zasemanzini (zangaphakathi ngqo emjelweni) khonukuze ingade ibangele ii- TPCs ezigqithileyo ze-biota linguqu kubukhulu beepethini zosasazo lweenkoko zeentlenga mayingade ibangele ii- TPCs ezigqithileyo kwizinto eziphilayo ezingenamathambo zolwandle Inguqu kulwakiwo lweentlenga (oluphakathi) neempawu Inguqu kwibathymetry e-avareji	I-avareji yokwakheka kweentlenga (i-% yamaqhezu) iphaya kwinguqu elindelekileyo echwebeni ukusuka kuleyo yesiqhelo (uza kuthatyathwa umlinganiselo wayo) ngo-30% (kuphando ngalunye) I-avareji yobunzulu phaya ngakwinguqu elindelekileyo kwelona-lona jelo ukusuka ku-30% waleyo yesiqhelo (iza kuqingqwa) (kwezi meko kulindeleke ukuba kubekho uguquguquko olubalaseleyo kwi-bathymetry ephakathi kweempuphuma namaxesha ovaleleko awandisiweyo)
I-Microalgae (ubulembu obuncinanana)	Gcina ubukhulu bendalo bobulembu be-phytoplankton/bolwandle buphakathi Thintela ukuvela kwee-phytoplankton ezidubulayo	i-Phytoplankton ngu > 3.5 µg/l (phakathi) i-Benthic microalgae >11 mg/m ² (phakathi) i-Phytoplankton ngu >20 µg/l /okanye ukushinyana kweeseli ngun>10 000 iiseli/ml (kanye nje)
Ii-Macrophytes	Gcina iindawo zokuphila ze-macrophyte zikwimeko entle. Thintela ukwanda kwezondlo eziza kudala ubulembu obudubulayo Beka iliso kwizityalo ezitshabalalisayo kumda wonxweme	Inguqu engaphezu kwe-20 % kummandla owogqunywe zii-macrophytes (iingcongolo neentlenga ziqquma ummandla oyi-3.14 ha, ii-macrophytes ezigubungelweyo nemigxobhozo yeetyuwa ekhoyo lityalo ezitshabalalisayo zogqume malunga ne- >5% yeendawo zokuphila zizonke

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
Izinto eziphilayo ezingenamathambo	Phanda ngobukho beeproni zentlabathi ze- <i>Callichirus kraussi</i> kunxweme lwentlabathi olusechwebeni elisemazantsi Phanda ngobukho bee- <i>Pseudodiptomus hessei</i> okanye ii- congeneric zechweba kwi-zooplankton yechweba	Imigqeku iphambuka kwi-avareji yesiqhelo (njengo uphando lwezihlandlo ezintathu lubonisa) malunga nangaphaya kwe- 30%
Iintlanzi	Iimigqeku yeentlanzi mayiuke amabakala anxulunyaniswa nechweba amahlanu nabonisa iimpawu ezifanayo (ngokwahluka nangobuninzi) kwezo ziphantsi kohlo. Ngokwamanani, ubugqeku mabuquke oku kulandelayo: Imigqeku yasechwebeni (i-50-80% lobuninzi babo bedibene) i-Ib yemigqeku yasemanzini nasechwebeni (i-5-20%) i-Ila yaleyo impilo yayo ixhomekeke echwebeni (i-10-20%) i-Ilb yemigqeku enxulunyaniswa nechweba (i-5- 15%), i-Ilc yezo zixhamla emanzini (i-20-80%) i-III yezibhadubhadu zasemanzini (ezingekho ngaphezulu kwe- 5%) i-IV yeentlanzi zomthonyama (i-1-5%) i-V yeentlobo ze-catastromous (ezo zifudukela elwandle ukuya kuzala (i-1-5%) Iintlobo zamabakala mazibe nemigqeku ezinzileyo eziintlobo ezingekho ngaphantsi kwesibini (umzekelo i- <i>G.aestuaria</i> , i- <i>Hyporampus capensis</i>). Amabakala aloo migqeku ampilo yayo ixhomekeke echwebeni makabenabameli abaziintlobo ezisetyenziswa kakhulu ezingekho ngaphantsi kwesibini (umzekelo, i- <i>L. lithognathus</i> , , i- <i>Lichia amia</i>). Iintlobo ze-REI (zasekudibaneni komlambo nechweba (i-River Estuary Interface) ezonganyelwe zizo zombini ii- <i>Myxus capensis</i> nee- <i>G. aestuaria</i> .	i-Ia yemigqeku yasechwebeni ngu <50% i-Ib yemigqeku yasemanzini nasechwebeni ngu- <10% i-Ila yemigqeku empilo yawo ixhomekeke echwebeni ngu <10% i-Ilb yemigqeku ebandakanyeka echwebeni ngu <5% i-Ilc yaloo migqeku ixhamla emanzini ngu < 20% i-III yezibhadu-bhadu zasemanzini ngu <5% i-IV yeentlanzi zomthonyama ngu <1% i-V yezo ntlobo zifudukayo ukusuka emanzini ziye elwandle ukuya kuzala ngu <1%
Iintaka	Gcina imigqeku yeqela lokuqala leentaka ezilapho echwebeni.	Amanani eentaka kulo naliphi na iqela (ngaphandle kwezo ntlobo zanda ngokwengingqi, njengedada laseYiphutha) ehla aye kutsho ngaphantsi kwemeko ephakathi yesiqhelo (oku kuqinisekiswa ziinkcukacha zexesha elidlulileyo okanye uphando lwakuqala/lwangaphambili) Amanani eentlobo ezithile / eentaka ezibalwa izihlandlo ezithathu ezilandelelanayo zehlobo okanye zobusika

UTafle 7.5: ii-EcoSpecs kwichweba iGroot (Wes)

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
I-Hydrology (Imeko yasemanzini)	Gcina ubuncikane isimo sexesha langoku sokuhamba kwamanzi (iza kuqinisekiswa)	I-MAR ayiguquki ngaphezu kwe- 10% kuleyo yexesha langoku limpuphuma (ziboniswa yi- 1:10 yexesha lomnyaka) azinciphi ngaphezulu kwe- 5% kwezo zexesha langoku Amanzi ahamba ezantsi akohluki ngaphezu kwe - 50% kulawo exesha langoku
I-Hydrodynamics (uguquguquko lwasemanzini)	Gcina isimo somlomo wechweba sisihle ukuze lendawo yokuphilisana izilungele iintaka, iintlanzi, ii- macrophytes, i-micro-algae (ubulembu obuncinanana) nekwaliiti yamanzi	Isimo somlomo ovalekileyo wechweba asinciphi nge- 10% kweso sexesha sangoku I-avareji yobunzulu bamanzi ngu < 10% kunaleyo yexesha langoku Iqondo lokudlanyaza kwamaza (xa umlomo wechweba uvulekile) ngu > 20%
Ikwaliiti yamanzi	Ukusasazwa kwamanzi anetyuwa makungade kubangele ii-TPCs ezigqithileyo ezintlanzini, kwizinto eziphilayo ezingenamathambo, kwii- macrophytes nee-microalgae Ubukho bodaka ne-oksijini enyibilikisiweyo mabungade bubangele ii-TPCs ezigqithileyo kwii-biota Ubukho be-DIN/be-DIP) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes nee-microalgae Izinto ezinyehfu mazingade zibangele ii-TPCs ezigqithileyo kwii-biota	Iqondo lobukho betyuwa ngasechwebeni lincipha ngo-5 ngaphantsi kwe-avareje yesiqhelo 9iza kuqingqwa) Iqondo lobukho betyuwa emanzini ngu < 10 kumphezulu wechweba (uluhlu olulindelekileyo ngu5-10 ikakhulu kuyo yonke le ndawo) I-oksijini enyibilikisiweyo (DO) ngu< 5 mg/l ecwhebeni Ubukho bodaka ngu > 10 NTU kumanzi ahamba kancinci i-Secchi: iya ezantsi DIN >100 µg/l kanye nje DIP > 20 µg/l kanye nje Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) Iqondo lobukho beentlenge libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi-Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Uguquguquko kweentlenge	Imeko yeempuphuma mayilandelee iipethini zosasazo lweentlenge nezeendawo zokuphilisana zasemanzini (zangaphakathi ngqo emjelweni) khonukuze ingade ibangele ii- TPCs ezigqithileyo ze-biota linguqu kubukhulu beepethini zosasazo lweenkoko zeentlenge mayingade ibangele ii- TPCs ezigqithileyo kwizinto eziphilayo ezingenamathambo zolwandle Inguqu kulwakihiwo lweentlenge (oluphakathi) neempawu Inguqu kwibathymetry e-avareji	I-avareji yokwakheka kweentlenge (i-% yamaqhezu) iphaya kwinguqu elindelekileyo echwebeni ukusuka kuleyo yesiqhelo (uza kuthatyathwa umlinganiselo wayo) ngo-30% (kuphando ngalunye) I-avareji yobunzulu phaya ngakwinguqu elindelekileyo kwelona-lona jelo ukusuka ku-30% waleyo yesiqhelo (iza kuqingqwa) (kwezi meko kulindeleke ukuba kubekho uguquguquko olubalaseleyo kwi-bathymetry ephakathi kweempuphuma namaxesha ovaleleko awandisiweyo)

CONTINUES ON PAGE 386 OF BOOK 4

Printed by and obtainable from the Government Printer, Bosman Street, Private Bag X85, Pretoria, 0001
Contact Centre Tel: 012-748 6200. eMail: info.egazette@gpw.gov.za
Publications: Tel: (012) 748 6053, 748 6061, 748 6065



Government Gazette Staatskoerant

REPUBLIC OF SOUTH AFRICA
REPUBLIEK VAN SUID AFRIKA

Vol. 686

26

August
Augustus

2022

No. 46798

PART 4 OF 4

N.B. The Government Printing Works will not be held responsible for the quality of "Hard Copies" or "Electronic Files" submitted for publication purposes

ISSN 1682-5845



9 771682 584003



AIDS HELPLINE: 0800-0123-22 Prevention is the cure

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
I-Microalgae (ubulembu obuncinanana)	Gcina ubukhulu bendalo bobulembu be-phytoplankton/bolwandle buphakathi Thintela ukuvela kwee-phytoplankton ezidubulayo	i-Phytoplankton ngu > 3.5 µg/l (phakathi) i-Benthic microalgae >11 mg/m ² (phakathi) i-Phytoplankton ngu >20 µg/l /okanye ukushinyana kweeseli ngun>10 000 iiseli/ml (kanye nje)
Ii-Macrophytes	Gcina iindawo zokuphila ze-macrophyte zikwimeko entle. Thintela ukwanda kwezondlo eziya kudala ubulembu obudubulayo Beka iliso kwizityalo ezitshabalalisayo kumda wonxweme	Inguqu engaphezu kwe-20 % kummandla owogqunywe zii-macrophytes (iingcongolo neentlenga ziqguma ummandla oyi-2.54 ha, nemigxobhozo yeetyuwa eyi-0.76 ha Iimacro-algal ezidubulayo ziqguma kangange- > 50% yommandla wamanzi avulekileyo kwezo meko zomlomo wechweba ovalekileyo Izityalo ezitshabalalisayo zogqume malunga ne->5% yeendawo zokuphila zizonke
Izinto eziphilayo ezingenamathambo	Phanda ngobukho beeproni zentlabathi ze-Callichirus kraussi kunxweme lwentlabathi olusechwebeni elisemazantsi Phanda ngobukho bee- copepod Pseudodiaptomus hessei okanye ii-congeneric zechweba kwi- zooplankton yechweba	Imigqeku iphambuka kwi-avareji yesiqhelo (njengo uphando lwezihlandlo ezintathu lubonisa) malunga nangaphaya kwe- 30%

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
lintlanzi	<p>Iimigqeku yeentlanzi mayi iquke amabakala anxulunyaniswa nechweba amahlanu nabonisa iimpawu ezifanayo (ngokwahluka nangobuninzi) kwezo ziphantsi kohlolo. Ngokwamanani, ubugqeku mabuquke oku kulandelayo:</p> <p>Iimigqeku yasechwebeni (i-50-80% lobuninzi babo bedibene)</p> <p>i-Ib yemigqeku yasemanzini nasechwebeni (i-5-20%)</p> <p>i-Ila yaleyo impilo yayo ixhomekeke echwebeni (i-10-20%)</p> <p>i-Ilb yemigqeku enxulunyaniswa nechweba (i-5- 15%),</p> <p>i-Ilc yezo zixhamla emanzini (i-20-80%)</p> <p>i-III yezibhadubhadu zasemanzini (ezingekho ngaphezulu kwe- 5%)</p> <p>i-IV yeentlanzi zomthonyama (i-1-5%)</p> <p>i-V yeentlobo ze-catadromous (ezo zifudukela elwandle ukuya kuzala (i-1-5%)</p> <p>lintlobo zamabakala mazibe nemigqeku ezinzileyo eziintlobo ezingekho ngaphantsi kwesibini (umzekelo i- <i>G.aestuaria</i>, i- <i>Hyporampus capensis</i>).</p> <p>Amabakala aloo migqeku ampilo yayo ixhomekeke echwebeni makabenabameli abaziintlobo ezisetyenziswa kakhulu ezingekho ngaphantsi kwesibini (umzekelo, i-<i>L. lithognathus</i>, i-<i>Lichia amia</i>).</p> <p>lintlobo ze-REI (zasekudibaneni komlambo nechweba (i-River Estuary Interface) ezonganyelwe zizo zombini ii- <i>Myxus capensis</i> nee-<i>G. aestuaria</i>.</p>	<p>i-Ia yemigqeku yasechwebeni ngu <50%</p> <p>i-Ib yemigqeku yasemanzini nasechwebeni ngu <10%</p> <p>i-Ila yemigqeku empilo yawo ixhomekeke echwebeni ngu <10%</p> <p>i-Ilb yemigqeku ebandakanyeka echwebeni ngu <5%</p> <p>i-Ilc yaloo migqeku ixhamla emanzini ngu < 20%</p> <p>i-III yezibhadu-bhadu zasemanzini ngu <5%</p> <p>i-IV yeentlanzi zomthonyama ngu <1%</p> <p>i-V yezo ntlobo zifudukayo ukusuka emanzini ziye elwandle ukuya kuzala ngu <1%</p>
lintaka	Gcina imigqeku yeqela lokuqala leentaka ezilapho echwebeni	<p>Amanani eentaka kulo naliphi na iqela (ngaphandle kwezo ntlobo zanda ngokwengingqi, njengedada laseYiphutha) ehla aye kutsho ngaphantsi kwemeko ephakathi yesiqhelo (oku kuqinisekiswa ziinkcukacha zexesha elidlulileyo okanye uphando lwakuqala/lwangaphambili)</p> <p>Amanani eentlobo ezithile / eentaka ezibalwa izihlandlo ezithathu ezilandelelanayo zehlobo okanye zobusika</p>

UTafale 7.6: ii-EcoSpecs nemida yeengxaki ezisenokwehla kwichweba iBloukrans

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
I-Hydrology (Imeko yasemanzini)	Gcina ubuncikane isimo sexesha langoku sokuhamba kwamanzi	yahluka malunga nangaphaya kwe- 10% ye-MAR
I-Hydrodynamics (uguquguquko lwasemanzini)	Gcina isimo somlomo wechweba sisihle ukuze lendawo yokuphilisana izilungele iintaka, iintlanzi, ii- macrophytes, i-microalgae (ubulembu obuncinanana) nekwaliti yamanzi	Umlomo wechweba uyavala
Ikwaliti yamanzi	Ukusasazwa kwamanzi anetyuwa makungade kubangele ii-TPCs ezigqithileyo ezintlanzini, kwizinto eziphilayo ezingenamathambo, kwii- macrophytes nee-microalgae Ubukho bodaka ne-oksijini enyibilikisiweyo mabungade bubangele ii-TPCs ezigqithileyo kwii-biota Ubukho be-DIN/be-DIP) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes nee-microalgae Izinto ezinetyhefu mazingade zibangele ii-TPCs ezigqithileyo kwii-biota	Iqondo lobukho betyuwa emanzini ngu < 10 kumphezulu wechweba (uluhu olulindelekileyo ngu >30 ikakhulu kuyo yonke le ndawo) I-oksijini enyibilikisiweyo (DO) ngu< 5 mg/l ecwhebeni Ubukho bodaka ngu > 10 NTU kumanzi ahamba kancinci i-Secchi: iya ezantsi DIN >100 µg/l kanye nje DIP > 20 µg/l kanye nje Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAf, 1995) Iqondo lobukho beentlenge libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi-Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
Uguquguquko lweentlenge	Imeko yeempuphuma mayilandele ipethini zosasazo lweentlenge nezeendawo zokuphilisana zasemanzini (zangaphakathi ngqo emjelweni) khonukuze ingade ibangele ii- TPCs ezigqithileyo ze-biota linguqu kubukhulu beepethini zosasazo lweenkoko zeentlenge mayingade ibangele ii- TPCs ezigqithileyo kwizinto eziphilayo ezingenamathambo zolwandle Inguqu kulwakiwo lweentlenge (oluphakathi) neempawu Inguqu kwibathymetry e-avareji	I-avareji yokwakheka kweentlenge (i-% yamaqhezu) iphaya kwinguqu elindelekileyo echwebeni ukusuka kuleyo yesiqhelo (uza kuthatyathwa umlinganiselo wayo) ngo-30% (kuphando ngalunye) I-avareji yobunzulu phaya ngakwinguqu elindelekileyo kwelona-lona jelo ukusuka ku-30% waleyo yesiqhelo (iza kuqingqwa) (kwezi meko kulindeleke ukuba kubekho uguquguquko olubalaseleyo kwi-bathymetry ephakathi kweempuphuma namaxesha ovaleleko awandisiweyo)
I-Microalgae (ubulembu obuncinanana)	Gcina ubukhulu bendalo bobulembu be-phytoplankton/bolwandle buphakathi Thintela ukuvula kwee-phytoplankton ezidubulayo	i-Phytoplankton ngu > 1.0 µg/l (phakathi) i-Benthic microalgae >11 mg/m ² (phakathi) i-Phytoplankton ngu >20 µg/l /okanye ukushinyana kweeseli ngun>10 000 iiseli/ml (kanye nje)
Ii-Macrophytes	Iindawo zokuphila ezisechwebeni zinamanxweme entlabathi/odaka kuphela (kangange-0.63 ha) nemijelo (kangange-2.88 ha), azikho ii-macrophytes	N/A

Icandelo le-ikholoji (lofundo ngendawo yokuphilisana nonxulumano lwezinto eziphila kuloo ndawo)	ii-EcoSpecs	Imida yeengxaki ezisenokwehla
Izinto eziphilayo ezingenamathambo	Phanda ngobukho beepironi zentlabathi ze-Callichirus kraussi kunxweme lwentlabathi olusechwebeni elisemazantsi Phanda ngobukho bee- copepod Pseudodiptomus hessei okanye ii-congeneric zechweba kwi- zooplankton yechweba	Ukuba imigqeku iphambuka kwi-avareji yesiqhelo (njengoko uphando lwezihlandlo ezintathu lubonisa) malunga nangaphaya kwe-30%
Iintlanzi	Iimigqeku yeentlanzi mayiquke amabakala anxulunyaniswa nechweba amahlanu nabonisa iimpawu ezifanayo (ngokwahluka nangobuninzi) kwezo ziphantsi kohlo. Ngokwamanani, ubugqeku mabuquke oku kulandelayo: Iimigqeku yasechwebeni (i-50-80% lobuninzi babo bedibene) i-Ib yemigqeku yasemanzini nasechwebeni (i-5-20%) i-Ila yaleyo impilo yayo ixhomekeke echwebeni (i-10-20%) i-Ilb yemigqeku enxulunyaniswa nechweba (i-5- 15%), i-Ilc yezo zixhamla emanzini (i-20-80%) i-III yezibhadubhadu zasemanzini (ezingekho ngaphezulu kwe- 5%) i-IV yeentlanzi zomthonyama (i-1-5%) i-V yeentlobo ze-catadromous (ezo zifudukela elwandle ukuya kuzala (i-1-5%) Iintlobo zamabakala mazibe nemigqeku ezinzileyo eziintlobo ezingekho ngaphantsi kwesibini (umzekelo i- <i>G. aestuaria</i> , i- <i>Hyporamphus capensis</i> , <i>Omobranchus woodii</i>). Amabakala aloo migqeku ampilo yayo ixhomekeke echwebeni makabenabameli abaziintlobo ezisetyenziswa kakhulu ezingekho ngaphantsi kwesibini (umzekelo, i- <i>L. lithognathus</i> , <i>P. commersonii</i> , i- <i>Lichia amia</i>). Iintlobo ze-REI (zasekudibaneni komlambo nechweba (i-River Estuary Interface) ezonganyelwe zizo zombini ii- <i>Myxus capensis</i> nee- <i>G. aestuaria</i> .	i-Ia yemigqeku yasechwebeni ngu <50% i-Ib yemigqeku yasemanzini nasechwebeni ngu- <10% i-Ila yemigqeku empilo yawo ixhomekeke echwebeni ngu <10% i-Ilb yemigqeku ebandakanyeka echwebeni ngu <5% i-Ilc yaloo migqeku ixhamla emanzini ngu < 20% i-III yezibhadu-bhadu zasemanzini ngu <5% i-IV yeentlanzi zomthonyama ngu <1% i-V yezo ntlobo zifudukayo ukusuka emanzini ziye elwandle ukuya kuzala ngu <1%
Iintaka	Gcina imigqeku yeqela lokuqala leentaka ezilapho echwebeni	Amanani eentaka kulo naliphi na iqela (ngaphandle kwezo ntlobo zanda ngokwengingqi, njengedada laseYiphutha) ehla aye kutsho ngaphantsi kwemeko ephakathi yesiqhelo (oku kuqinisekiswa ziinkcukacha zexesha elidlulileyo okanye uphando lwakuqala/lwangaphambili) Amanani eentlobo ezithile / eentaka ezibalwa izihlandlo ezithathu ezilandelelanayo zehlobo okanye zobusika

UTafle 7.7: ii-EcoSpecs nee-TPCs kwichweba iGoukou

ii-EcoSpecs	i-TPC
Ikwali yamanzi	
Usasazo lobukho beetyuwa emanzini malungade lubangele ii-TPCs ezigqithileyo kwi-biota.	<ul style="list-style-type: none"> ▪ Ubukho betyuwa emanzini ngu > 0 kumphezulu wechweba. ▪ I-avareji yobukho betyuwa emanzini kuMda D ngu > 5. ▪ I-avareji yobukho betyuwa emanzini kuMda C ngu > 20. ▪ I-avareji yobukho betyuwa emanzini ukusuka kwi- 5 km kumphezulu womjelo ukusukaemlonyeni wechweba ngu > 20 ngaphezu kweenyanga ezintathu enyakeni.
Ukwahluka kweemeko (ngokwe-pH, i-oksijini enyibilikileyo nobukho bodaka) makungade kubangele ii-TPCs ezigqithileyo kwi-biota.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Nge-6.0 < yepH > ngu-7.5. ▪ i-DO ngu < 5 mg/L. ▪ Izinto eziqinileyo ezirhoxiswayo ngu > 5 mg/L (ngamanzi ahamba kancinci). <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu > 10 Nephelometric Turbidity Units (NTU) (ngamanzi ahamba kancinci). ▪ I-avareji nge- 6.0 < yepH > ngu-8.5 (isanda ngokwenyuka kweqondo lobukho beetyuwa). ▪ I-avareji ye- DO ngu < 5 mg/L.
Ubukho bezondlo ezingezozendalo (njenge-NO ₃ -N, i-NH ₃ -N ne- PO ₄ -P) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes ne-microalgae.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ i-NO_x-N ngu > 150 µg/L kwiinyanga ezimbini ezilandeelanayo. ▪ i-NH₃-N ngu > 20 µg/L kwiinyanga ezimbini ezilandeelanayo.. ▪ I-PO₄-PP ngu > 20 µg/L kwiinyanga ezimbini ezilandeelanayo.. <p>Echwebeni (ngaphandle kwamaxa kukho iimpuphuma):</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu > 150 µg/L ngomlinganiselo omnye > 200 µg/L. ▪ I-avareji ye-NH₃-N ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 100 µg/L. ▪ I-avareji ye- PO₄-P ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 50 µg/L.
Ubukho bezinto ezinyehfu (umzekelo iimethali zokuchonga, izibulala zinambuzane namayeza athile) mabungade bubangele ii-TPCs ezigqithileyo kwi-biota.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Iimethali zokuchonga (mazicokiswe futhi ziqinisekise kuhlolo lwexesha elizayo). ▪ Izibulala-vumba namayeza athile (mazicokiswe futhi ziqinisekise kuhlolo lwexesha elizayo)). <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Iqondo lobukho beentleze libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

UTafale 7.8: ii-EcoSpecs zekwaliti yamanzi nee-TPCs kwichweba iGouritz

Icandelo	ii-EcoSpecs	Imida yengxaki esenokwehla
Ikwaliiti yamanzi	<p>Usasazo lobukho beetyuwa emanzini malungade lubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<ul style="list-style-type: none"> ▪ Ubukho betyuwa emanzini ngu > 0 kumphezulu wechweba. ▪ I-avareji yobukho betyuwa emanzini kwiSikhundla 11.1 km ngu > 5. ▪ I-avareji yobukho betyuwa emanzini kuMda C ngu > 20. ▪ I-avareji yobukho betyuwa emanzini ukusuka kwi- 11 km kumphezulu womjelo ukusukaemlonyeni wechweba ngu > 20 ngaphezu kweenyanga ezintathu enyakeni. ▪ Ubukho betyuwa emanzini ngu > 40 kwiintlenge zomgobhoxo weetyuwa (oku kunxulunyaniswa nokuncipha kobumanzi nokoma kwendawo yokuphila elithafa leempuphuma)
	<p>Ukwahluka kweemeko (ngokwe-pH, i-oksijini enyibilikileyo nobukho bodaka) makungade kubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Nge-7.0 < yepH > ngu-8.3. ▪ i-DO ngu < 5 mg/L. ▪ Izinto eziqinileyo ezirhoxiswayo ngu>5 mg/L (ngamanzi ahamba kancinci). <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>10 NTU (ngamanzi ahamba kancinci, azolileyo, xa kuxube nomoya kungalinyusa iqondo lobukho bodaka liye ku- 20-40 NTU). ▪ I-avareji nge- 7.0 < yepH > ngu-8.5 (isanda ngokwenyuka kweqondo lobukho beetyuwa). ▪ I-avareji ye- DO ngu< 5 mg/L.
	<p>Ubukho bezondlo ezingezozendalo (njenge-NO₃-N, i-NH₃-N ne- PO₄-P) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes ne-microalgae.</p>	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ i-NO_x-N ngu >100 µg/L kwiinyanga ezimbini ezilandelelanayo. ▪ i-NH₃-N ngu > 20 µg/L kwiinyanga ezimbini ezilandelelanayo.. ▪ I-PO₄-PP ngu> 20 µg/L kwiinyanga ezimbini ezilandelelanayo. <p>Echwebeni (ngaphandle kwamaxa kukho iimpuphuma):</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >100 µg/L ngomlinganiselo omnye ngu > 100 µg/L. ▪ I-avareji ye- NH₃-N ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 100 µg/L. ▪ I-avareji ye- PO₄-P ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 50 µg/L.
	<p>Ubukho bezinto ezinetyhefu (umzekelo iimethali zokuchonga, izibulala zinambuzane namayeza athile) mabungade bubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Iimethali zokuchonga (ziza kuqinisekiswa) ▪ Izibulala-zinambuzane namayeza athile (ziza kuqinisekiswa)) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAf, 1995) ▪ Iqondo lobukho beentlenge libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi-Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

UTafale 7.9: ii-Eco Specs nee-TPCs kwichweba iKlein Brak

Icandelo	ii-EcoSpecs	Imida yengxaki enokwehla
Ikwali yamanzi	Usasazo lobukho beetyuwa emanzini malungade lubangele ii-TPCs ezigqithileyo kwi-biota.	<ul style="list-style-type: none"> ▪ Akukho bungqina babukho beetyuwa emanzini kumphezulu wonxweme (kwiMida D no-F) ▪ Akukho REI kumphezulu wonxweme (kwiMida D no-F) ▪ Iqondo lobukho beetyuwa emanzini ngu > 35
	Ukwahluka kweemeko (ngokwe-pH, i-oksijini enyibilikileyo nobukho bodaka) makungade kubangele ii-TPCs ezigqithileyo kwi-biota.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ I-7.0 < ye-pH > ngu- 8.5 ▪ I-DO < ngu5 mg/l ▪ Izinto eziqinileyo ezirhoxiswayo ngu > 5 mg/ l (kumanzi ahamba kancinci) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>10 NTU (ngamanzi ahamba kancinci, azolileyo, xa kuxube nomoya kungalinyusa iqondo lobukho bodaka liye ku- 20-40 NTU). ▪ I-avareji nge- 7.0 < yepH > ngu-8.5 (isanda ngokwenyuka kweqondo lobukho beetyuwa). ▪ I-avareji ye- DO ngu< 5 mg/l
	Ubukho bezondlo ezingezozendalo (njenge-NO ₃ -N, i-NH ₃ -N ne- PO ₄ -P) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes ne-microalgae.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ i-NO_x-N ngu >150 µg/L kwiinyanga ezimbini ezilandelelanayo. ▪ i-NH₃-N ngu > 20 µg/L kwiinyanga ezimbini ezilandelelanayo.. ▪ I-PO₄-PP ngu> 20 µg/L kwiinyanga ezimbini ezilandelelanayo. <p>Echwebeni (ngaphandle kwamaxa kukho iimpuphuma):</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >150 µg/L ngomlinganiselo omnye ngu > 200 µg/L. ▪ I-avareji ye- NH₃-N ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 100 µg/L. ▪ I-avareji ye- PO₄-P ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 50 µg/L.
	Ubukho bezinto ezinetyhefu (umzekelo iimethali zokuchonga, izibulala zinambuzane namayeza athile) mabungade bubangele ii-TPCs ezigqithileyo kwi-biota.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Iimethali zokuchonga (ziza kuqinisekiswa) ▪ Izibulala-zinambuzane namayeza athile (ziza kuqinisekiswa)) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Iqondo lobukho beentleze libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

UTafle 7.10: ii-EcoSpecs nee-TPCs kwi-Wilderness System

Icandelo	ii-EcoSpecs	Imida yengxaki esenokwehla
Ikwalityi yamanzi	Ubukho betyuwa emanzini	<p>Ichweba likwimo yovaleko:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho betyuwa emanzini kuMda A ngu < 12, ▪ I-avareji yobukho betyuwa emanzini kuMda B ngu: < 10 ▪ I-avareji yobukho betyuwa emanzini kuMda C ngu < 5 <p>i-avareji yobukho betyuwa emanzini kumachibi ngu- +2 ukusuka kuleyo yesiqhelo (yango2013) futhi ke ukwahluka kona akwandi, njengalapha ezantsi:</p> <ul style="list-style-type: none"> ▪ iSerpentine: 12 ± 10 ▪ iEilandvlei: 8 ± 5 ▪ iLangvlei: 10 ± 4 ▪ iRondevlei: 10 ± 5
	Ukwahluka kweemeko (ngokwe-pH, i-oksijini enyibilikileyo nobukho bodaka) makungade kubangele ii-TPCs ezigqithileyo kwi-biota.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ I-6.0 < ye-pH > ngu- 7.0 (kwiTouw) ▪ I-7.0 < ye-pH > ngu- 8.0 (kwiDuiwe) ▪ I-DO < ngu5 mg/l ▪ Izinto eziqinileyo ezirhoxiswayo ngu > 5 mg/ l (kumanzi ahamba kancinci) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>5 NTU (ngamanzi ahamba kancinci. ▪ I-avareji nge- 6.0 < yepH > ngu-8.5 (isanda ngokwenyuka kweqondo lobukho beetyuwa). ▪ I-avareji ye- DO ngu< 5 mg/l ▪ DO < 5 mg/l <p>Kumachibi:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>5 NTU ▪ I-avareji nge7.0 < yepH > ngu> 8.5 ▪ I-avareji ye- DO ngu< 5 mg/l
	Ubukho bezondlo ezingezozendalo (njenge- NO ₃ -N, i-NH ₃ -N ne- PO ₄ -P) mabungade bubangele ii-TPCs ezigqithileyo kwi-macrophytes ne-microalgae.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ i-NO_x-N ngu >50 µg/L kwiinyanga ezimbini ezilandelelanayo. ▪ i-NH₃-N ngu > 10 µg/L kwiinyanga ezimbini ezilandelelanayo.. ▪ I-PO₄-PP ngu> 10 µg/L kwiinyanga ezimbini ezilandelelanayo. <p>Echwebeni (ngaphandle kwamaxa kukho iimpuphuma):</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >50 µg/L ngomlinganiselo omnye ngu > 100 µg/L. ▪ I-avareji ye-NH₃-N ngu > 10 µg/L ngethuba lophando, ngomlinganiselo omnye > 100 µg/L. ▪ I-avareji ye- PO₄-P ngu > 10 µg/L ngethuba lophando, ngomlinganiselo omnye > 50 µg/L. <p>Kumachibi:</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >50 µg/L ngethuba lophando, ngomlinganiselo omnye ngu > 100 µg/L. ▪ I-avareji ye-NH₃-N ngu > 20 µg/l ngethuba lophando (iza kuqinisekiswa) ▪ I-avareji ye- PO₄-P ngu> 20 µg/l ngethuba lophando (iza kuqinisekiswa)
	Ubukho bezinto ezinyehfu (umzekelo iimethali zokuchonga, izibulala zinambuzane	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Iimethali zokuchonga (iza kuqinisekiswa)

	namayeza athile) mabungade bubangele ii-TPCs ezigqithileyo kwi-biota.	<ul style="list-style-type: none"> ▪ Izibulala-zinambuzane namayeza athile (ziza kuqinisekiswa) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Iqondo lobukho beentlenge libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
--	---	---

UTafle 7.11: Uvavanyo lwesimo sangoku sekwaliti yamanzi kwi-H8DUIW-EWR1

IZithako zeKwaliti yaManzi	Ixabiso le-PES	IBakala/Amagqabantshintshi
Ii-ioni zeetyuwa ezingezozendalo (mg/l)		
Sulphate as SO ₄	N/A	-
Sodium as Na	382.2	Ibetha ngaphaya ko ≤ 70 mg/L (TWQR) osetyenziswa kwezolimo: kunkcukeshelo.
Magnesium as Mg	67.4	Asikho isikhokelo.
Calcium as Ca	55.0	Asikho isikhokelo.
Chloride as Cl	805.4	Ibetha ngaphaya ko ≤ 100 mg/L (TWQR) osetyenziswa kwezolimo: kunkcukeshelo.
Potassium as K	9.25	Asikho isikhokelo.
Ukutsala umbane (mS/m)		
	272	E/F: RC = 80 mS/m.
Izondlo (mg/l)		
SRP	0.014	A
TIN	0.118	A
Ukwahluka kwendalo		
pH (5 th + 95 th %ile)	u-6.6 no- 8.1	B
Ubushushu (°C)	N/A	A/B. likho ifuthe elilindelekileyo kumanzi ahamba kancinci.
i-oksijini enyibilikisiweyo (mg/L)	N/A	A/B. likho ifuthe elilindelekileyo kumanzi ahamba kancinci.
Ubukho bodaka (NTU)	N/A	B. iinguqu kwinqondo lobukho bodaka zinxulunyaniswa ikakhulu nezendalo, apho kuye kubekho neenguqulelo ezenziwa ngabantu ezingephi - umzekelo olwaa shishino ngegraveli phaya emantla omjelo
Ukwahluka ngokwemvakalelo		
Chl-a: phytoplankton (ug/L)	N/A	N/A
Inqaku lobukhulu bezinto eziphilayo ezingenamathatho (i-MIRAI)	50.7%	D
Inqaku le-SASS	78	
Inqaku le-ASPT	56	
ii-Diatoms	11.1	C/D (n = 1, Jan 2014)
Inqaku leentlanzi (i-FRAI)	51.6%	D (izonke ezo zechweba ezafudukela kumanzi ahlaziyekileyo nezo ke ingezizo zomthonyama).
Iityhefu		
Ammonia (as N)	0.003	A
Fluoride (as F)	0.33	A
LULONKE UHLELO LWESIKHUNDLA		C (73.2%)

Izithako zeKwaliti yaManzi	Ixabiso le-PES	IBakala/Amagqabantshintshi
(Umzekelo we-PAI)		

(a) N/A- bezingekho iinkcukacha kolu vavanyo.

UTafle 7.12: H8DUIW-EWR1: ii-EcoSpecs nee- TPCs zekwaliti yamanzi (iBakala C)

ii-Metrics	ii-EcoSpecs	ii-TPCs
ii-ioni zeetyuwa ezingezozendalo		
Sulphate as SO ₄	N/A	N/A
Sodium as Na	i-95 th percentile yeenkukacha mayibe ngu- ≤ 380 mg/L.	i-95 th percentile yeenkukacha mayibe ngu- 300 - 380 mg/L.
Magnesium as Mg	i-95 th percentile yeenkukacha mayibe ngu- ≤ 67 mg/L.	i-95 th percentile yeenkukacha mayibe ngu- 53.5 - 67 mg/L.
Calcium as Ca	i- 95 th percentile yeenkukacha mayibe ngu- ≤ 55 mg/L.	i- 95 th percentile yeenkukacha mayibe ngu- 44 - 55 mg/L.
Chloride as Cl	i- 95 th percentile yeenkukacha mayibe ngu- ≤ 800 mg/L.	i-95 th percentile yeenkukacha mayibe ngu- 640 - 800 mg/L.
Potassium as K	i-95 th percentile yeenkukacha mayibe ngu- ≤ 9 mg/L.	i- 95 th percentile yeenkukacha mayibe ngu- 7 - 9 mg/L.
Ukwahluka kwendalo		
Ukutsala umbane (mS/m)	i- 95 th percentile yeenkukacha mayibe ngu- ≤ 270 mS/m.	i-95 th percentile yeenkukacha mayibe ngu- 210 - 270 mS/m.
ipH	i- 5 th percentile yeenkukacha mayibe ngu- 6.5 - 8.0, ze i- 95 th percentile ibe ngu-8.0 - 8.8.	i- 5 th percentile yeenkukacha - ≤ 6.3 ze i-95 th percentile ibe ngu- ≥ 8.6.
ubushushu ^(a)	Uluhlu lobushushu bendalo.	Qalisa uhlolo lwesiqhelo loku kwahluka.
i-oksijini enyibilikisiweyo ^(a) (DO)	i-5 th percentile yeenkukacha mayibe ngu- ≥ 7.0 mg/L.	i-5 th percentile yeenkukacha mayibe ngu- 7.2 - 7.0 mg/L. Qalisa uhlolo lwesiqhelo loku kwahluka.
Ubukho bodaka ^(a)	Iinguqu kwiqondo lobukho bodaka bunxulunyaniswa ikakhulu neenguqulelo ezenziwa ngabantu ezingephi (umzekelo olwaa shishino lwegraveli kumantla omjelo gravel mining upstream). Iintenge ziiindelekile kwiindawo zokuphila.	Qalisa uhlolo lwesiqhelo loku kwahluka.
Izondlo		
TIN	i-50 th percentile yeenkukacha mayibe ngu ≤ 0.25 mg/L.	i- 50 th percentile yeenkukacha mayibe ngu 0.2 - 0.25 mg/L.
PO ₄ -P	i-50 th percentile yeenkukacha mayibe ngu ≤ 0.015 mg/L.	i-50 th percentile yeenkukacha mayibe ngu 0.012 - 0.015 mg/L.
Ukwahluka ngokwemvakalelo		
Chl-a phytoplankton	i-50 th percentile yeenkukacha mayibe ngu < 15 µg/L.	i- 50 th percentile yeenkukacha mayibe ngu 12 - 15 µg/L.
Chl-a periphyton	i-50 th percentile yeenkukacha mayibe ngu ≤ 12 mg/m ² .	i-50 th percentile yeenkukacha mayibe ngu 10 - 12 mg/m ² .
Iityhefu		
Fluoride	i-50 th yeenkukacha mayibe ngu ≤ 1.5 mg/L.	i-50 th percentile yeenkukacha mayibe ngu 1.2 - 1.5 mg/L.
Ammonia (NH ₃ -N)	i-50 th percentile yeenkukacha mayibe ngu ≤ 0.015 mg/L.	i- 50 th percentile yeenkukacha mayibe ngu 0.012 - 0.015 mg/L.
Other toxics	i- 95 th percentile yeenkukacha mayibe phakathi koluhlu lwekwaliti yamanzi	Likho ifuthe eliiindelekileyo ukuba i- 95 th percentile yeenkukacha ibetha ngaphaya

ii-Metrics	ii-EcoSpecs	ii-TPCs
	engqaliweyo (i-Target Water Quality Range (TWQR) njengoko kubhaliwe kuDWAF (1996) okanye umda weBakala A njengoko kubhaliwe kuDWAF (2008).	kwe- TWQR njengoko kubhaliwe kuDWAF (1996) okanye kumda ophezulu weBakala A njengoko kubhaliwe kuDWAF (2008).

(a) N/A- Bezingekho iinkcukacha kolu vavanyo

UTafle 7.13: ii-EcoSpecs nee-TPCs kwichweba iDuiwenhoks

Icandelo	ii-EcoSpecs	Imida yengxaki esenokwehla
Ikwalityi yamanzi	Usasazo lobukho beetyuwa emanzini malungade lubangele ii-TPCs ezigqithileyo kwi-biota.	<ul style="list-style-type: none"> ▪ Ubukho betyuwa emanzini ngu > 0 kumphezulu wechweba. ▪ I-avareji yobukho betyuwa emanzini kuMda D ngu > 5 ▪ I-avareji yobukho betyuwa emanzini kuMda C ngu > 20. ▪ I-avareji yobukho betyuwa emanzini ukusuka kwi- 5 km kumphezulu womjelo ukusuka emlonyeni wechweba ngu > 20 ngaphezu kweenyanga ezintathu enyakeni.
	Ukwahluka kweemeko (ngokwe-pH, i-oksijini enyibilikileyo nobukho bodaka) makungade kubangele ii-TPCs ezigqithileyo kwi-biota.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ I-6.0 < ye-pH > ngu- 7.5 ▪ I-DO < ngu5 mg/l ▪ Izinto eziqinileyo ezirhoxiswayo ngu > 5 mg/l (kumanzi ahamba kancinci) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>10 NTU (ngamanzi ahamba kancinci. ▪ I-avareji nge- 6.0 < yepH > ngu-8.5 (isanda ngokwenyuka kweqondo lobukho beetyuwa). ▪ I-avareji ye- DO ngu< 5 mg/l
	Ubukho bezondlo ezingezozendalo (njenge-NO ₃ -N, i-NH ₃ -N ne- PO ₄ -P) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes ne-microalgae.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ i-NO_x-N ngu >150 µg/L kwiinyanga ezimbini ezilandelelanayo. ▪ i-NH₃-N ngu > 20 µg/L kwiinyanga ezimbini ezilandelelanayo.. ▪ I-PO₄-PP ngu> 20 µg/L kwiinyanga ezimbini ezilandelelanayo. <p>Echwebeni (ngaphandle kwamaxa kukho iimpuphuma):</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >150 µg/L ngomlinganiselo omnye ngu > 200 µg/L. ▪ I-avareji ye-NH₃-N ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 100 µg/L. ▪ I-avareji ye- PO₄-P ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 50 µg/L.

	<p>Ubukho bezinto ezinyehfu (umzekelo iimethali zokuchonga, izibulala zinambuzane namayeza athile) mabungade bubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Iimethali zokuchonga (ziza kuqinisekiswa) ▪ Izibulala-zinambuzane namayeza athile (ziza kuqinisekiswa) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Iqondo lobukho beentlele libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
--	---	---

UTafle 7.14: ii-EcoSpecs nee-TPCs kwichweba iGoukou

ii-EcoSpecs	i-TPC
Ikwaliti yamanzi	
<p>Usasazo lobukho beetyuwa emanzini malungade lubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<ul style="list-style-type: none"> ▪ Ubukho betyuwa emanzini ngu > 0 kumphezulu wechweba. ▪ I-avareji yobukho betyuwa emanzini kuMda D ngu > 5 ▪ I-avareji yobukho betyuwa emanzini kuMda C ngu > 20. ▪ I-avareji yobukho betyuwa emanzini ukusuka kwi- 5 km kumphezulu womjelo ukusuka emlonjeni wechweba ngu > 20 ngaphezu kweenyanga ezintathu enyakeni.
<p>Ukwahluka kweemeko (ngokwe-pH, i-oksijini enyibilikileyo nobukho bodaka) makungade kubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ I-6.0 < ye-pH > ngu- 7.5 ▪ I-DO < ngu5 mg/l ▪ Izinto eziqinileyo ezirhoxiswayo ngu > 5 mg/ l (kumanzi ahamba kancinci) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>10 NTU (ngamanzi ahamba kancinci). ▪ I-avareji nge- 6.0 < yepH > ngu-8.5 (isanda ngokwenyuka kweqondo lobukho beetyuwa). ▪ I-avareji ye- DO ngu< 5 mg/l
<p>Ubukho bezondlo ezingezozendalo (njenge- NO₃-N, i-NH₃-N ne- PO₄-P) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes ne-microalgae.</p>	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ i-NO_x-N ngu >150 µg/L kwiinyanga ezimbini ezilandelelanayo. ▪ i-NH₃-N ngu > 20 µg/L kwiinyanga ezimbini ezilandelelanayo.. ▪ I-PO₄-PP ngu> 20 µg/L kwiinyanga ezimbini ezilandelelanayo. <p>Echwebeni (ngaphandle kwamaxa kukho iimpuphuma):</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >150 µg/L ngomlinganiselo omnye ngu > 200 µg/L. ▪ I-avareji ye-NH₃-N ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 100 µg/L. ▪ I-avareji ye- PO₄-P ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 50 µg/L.

ii-EcoSpecs	i-TPC
<p>Ubukho bezinto ezinyehfu (umzekelo iimethali zokuchonga, izibulala zinambuzane namayeza athile) mabungade bubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Iimethali zokuchonga (mazicokiswe futhi ziqinisekiswe kuhlo lwexesha elizayo). ▪ Izibulala-vumba namayeza athile (mazicokiswe futhi ziqinisekiswe kuhlo lwexesha elizayo)). <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAf, 1995) ▪ Iqondo lobukho beentlenge libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

UTafle 7.15: ii- EcoSpecs zekwaliti yamanzi nee-TPCs kwichweba iGouritz

Icandelo	ii-EcoSpecs	Imida yengxaki esenokwehla
Ikwality yamanzi	<p>Usasazo lobukho beetyuwa emanzini malungade lubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<ul style="list-style-type: none"> ▪ Ubukho betyuwa emanzini ngu > 0 kumphezulu wechweba. ▪ I-avareji yobukho betyuwa emanzini kwiSikhundla 11, 1 km kumphezulu womjelo webhulorho ngu > 5 ▪ I-avareji yobukho betyuwa emanzini kuMda C ngu > 20. ▪ I-avareji yobukho betyuwa emanzini ukusuka kwi- 11 km kumphezulu womjelo ukusuka emlonyeni wechweba ngu > 20 ngaphezu kweenyanga ezintathu enyakeni. ▪ Iqondo lobukho betyuwa emanzini ngu > 40 kwiintlenge zomgobhozo weetyuwa (linxulunyaniswa nokuncipha kobumanzi nokoma kwendawo yokuphila kwithafa leempophuma.
	<p>Ukwahluka kweemeko (ngokwe-pH, i-oksijini enyibilikileyo nobukho bodaka) makungade kubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ I-7.0 < ye-pH > ngu- 8.3 ▪ I-DO < ngu5 mg/l ▪ Izinto eziqinileyo ezirhoxiswayo ngu > 5 mg/ l (kumanzi ahamba kancinci) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>10 NTU (ngamanzi ahamba kancinci, azolileyo, xa kuxube nomoya kungalinyusa iqondo lobukho bodaka liye ku- 20-40 NTU). ▪ I-avareji nge- 7.0 < yepH > ngu-8.5 (isanda ngokwenyuka kweqondo lobukho beetyuwa). ▪ I-avareji ye- DO ngu< 5 mg/l

Icandelo	ii-EcoSpecs	Imida yengxaki esenokwehla
	Ubukho bezondlo ezingezozendalo (njenge-NO ₃ -N, i-NH ₃ -N ne- PO ₄ -P) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes ne-microalgae.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ i-NO_x-N ngu >100 µg/L kwiinyanga ezimbini ezilandelelanayo. ▪ i-NH₃-N ngu > 20 µg/L kwiinyanga ezimbini ezilandelelanayo.. ▪ I-PO₄-PP ngu> 20 µg/L kwiinyanga ezimbini ezilandelelanayo. <p>Echwebeni (ngaphandle kwamaxa kukho iimpuphuma):</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >100 µg/L ngomlinganiselo omnye ngu > 150 µg/L. ▪ I-avareji ye-NH₃-N ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 100 µg/L. ▪ I-avareji ye- PO₄-P ngu > 20 µg/L ngethuba lophando, ngomlinganiselo omnye > 50 µg/L.
	Ubukho bezinto ezinetyhefu (umzekelo iimethali zokuchonga, izibulala zinambuzane namayeza athile) mabungade bubangele ii-TPCs ezigqithileyo kwi-biota.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Iimethali zokuchonga (ziza kuqinisekiswa) ▪ Izibulala-zinambuzane namayeza athile (ziza kuqinisekiswa)) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAf, 1995) ▪ Iqondo lobukho beentlenga libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)

UTafale 7.16: ii-EcoSpecs nee- TPCs kwi- Wilderness System

Icandelo	ii-EcoSpecs	Imida yengxaki esenokwehla
Ikwali yamanzi	Ubukho beetyuwa emanzini	<p>Ichweba likwimo yovaleko:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho betyuwa emanzini kuMda A ngu < 12, ▪ I-avareji yobukho betyuwa emanzini kuMda B ngu: < 10 ▪ I-avareji yobukho betyuwa emanzini kuMda C ngu < 5 <p>i-avareji yobukho betyuwa emanzini kumachibi ngu- +2 ukusuka kuleyo yesiqhelo (yango2013) futhi ke ukwahluka kona akwandi, njengalapha ezantsi:</p> <ul style="list-style-type: none"> ▪ iSerpentine: 12 ± 10 ▪ iEilandvlei: 8 ± 5 ▪ iLangvlei: 10 ± 4 ▪ iRondevlei: 10 ± 5
	Ukwahluka kweemeko (ngokwe-pH, i-oksijini enyibilikileyo nobukho bodaka) makungade kubangele ii-TPCs ezigqithileyo kwi-biota.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ I-6.0 < ye-pH > ngu- 7.0 (kwiTouw) ▪ I-7.0 < ye-pH > ngu- 8.0 (kwiDuiwe) ▪ I-DO < ngu5 mg/l ▪ Izinto eziqinileyo ezirhoxiswayo ngu > 5 mg/ l (kumanzi ahamba kancinci) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>5 NTU (ngamanzi ahamba kancinci. ▪ I-avareji nge- 6.0 < yepH > ngu-8.5 (isanda ngokwenyuka kweqondo lobukho beetyuwa). ▪ I-avareji ye- DO ngu< 5 mg/l ▪ DO < 5 mg/l <p>Kumachibi:</p> <ul style="list-style-type: none"> ▪ I-avareji yobukho bodaka ngu>5 NTU ▪ I-avareji nge7.0 < yepH > ngu> 8.5 ▪ I-avareji ye- DO ngu< 5 mg/l
	Ubukho bezondlo ezingezozendalo (njenge-NO ₃ -N, i-NH ₃ -N ne- PO ₄ -P) mabungade bubangele ii-TPCs ezigqithileyo kwii-macrophytes ne-microalgae.	<p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ i-NO_x-N ngu >50 µg/L kwiinyanga ezimbini ezilandelelanayo. ▪ i-NH₃-N ngu > 10 µg/L kwiinyanga ezimbini ezilandelelanayo.. ▪ I-PO₄-PP ngu> 10 µg/L kwiinyanga ezimbini ezilandelelanayo. <p>Echwebeni (ngaphandle kwamaxa kukho iimpuphuma):</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >50 µg/L ngomlinganiselo omnye ngu > 100 µg/L. ▪ I-avareji ye-NH₃-N ngu > 10 µg/L ngethuba lophando, ngomlinganiselo omnye > 100 µg/L. ▪ I-avareji ye- PO₄-P ngu > 10 µg/L ngethuba lophando, ngomlinganiselo omnye > 50 µg/L. <p>Kumachibi:</p> <ul style="list-style-type: none"> ▪ I-avareji ye- NO_x-N ngu >50 µg/L ngethuba lophando, ngomlinganiselo omnye ngu > 100 µg/L. ▪ I-avareji ye-NH₃-N ngu > 20 µg/l ngethuba lophando (iza kuqinisekiswa)


	<p>Ubukho bezinto ezinyehfu (umzekelo iimethali zokuchonga, izibulala zinambuzane namayeza athile) mabungade bubangele ii-TPCs ezigqithileyo kwi-biota.</p>	<ul style="list-style-type: none"> ▪ I-avareji ye- PO₄-P ngu> 20 µg/l ngethuba lophando (iza kuqinisekiswa) <p>Amanzi angena emlanjeni:</p> <ul style="list-style-type: none"> ▪ Iimethali zokuchonga (ziza kuqinisekiswa) ▪ Izibulala-zinambuzane namayeza athile (ziza kuqinisekiswa) <p>Echwebeni:</p> <ul style="list-style-type: none"> ▪ Iqondo lobukho bamanzi kweli candelo libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi- SA Water Quality Guidelines for coastal marine waters (DWAF, 1995) ▪ Iqondo lobukho beentlenge libetha ngaphaya kwamaxabiso angqaliweyo ngokubhaliweyo kwi-Indian Ocean (WIO) Region guidelines (UNEP/Nairobi Convention Secretariat and CSIR, 2009)
--	---	--

8. IMIWONYO

UTafale 8.1: Imiwonyo – Umqathango wokhuselo lwendawo yoboniselo ngamanzi yeminyaka emine

Indawo yoboniselo ngamanzi yeminyaka emine	i-EIS	i-PES	i-REC	Ungayizuza njani i- REC
K10A	Phakathi	C	C	Beka iliso kutyani oluzizityalo ezingezizo zomthonyama nezitshabalalisayo, kukhukuliso lomhlaba nezinye izinto ezingenelelayo kusetyenziso lomhlaba.
K10B	Phakathi	C	C	
K10C	Phakathi	B/C	B/C	
K10D	Phakathi	B/C	B/C	
K10E	Phakathi	B/C	B/C	
K10F	Phakathi	C	C	
K20A	Phakathi	C	C	
K30A	Phezulu	C	C	Kubekho izidambisi kwimimandla yeedolophu neyezolimo; beka iliso kwikwaliti yamanzi, kukhukuliso lomhlaba nakutyani olutshabalalisayo.
K30B	Phezulu	D	C/D	
K30C	Phakathi	D	D	Beka iliso kutyani oluzizityalo ezingezizo zomthonyama nezitshabalalisayo, kukhukuliso lomhlaba nezinye izinto ezingenelelayo kusetyenziso lomhlaba.
K30D	Phezulu kakhulu	B	B	
K40A	Phakathi	D	D	
K40B	Phakathi	C	C	
K40C	Phakathi	C	C	
K40D	Phezulu kakhulu	B	B	
K40E	Phakathi	B/C	B/C	
K50A	Phakathi	B/C	B/C	Khusela uphucule imeko yaloo mabala emiwonyo , beka iliso kutyani olutshabalalisayo.
K50B	Phezulu	C/D	C	
K60A	Phakathi	B	B	Beka iliso kutyani oluzizityalo ezingezizo zomthonyama nezitshabalalisayo, kukhukuliso lomhlaba nezinye izinto ezingenelelayo kusetyenziso lomhlaba.
K60B	Phakathi	B	B	
K60C	Phakathi	B	B	
K60D	Phezulu	A	A	
K60E	Phezulu	C	C	
K60F	Phezulu	C	C	
K60G	Phakathi	C	C	
K70A	Phakathi	C	C	
K70B	Phantsi	A	A	
H80A	Phezulu	C/D	C	
H80B	Phakathi	C	C	
H80C	Phakathi	D	D	
H80D	Phakathi	D	D	

Indawo yoboniselo ngamanzi yeminyaka emine	i-EIS	i-PES	i-REC	Ungayizuza njani i- REC
H80E	Phakathi	C/D	C/D	
H90A	Phakathi	C	C	
H90B	Phakathi	D	D	
H90C	Phakathi	D	D	
H90D	Phakathi	C	C	
H90E	Phakathi	C/D	C/D	
J11D	Phantsi	C	C	
J11F	Phakathi	C	C	
J11G	Phakathi	B	B	
J12A	Phakathi	B	B	
J12B	Phakathi	B	B	
J12J	Phakathi	B	B	
J12K	Phakathi	B	B	
J12L	Phakathi	C	C	
J21A	Phakathi	B/C	B/C	
J21B	Phakathi	B	B	
J22B	Phakathi	B	B	
J22G	Phakathi	B	B	Control invasive alien vegetation, erosion and land-use encroachment.
J22K	Phantsi	B/C	B/C	
J23E	Phantsi	C	C	
J23J	Phakathi	B	B	
J24F	Phantsi	C	C	
J25A	Phantsi	B	B	
J33B	Phantsi	C	C	
J33E	Phantsi	C	C	
J34C	Phantsi	C	C	
J34D	Phantsi	C	C	
J34E	Phantsi	C/D	C/D	
J34F	Phantsi	D	D	
J40B	Phantsi	B	B	
J40C	Phakathi	C/D	C/D	
J40D	Phakathi	D	D	
J40E	Phezulu	C	C	



Umzobo 1: iindawo ngeendawo zengingqi i- Gouritz, imilambo/iindawo adibana kuzo amachweba nezikhundla ze-EWR sites



Umzobo 2: iindawo ngeendawo zengingqi i-Breede-Overberg, imilambo/iindawo adibana kuzo amachweba nezikhundla ze-EWR sites